

PM 04 16-12

3/24/96

P9174



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other

OPP Identifier Number 229746

Application for Pesticide - Section I

1. Company/Product Number 16-12... 2. EPA Product Manager Edwards... 3. Proposed Classification [X] None [] Restricted... 4. Company/Product Name BLUE DRAGON Garden Dust... 5. Name and Address of Applicant... 6. Expedited Review...

Section - II

Amendment - Explain below. Resubmission in response to Agency letter dated... Notification - Explain below. Final printed labels in response to Agency letter dated... "Me Too" Application. Other - Explain below.

NOTIFICATION

Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification to add the following section headings to label: Precautionary Statements Hazards to Humans & Domestic Animals Directions for Use

MAR 24 1996

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging... 2. Type of Container: Metal, Plastic, Glass, Paper, Other... 3. Location of Net Contents Information... 4. Size(s) Retail Container... 5. Location of Label Directions... 6. Manner in Which Label is Affixed to Product...

Section - IV

1. Contact Point: Name Rebecca M. Horton, Title Regulatory & Sales Support Manager, Telephone No. (540)362-3657... Certification: I certify that the statements I have made on this form... 2. Signature: Rebecca M. Horton... 3. Title: Regulatory & Sales Support Manager... 4. Typed Name: Rebecca M. Horton... 5. Date: 2/28/96... 6. Date Application Received: (Stamped) SIG 6/3/96

AFTER 2/4

[FRONT PANEL]

**DRAGON®
BLUE DRAGON® GARDEN DUST**

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects

Contains SEVIN® brand carbaryl insecticide

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 2%

INERT INGREDIENTS: 98%

TOTAL: 100%

Keep Out of Reach of Children

CAUTION

See back/side panels for additional cautions.

NET WEIGHT 4 LB

[BACK/SIDE/OTHER PANEL(S)]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

FOR HOMEOWNER USE ONLY. BLUE DRAGON® GARDEN DUST controls certain listed insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces, if needed, repeat 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1 ¼ lb. per 1,000 sq. ft. For control of Lygus, Stink Bugs, Corn Earworm, Tarnish Plant Bug, apply approximately 1 ¼ lb. per 1,000 sq. ft. Do not feed treated bean vines to domestic animals.

CORN - Use approximately 1 ¼ lb. to 2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation.

Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS - For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, apply approximately 1 1/4 lb. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew) - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH - For control of Flea Beetles, Leafhoppers, Harlequin Bug, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For six-spotted Leafhoppers, Corn Earworm, Armyworm and Stink Bugs, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce - 3 days).

ASPARAGUS - For control of Asparagus Beetle on spears, apply approximately 1 1/4 to 2 1/2 lbs. per 1,000 sq. ft. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

* **CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS** - For control of Flea Beetles, Harlequin Bugs, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For Corn Earworm, Stink Bug, Armyworm, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEAS AND OTHER ORNAMENTALS - For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillar, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

DOGS, CATS - For control of Fleas, Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens or puppies under 4 weeks old. Do not use on pregnant dogs.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. DO NOT take internally. Wear regular long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

4/4

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

