



CAUTION

GARMENT, IF INHALED OR SWALLOWED DO NOT breathe dust. DO NOT take or eat, drink. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Contact your doctor, recheck cholesterol and visit your doctor as advised. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wet lands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

STORAGE & DISPOSAL

Storage: Store only in original container in a cool, dry place inaccessible to children and pets and away from feed and feedstuffs. **Pesticide disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container disposal:** Do not reuse empty bag. Discard bag in trash.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SUPER BLUE DRAGON[®] GARDEN DUST

5% SEVIN[®]

KILLS: Japanese Beetles, Mexican Bean Beetles,
Flea Beetles, Tomato Hornworm, Corn Earworm,
Fleas, Brown Dog Ticks,
and certain other listed insects.

ACTIVE INGREDIENT	BY WEIGHT
*Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional cautions

NET WEIGHT 4 LB.



SUPER BLUE DRAGON 5% SEVIN DUST

SEVIN is a registered trademark of Ciba-Geigy Corporation, Basle, Switzerland.

ACCEPTED

JUL 1 1 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 16-15

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

SUPER BLUE DRAGON is a dust which is effective against a wide range of insect pests of plants and animals. It is effective against many of the most common insects which cause damage to crops and livestock. It is also effective against many of the most common insects which cause damage to animals. It is also effective against many of the most common insects which cause damage to humans. It is also effective against many of the most common insects which cause damage to property.

DIRECTIONS

It is a violation of Federal law to apply this product to any animal which is not listed with a labeling statement on the labeling of food and feed products. Do not apply this product to crops on loss or other animal labels.

- ✓ **BEANS** - For control of Mexican bean beetle, leaf miner, leaf beetle, apply approximately 1/2 lb./1000 sq. ft. DO NOT treat beans before they are ready for harvest.
- ✓ **LETTUCE** - For control of pyralid, apply approximately 1 lb./1000 sq. ft. (DO NOT apply to lettuce within 3 days of harvest or within 14 days of leaf lettuce).
- ✓ **CORN** - Use approximately 3/4 to 1000 sq. ft. for control of corn earworm, fall armyworm, flea beetles, Japanese beetles, Sap beetle, leafhopper, and European cornborer. For corn borer and Japanese beetles (*bothworm damage*) and for *leaf beetles*, make one or more applications to entire plant beginning when borer starts to enter and depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper treatment treatments. For borer or leafhopper, apply *from early in the week when plants are about knee high*. Repeat at weekly intervals until the plants are ready for harvest. For insects other than borer and leafhopper, apply 1/2 to 3/4 lb. per acre starting when leaf flies appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Do apply immediately after rain. Allow 7 days before harvest or grazing of the crop. Application of this product to the foliage of corn during the pollen shed period will reduce bee populations.
- ✓ **POTATOES, TOMATOES, EGGPLANT AND PEPPERS** - For control of Colorado potato beetle, leafhoppers, and flea beetle, apply approximately 1/2 to 1000 sq. ft. For control of European cornborer, fall armyworm, tomato leaf worm, European cornborer, flea beetle and leafhopper, apply approximately 3/4 lb./1000 sq. ft.
- ✓ **CUCUMBERS, SUMMER SQUASH** - For control of pickleworm, melonworm, cucumber beetle, squash bugs, flea beetles, apply approximately 1/2 lb./1000 sq. ft.
- ✓ **GRAPES** - For control of leafhoppers, Japanese beetle, grape berry moth, apply approximately 1 lb./1000 sq. ft.
- ✓ **STRAWBERRIES** - For control of meadow spittlebug, leaf roller, apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.
- ✓ **ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS** - For control of Japanese beetle, periodical cicada, aphid, tent caterpillar, lace bugs, boxwood leaf miner, apply approximately 1-1 1/4 lbs/1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.
- ✓ **DOGS, CATS** - For control of fleas and brown dog ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. DO NOT treat kittens and puppies under 4 weeks old. DO NOT apply to animals more than once weekly. Do not use on pregnant dogs.

EPA Reg. No. 16-15

EPA Est. 16 VA 1

Manufactured By
DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24019