

PM-19

Reg # 16-15

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

SS15149

19 14
200
17

FEB 24 1997

Rebecca M. Horton
Dragon Corporation
P.O. Box 7311
Roanoke, VA 24019-7311

Subject: Super Blue Dragon Garden Dust
EPA Registration No. 16-15
Amendment dated November 12, 1996
Response to Agency letter dated 10/1/96

Dear Ms. Horton:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:

a) Add a FIRST AID section (or STATEMENT OF PRACTICAL TREATMENT) to your label as follows:

"FIRST AID

IF SWALLOWED: Induce vomiting by touching back of throat with finger and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes with plenty of water. Wash skin thoroughly with soap and water.

2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

151

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

		CONCURRENCES						
SYMBOL	Enclosure							
SURNAME								
DATE								

Edwards A-114/15.1 2/16/97

29 4

ACCEPTED
with COMMENTS
in EPA Letter Dated

[FRONT PANEL]

SUPER BLUE DRAGON® GARDEN DUST

5% SEVIN®

FEB 24 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

16-15

Nothing Beats The Dragon!

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm, Fleas, Brown Dog Ticks and certain other listed insects.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm, Fleas and Brown dog ticks.]

ACTIVE INGREDIENT:	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS:	<u>95%</u>
TOTAL:	100%

Keep Out of Reach of Children
CAUTION
See back/side panels for additional cautions.

NET WT. 4 LB

[BACK/SIDE/OTHER PANEL(S)]
READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

REC'D
NOV 20 1996
P 3 32 PM
DPD1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. **DO NOT** take internally. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.** [NOTE to EPA: When marketed for use on dogs and cats, the following statement will be included on the label: Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.]

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

NOTE TO PHYSICIAN [Note to EPA: When marketed for use on dogs and cats, the following will be included on the label: & **VETERINARIAN**]: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOMEOWNER USE ONLY. SUPER BLUE DRAGON® controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using **DRAGON® SEVIN® Liquid**. Do not repeat application more than once a week. Do not use with lime or other alkaline materials. **SUPER BLUE DRAGON®** may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles - apply approximately 1/2 lb. per 1,000 sq. ft. Do not feed treated bean vines to domestic animals.

LETTUCE - For control of lygus and stink bugs, apply approximately 1 lb. per 1,000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN - Use approximately 3/4 lb. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, Leafhoppers and European corn borer. For corn borers and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS - For control of Colorado Potato Beetle, Leafhoppers and Flea Beetles, apply approximately 1/2 lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bug and Lygus Bugs, apply approximately 3/4 lb. per 1,000 sq. ft.

CUCUMBERS, SUMMER SQUASH - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, apply approximately 1/2 lb. per 1,000 sq. ft.

49 4

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, apply approximately 1 lb. per 1,000 sq. ft.

STRAWBERRIES - For control of Meadow Spittle Bug, Leaf Roller, apply approximately 1 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS - For control of Japanese Beetle, Periodical Cicada, Aphid, Tent Caterpillar, Lace Bugs, Boxwood Leaf Miner, apply approximately 1-1 1/4 lbs. per 1,000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.

DOGS, CATS - READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS OR CATS.
To control (kill) Fleas and Brown Dog Ticks: Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pets eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Do not repeat treatment for 1 week. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals. Dust bedding and doghouse using a shaker can or other dust applicator.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.