

16-27

4-3-1971

FRONT PANEL

1/5

Best Available Copy

DRAGON
5%
SEVIN* DUST

NET WEIGHT

FOR CONTROL OF JAPANESE BEETLE, FLEA BEETLE, MEXICAN LEAF BEETLE, TOMATO PINworm,
ORN EARWORM, AND CERTAIN OTHER INSECTS ATTACKING VEGETABLES, GRAPES, ORNAMENTALS,
AND SHRUBS.

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate).....	5%
INERT INGREDIENTS.....	95%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional cautions.

Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

MANUFACTURED BY
DRAGON CHEMICAL CORPORATION
ROANOKE, VIRGINIA 24019

EPA REG. NO. 16-27

DRAGON
5
SEVIN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON 5 SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

DIRECTIONS

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, apply approximately $\frac{1}{2}$ lb./1000 sq. ft. DO NOT feed treated bean vines to dairy or pet animals.

LETTUCE - For control of Lygus and Stink Bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 10 days to leaf lettuce.

CORN - For control of Corn Earworm, apply approximately $\frac{3}{4}$ to $\frac{9}{10}$ lb./1000 sq. ft. Make 3-4 applications at 2 to 3 day intervals beginning when silks first appear. For control of European Corn Borer, apply approximately $\frac{3}{4}$ to $\frac{9}{10}$ lb./1000 sq. ft. Make application into the whorls and to the entire plant. When forage is to be fed to livestock, DO NOT apply within 7 days of harvest.

POTATOES, TOMATOES, EGGPLANT and PEPPERS - For control of Colorado Potato Beetle, Leafhoppers and Flea Beetles, apply approximately $\frac{1}{2}$ to $\frac{1}{2}$ lb./1000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, and Lygus Bugs, apply approximately $\frac{3}{4}$ to $\frac{9}{10}$ lbs./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH - For control of Pickleworm, Melonworm, Cucumber Beetle, Squash Bugs, Flea Beetles, apply approximately $\frac{1}{2}$ lb./1000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES - For control of Meadow Spittlebug, Leaf Roller, apply approximately $\frac{3}{4}$ to $\frac{9}{10}$ lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS - For control of Japanese Beetles, Periodical Cicada, Aphid, Tent Caterpillars, Lace Bugs, Boxelder Leaf Miner and certain other insects, apply approximately 1 lb./1000 sq. ft.

ANIMALS, DOG, CATS - For control of Fleas and Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Use sleeping quarters and infested areas weekly. DO NOT treat kittens under 4 weeks old. DO NOT apply to animals more than once weekly.

SIDE PANEL

CAUTION

HARMFUL IN INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid prolonged and repeated contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

DO NOT reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

20.11.14, 1970

20.11.14, 1970

DRAGON
5%
SEVIN* DUST

<p>ACCEPTED</p> <p>APR 3 1971</p> <p>UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER- ED UNDER NO. <u>16-27</u> SUBJECT TO ATTACHED COMMENTS.</p>
--