

16-27

1-4-1972

1/5

October 1, 1961

USDA REG. NO. 16-27

DRAGON
5%
SEVIN* DUST

ACCEPTED
JAN 4 1972
UNDER THE FEDERAL INSECTICIDE
FUNDING AND REIMBURSEMENT
FOR ECONOMIC PESTION REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED CONDITIONS.

16-27

16-27
16-27

FRONT PANEL

2/5

**DRAGON
5%
SEVIN* DUST**

**NET WEIGHT
4 lb.**

**FOR CONTROL OF JAPANESE BEETLE, FLEA BEETLE, MEXICAN BEAN BEETLE, TOMATO HORNWORM,
CORN EARWORM, AND CERTAIN OTHER INSECTS ATTACKING VEGETABLES, GRAPES, ORNAMENTALS
AND SHRUBS.**


ACTIVE INGREDIENTS	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate).....	5%
INERT INGREDIENTS.....	95%
Total	100%

**CAUTION: KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional
cautions.**

*** Trade Mark of UCC for 1-naphthyl N-methylcarbamate.**

**Manufactured By
DRAGON CHEMICAL CORPORATION
Roanoke, Virginia 24019**

USDA REG. NO. 16-27



BACK PANEL

3/5

**DRAGON
5%
SEVIN* DUST**

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON 5% SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

DIRECTIONS

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, apply approximately 1/2 lb./1000 sq. ft. DO NOT feed treated bean vines to dairy or meat animals.

LETTUCE - For control of Lygus and Stink Bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN - For control of Corn Earworm, apply approximately 3/4 to 9/10 lb./1000 sq. ft. Make 3-4 applications at 2 to 3 day intervals beginning when silks first appear. For control of European Corn Borer, apply approximately 3/4 to 9/10 lb./1000 sq. ft. Make application into the whorls and to the entire plant. When forage is to be fed to livestock, DO NOT apply within 7 days of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

POTATOES, TOMATOES, EGGPLANT and PEPPERS - For control of Colorado Potato Beetle, Leafhoppers and Flea Beetles, apply approximately 1/2 to 1/2 lb./1000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Corer, Stink Bugs, and Lygus Bugs, apply approximately 3/4 to 9/10 lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, apply approximately 1/2 lb./1000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES - For control of Meadow Spittlebug, Leaf Roller, apply approximately 3/4 to 9/10 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS - For control of Japanese Beetles, Periodical Cicada, Aphid, Tent Caterpillars, Lace Bugs, Boxwood Leaf Miner and certain other insects, apply approximately 1 lb./1000 sq. ft.

TOBACCO*- For control of Hornworms, Budworms, Flea Beetles, June Beetles and Japanese Beetles, apply 20-40 lbs./acre. In plant beds for control of Flea Beetle, use at the rate of 1 to 2 lbs./100 sq. ft.

ANIMALS, DOGS, CATS - For control of Fleas and Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Dust only outside sleeping quarters and infested areas weekly. **DO NOT** treat infested areas in the home. **DO NOT** treat kittens or puppies under 4 weeks old. **DO NOT** apply to animals more than once weekly.

* Directions for TOBACCO will appear only on 25 lb. bags.

5/5

SIDE PANEL

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid prolonged and repeated contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

DO NOT re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.