

m-21642-00 16-27

8-16-1972

1/3

4 LBS. NET WT.



CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.** DO NOT store near feed or foodstuffs.

5%

SEVIN* DUST

LISTED

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

FOR THE CONTROL OF JAPANESE BEETLE, FLEA BEETLE, MEXICAN BEAN BEETLE, TOMATO HORNWORM, CORN EARWORM, AND CERTAIN OTHER INSECTS ATTACKING VEGETABLES, GRAPES, ORNAMENTALS, AND SHRUBS.

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methylcarbamate)

INERT INGREDIENTS

BY WT.

5%

95%

TOTAL 100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK/SIDE PANELS FOR ADDITIONAL CAUTIONS.

*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

EPA REG. NO. 16-27

**MANUFACTURED BY
DRAGON CHEMICAL CORPORATION
ROANOKE, VIRGINIA**

2/3



5% SEVIN* DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON 5% SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. **DO NOT** use with lime or other alkaline material. May be applied up to harvest except where noted.

DIRECTIONS

BEANS—For control of Mexican bean beetles, leafhoppers, flea beetles, apply approximately 1/2 lb./1000 sq. ft. **DO NOT** feed treated bean vines to dairy or meat animals.

LETTUCE—For control of lygus and stink bugs, apply approximately 1 lb./1000 sq. ft. **DO NOT** apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN—Use approximately 3/4 lb./1000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS—For control of Colorado potato beetles, leafhoppers, and flea beetles, apply approximately 1/2 lb./1000 sq. ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bug and lygus bugs, apply approximately 3/4 lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH—For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles, apply approximately 1/2 lb./1000 sq. ft.

GRAPES—For control of leafhoppers, Japanese beetle, grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES—For control of meadow spittlebug, leaf roller, apply approximately 1 lb./1000 sq. ft. **DO NOT** apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS—For control of Japanese beetles, periodical cicada, aphid, tent caterpillars, lace bugs, boxwood leaf miner, apply approximately 1-1 1/4 lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. **DO NOT** use on Boston Ivy.

ANIMALS, DOGS, CATS—For control of fleas and brown dog ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. **DO NOT** treat kittens or puppies under 4 weeks old. **DO NOT** apply to animals more than once weekly.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. **DO NOT** breathe dust. **DO NOT** take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.** **DO NOT** store near feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.



SIFT-GARD

0000

7072

ACCEPTED

AUG 16 1972

BY THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FEDERAL REGISTER
 SUBJECT
 TO ATTACHED COMMENTS.

16-27