



CAUTION
 HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust.
 DO NOT take internally. Avoid contact with skin. If on skin
 wash with soap. Wash hands before eating. Take shower if
 on shirt. Wash laundry before wearing. Change to clean clothing
 after use. **DO NOT FEED LIVESTOCK.** Carbaryl is a
 contact, stomach and respiratory irritant. Absorption is con-
 siderable. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**
 DO NOT store near food or feedstuffs.

This product is highly toxic to bees exposed to direct treatment or
 residues on crops. Protective information may be obtained from
 your Cooperative Agricultural Extension Service.
 Do not re-use empty container. Destroy it by burning with waste or
 burning. Stay away from smoke and fumes.

NOTICE
 Buyer assumes all risks of use, storage or handling of this material
 and in strict accordance with DIRECTIONS and CAUTIONS stated
 on this label.

5% SEVIN* DUST

CAUTION
 HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust.
 DO NOT take internally. Avoid contact with skin. If on skin
 wash with soap. Wash hands before eating. Take shower if
 on shirt. Wash laundry before wearing. Change to clean clothing
 after use. **DO NOT FEED LIVESTOCK.** Carbaryl is a
 contact, stomach and respiratory irritant. Absorption is con-
 siderable. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**
 DO NOT store near food or feedstuffs.

This product is highly toxic to bees exposed to direct treatment or
 residues on crops. Protective information may be obtained from
 your Cooperative Agricultural Extension Service.
 Do not re-use empty container. Destroy it by burning with waste or
 burning. Stay away from smoke and fumes.

NOTICE
 Buyer assumes all risks of use, storage or handling of this material
 and in strict accordance with DIRECTIONS and CAUTIONS stated
 on this label.

FOR THE CONTROL OF JAPANESE BEETLE FLEA BEETLE MEXICAN BEAN BEETLE
 TOMATO HORNWORM CORN EARWORM AND CERTAIN OTHER LISTED INSECTS
 ON VEGETABLES GRAPES ORNAMENTALS AND SHRUBS

ACTIVE INGREDIENT	BY WT
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
	TOTAL 100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
 SEE BACK SIDE PANELS FOR ADDITIONAL CAUTIONS

MANUFACTURED BY
DRAGON CHEMICAL CORPORATION
 ROANOKE, VIRGINIA

READ
 WARN
 STATE
 DRA
 meet
 weigh
 7:10
 to ho
 READ
 warn
 STATE
 DRA
 meet
 weigh
 7:10
 to ho
 READ
 warn
 STATE
 DRA
 meet
 weigh
 7:10
 to ho

5% SEVIN* DUST

READ INSTRUCTIONS ON THIS PRODUCT CAREFULLY
WARNING: AVOID BREATHING DUST AND CONTACT WITH EYES
STATE REGULATIONS

GENERAL INFORMATION

DRAGON 5% SEVIN DUST

most effective insecticide for control of insects on
plants. It is applied to the upper and lower surfaces of leaves
7 TO 8 DAYS DO NOT use with rain or irrigation
to harvest except where noted

DIRECTIONS

BEANS For control of *Acanthosoma* and *Trioxys*
apply 1 lb. per acre to the soil in the row
before the plants are set in the ground

LETTUCE For control of *Spilonota* and *Trioxys*
apply 1 lb. per acre to the soil in the row
before the plants are set in the ground

CORN Use for control of *Acanthosoma* and *Trioxys*
Japanese beetles, leaf beetles, grasshoppers,
and other insects which damage the leaves
to entire plant beginning when the plants are
and the degree of protection desired against
proper timing of treatments. For some insects
also plants are not to be sprayed before
insects affecting silks and ears appear

and continuing until silks begin to dry. For control
on severity of infestation. Do not use on plants
growing of forage. Application of this product
and period will certainly reduce the number of

POTATOES, TOMATOES, CUCUMBERS AND PEPPERS
For control of leafhoppers and grasshoppers
apply 1 lb. per acre to the soil in the row
before the plants are set in the ground

CUCUMBERS SUMMER SQUASH For control of
leaf beetles apply approximately 1 lb. per
acre to the soil in the row before the plants
are set in the ground

GRAPES For control of leafhoppers, apply
approximately 1 lb. per acre to the soil in
the row before the plants are set in the ground

STRAWBERRIES For control of grasshoppers
apply 1 lb. per acre to the soil in the row
before the plants are set in the ground

ROSES, AZALEAS, SHADY TREES AND OTHER ORNAMENTALS
For control of grasshoppers, apply
approximately 1 lb. per acre to the soil in
the row before the plants are set in the ground

ROSES For control of grasshoppers, apply
approximately 1 lb. per acre to the soil in
the row before the plants are set in the ground