

| | | | |
|--|--|--|-------------------------------------|
| | United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460 | <input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other | OPP Identifier Number 188526 |
| | Application for Pesticide: | | |

Section I

| | | |
|--|-----------------------------------|--|
| 1. Company/Product Number 16-27 | 2. EPA Product Manager Edwards | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) DRAGON ^(R) 5% SEVIN ^(R) Dust | PM# 19 | |
| 5. Name and Address of Applicant (Include ZIP Code) Dragon Corporation P. O. Box 7311 Roanoke, VA 24019 <input type="checkbox"/> Check if this is a new address. | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ |

Section II

| | |
|---|---|
| <input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below. |
|---|---|

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Compliance with PR Notice 93-3 -- Labeling Statement Prohibiting Application to Water

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3

DATE 7/12/93

Section III

| | | | | | |
|---|--|---|-------------------|---|-------------------|
| 1. Material This Product Will Be Packaged In: | | | | 2. Type of Container | |
| Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | | <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ | |
| * Certification must be submitted. | | If "Yes," Unit Package wgt. | No. per container | If "Yes," Package wgt. | No. per container |
| 3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container | | 4. Size(s) of Retail Container 1LB / 4LB / 25LB | | 5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product | |
| 6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | | | |

Section IV

| | | |
|---|---------------------------------|--|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name Rebecca M. Horton | Title Executive Assistant | Telephone No. (Include Area Code) (703) 362-3657 |
| Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | 6. Date Application Received (Stamped) |
| 2. Signature | 3. Title Executive Assistant | |
| 4. Typed Name Rebecca M. Horton | 5. Date 6/21/93 | |

DRAGON®
5% SEVIN®
brand carbaryl insecticide
DUST

NOTIFICATION
LABEL NOT REVIEWED
PER FIF NOTICE 84-8
DATE 7/12/93

For the control of Japanese Beetles, Flea Beetles, Mexican Bean Beetles, Tomato Hornworm, Corn Earworm, and certain other listed insects attacking vegetables, grapes, ornamentals and shrubs.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm, Fleas and Brown dog ticks.]

ACTIVE INGREDIENT:

| | |
|---|------------|
| Carbaryl (1-naphthyl N-methylcarbamate) | 5% |
| INERT INGREDIENTS: | <u>95%</u> |
| TOTAL: | 100% |

Keep Out of Reach of Children
CAUTION
See back/side/other panel for additional cautions.

NET WEIGHT 1 LB/4 LB/25 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON 5% SEVIN® DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. **DO NOT** use with lime or other alkaline material. May be applied up to harvest except where noted.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: For control of Mexican bean beetles, Leafhoppers, Flea beetles, apply approximately 1/2 lb./1000 sq. ft. **DO NOT** feed treated bean vines to dairy or meat animals.

LETTUCE: For control of Lygus and Stink bugs, apply approximately 1 lb./1000 sq. ft. **DO NOT** apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN: Use approximately 3/4 lb./1000 sq. ft. for Corn earworm, Fall armyworm, Flea beetles, Japanese beetles, Sap beetles, Leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are

about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For control of Colorado potato beetles, Leafhoppers and Flea beetles, apply approximately ½ lb./1000 sq. ft. For control of Tomato fruitworms, Fall armyworm, Tomato hornworm, European corn borer, Stink bug and Lygus bugs, apply approximately ¾ lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH: For control of Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetles, apply approximately ½ lb./1000 sq. ft.

GRAPES: For control of Leafhoppers, Japanese beetle, Grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES: For control of Meadow spittlebug, Leaf roller, apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS: For control of Japanese beetles, Periodical cicada, Aphid, Tent caterpillars, Lace bugs, Boxwood leaf miner, apply approximately 1-1¼ lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.

DOGS, CATS: For control of fleas and brown dog ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. DO NOT treat kittens or puppies under 4 weeks old. DO NOT apply more than once weekly. DO NOT use on pregnant dogs.

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. DO NOT take internally. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

0/04/1/

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.


PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019



Label of Record - as submitted under 6/25/92
Notification

5-11

DRAGON®
5% SEVIN®
brand carbaryl insecticide
DUST

EPA Reg. No. 16-27

EPA Est. No. 16-VA-1

For the control of Japanese Beetles, Flea Beetles, Mexican Bean Beetles, Tomato Hornworm, Corn Earworm, and certain other listed insects attacking vegetables, grapes, ornamentals and shrubs.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm, Fleas and Brown dog ticks.]

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%

TOTAL: 100%

Keep Out of Reach of Children

CAUTION

See back/side/other panel for additional cautions.

NET WEIGHT 1 LB/4 LB/25 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. DO NOT take internally. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

GENERAL INFORMATION

DRAGON 5% SEVIN® DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: For control of Mexican bean beetles, Leafhoppers, Flea beetles, apply approximately $\frac{1}{2}$ lb./1000 sq. ft. DO NOT feed treated bean vines to dairy or meat animals.

LETTUCE: For control of Lygus and Stink bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN: Use approximately $\frac{1}{4}$ lb./1000 sq. ft. for Corn earworm, Fall armyworm, Flea beetles, Japanese beetles, Sap beetles, Leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage; Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For control of Colorado potato beetles, Leafhoppers and Flea beetles, apply approximately $\frac{1}{2}$ lb./1000 sq. ft. For control of Tomato fruitworms, Fall armyworm, Tomato hornworm, European corn borer, Stink bug and Lygus bugs, apply approximately $\frac{1}{4}$ lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH: For control of Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetles, apply approximately ½ lb./1000 sq. ft.

GRAPES: For control of Leafhoppers, Japanese beetle, Grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES: For control of Meadow spittlebug, Leaf roller, apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS: For control of Japanese beetles, Periodical cicada, Aphid, Tent caterpillars, Lace bugs, Boxwood leaf miner, apply approximately 1-1½ lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.

DOGS, CATS: For control of fleas and brown dog ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. DO NOT treat kittens or puppies under 4 weeks old. DO NOT apply more than once weekly. DO NOT use on pregnant dogs.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019