

[FRONT PANEL]

DRAGON®
5% SEVIN® DUST
brand carbaryl insecticide

[Additional Brand Name: DRAGON SEVIN 5% Dust]

Nothing Beats The Dragon!

For the control of Japanese Beetles, Flea Beetles, Mexican Bean Beetles, Tomato Hornworm, Corn Earworm, and certain other listed insects attacking vegetables, grapes, ornamentals and shrubs.
Broad spectrum outdoor insect control.
Use on vegetables, grapes, ornamentals and others listed.
Kills fleas and ticks on dogs and cats.
Controls listed insects attacking vegetables, grapes, ornamentals and shrubs.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm, Fleas and Brown dog ticks.]

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%
..... 100%

ACCEPTED
FEB 28 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, is amended, for the pesticide registered under EPA Reg. No. 16-27

Keep Out of Reach of Children
CAUTION
See back/side/other panel for additional cautions.
NET WEIGHT 1 LB/4 LB/20 LB

REC'D
NOV 20 1996
P/DPD1

[BACK/SIDE/OTHER PANEL(S)]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. DO NOT take internally. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. [NOTE to EPA: When marketed for use on dogs and cats, the following statement will be included on the label: Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.]

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

BEST COPY AVAILABLE

NOTE TO PHYSICIAN [Note to EPA: When marketed for use on dogs and cats, the following will be included on the label: & **VETERINARIAN**]: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

[VERSION A - text "lined-out" format]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOMEOWNER USE ONLY. DRAGON® 5% SEVIN® DUST controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using **DRAGON® SEVIN® Liquid**. Do not repeat application more than once a week. Do not use with lime or other alkaline materials. **DRAGON® 5% SEVIN Dust** may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: To control (kill) Mexican bean beetles, Leafhoppers, Flea beetles, apply approximately 1/2 lb./1000 sq. ft. DO NOT feed treated bean vines to domestic animals.

LETTUCE: To control (kill) Lygus and Stink bugs, apply approximately 1 lb./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN: Use approximately 3/4 lb./1000 sq. ft. for Corn earworms, Fall armyworms, Flea beetles, Japanese beetles, Sap beetles, Leafhoppers and European corn borers. For corn borers and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: To control (kill) Colorado potato beetles, Leafhoppers and Flea beetles, apply approximately 1/2 lb./1000 sq. ft. To control (kill) Tomato fruitworms, Fall armyworms, Tomato hornworms, European corn borers, Stink bugs and Lygus bugs, apply approximately 3/4 lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH: To control (kill) Pickleworms, Melonworms, Cucumber beetles, Squash bugs, Flea beetles, apply approximately 1/2 lb./1000 sq. ft.

GRAPES: To control (kill) Leafhoppers, Japanese beetles, Grape berry moths, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES: To control (kill) Meadow spittlebugs and Leaf rollers, apply approximately 1 lb./1000 sq. ft. DO NOT apply within 1 day of harvest.

BEST COPY AVAILABLE

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS: To control (kill) Japanese beetles, Periodical cicada, Aphids, Tent caterpillars, Lace bugs and Boxwood leaf miners, apply approximately 1-1¼ lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. **DO NOT** use on Boston Ivy.

DOGS, CATS - READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS OR CATS.
To control (kill) Fleas and Brown Dog Ticks: Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pets eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Do not repeat treatment for 1 week. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals. Dust bedding and doghouse using a shaker can or other dust applicator. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS** and **CAUTIONS** stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

[VERSION B - "table" format]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOMEOWNER USE ONLY. DRAGON® 5% SEVIN® DUST controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using DRAGON® SEVIN® Liquid. Do not repeat application more than once a week. Do not use with lime or other alkaline materials. DRAGON® 5% SEVIN Dust may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEST COPY AVAILABLE

475

VEGETABLE GARDENS		
Crop	Insects	Use Rate per 1,000 sq. ft.
BEANS Do not feed treated bean vines to domestic animals.	Mexican Bean Beetles, Leafhoppers, Flea Beetles	½ lb.
LETTUCE Do not apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.	Lygus and Stink Bugs	1 lb.
CORN For corn borers and larvae in whorls (budworm damage) and for foliage feeders, apply	Corn Earworms, Fall Armyworms, Flea Beetles, Japanese Beetles, Sap Beetles, Leafhoppers and European Corn Borers	¾ lb.
thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.		
POTATOES, TOMATOES, EGGPLANT, PEPPERS	Colorado Potato Beetles, Flea Beetles and Leafhoppers	½ lb.
	Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers, Stink Bugs and Lace Bugs	¾ lb.
CUCUMBERS, SUMMER SQUASH	Pickleworms, Melonworms, Cucumber Beetles, Squash Bugs, and Flea Beetles	½ lb.

FRUITS		
Crop	Insects	Use Rate per 1,000 sq. ft.
GRAPES	Leafhoppers, Japanese Beetles, and Grape Berry Moths	1 lb.
STRAWBERRIES Do not apply within 1 day of harvest.	Meadow Spittle Bugs and Leaf Rollers	1 lb.

BEST COPY AVAILABLE

BEST COPY AVAILABLE

5 7 5

ROSES, SHADE TREES & ORNAMENTAL GARDENS		
Site	Insects	Use Rate per 1,000 sq. ft.
ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. Do not use on Boston Ivy.	Japanese Beetles, Periodical Cicada, Aphids, Tent Caterpillars, Lace Bugs, and Boxwood Leaf Miners	1 to 1 1/4 lb.

DOGS & CATS	
READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS OR CATS.	
Pest	Directions
FLEAS, BROWN DOG TICKS	Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pets eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Do not repeat treatment for 1 week. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals. Dust bedding and doghouse using a shaker can or other dust applicator. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
 Roanoke, VA 24019