



## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

FEB 10 1921

Dragon Corporation P.O. Box 7311 Roanoke, VA 24019

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Subject:

Label Amendment Submission of 7/29/93 in Response to PR Notice 93-7

EPA Reg. No. 16-28 Dragon 1-3/4% Sevin

#### Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed. The comments follow.

#### WHAT THIS ACCEPTANCE MEANS:


Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

#### WHAT YOU NEED TO DO NEXT:

Send to EPA one (1) copy of the final printed labeling:



- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.



Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

James A. Downing Firm Tompkins, Deputy Chief

Registration Support Branch

Registration Division (7505W)

Attachment

[FRONT PANEL]

# DRAGON® 13/4% SEVIN® brand carbaryl insecticide DUST

ACCEPTED with COMMENTS In EPA Letter Dated FEB 1 0 1994

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Rog. No.

Nothing Beats The Dragon!

Kills Japanese Beetles, Flea Beetles, Mexican Bean Beetles, Tomato Hornworm, Corn Earworm, and certain other listed insects attacking vegetables, ornamentals and shrubs.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm.]

#### **ACTIVE INGREDIENT:**

Carbaryl (1-naphthyl N-methylcarbamate)	1.75%
INERT INGREDIENTS:	
TOTAL:	00.00%

## Keep Out of Reach of Children CAUTION

See back/side/other panel(s) for additional cautions.

#### **NET WT. 4 LBS**

[BACK/SIDE/OTHER PANEL(S)]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

#### **GENERAL INFORMATION**

FOR HOMEOWNER USE ONLY. DRAGON 13/4% SEVIN® DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted. Subsequent planting of domestic food crops in treated soil must be made with plants on this or other carbaryl labels.

#### **DIRECTIONS**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

BEANS: For Mexican bean beetles, Leafhoppers, Flea beetles, apply approximately 1½ lbs./1000 sq. ft. DO NGT feed treated bean vines to domestic animals.

LETTUCE: For Lygus and Stink bugs, apply approximately 2-2½ lbs./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN: For Corn earworm, Fall armyworm, Flea beetles, Japanese beetles, Sap beetles, Leafhoppers and European corn borer apply approximately 2-2½ lbs./1000 sq. ft. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear

and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

- POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For Colorado potato beeties, Leashoppers and Flea beetles, apply approximately 3/4-11/4 lbs./1000 sq. ft. For Tomato fruitworm, Fall armyworm, Tomato hornworm, European corn borer, Stink bug and Lygus bugs, apply approximately 2-21/2 lbs./1000 sq. ft.
- CUCUMBERS, SUMMER SQUASH: For Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetles, apply approximately 1% lbs./1000 sq. ft.
- GRAPES: For Leafhoppers, Japanese beetle, Grape berry moth, apply approximately 2<sup>3</sup>/<sub>4</sub> lbs./1000 sq. ft.
- STRAWBERRIES: For Meadow spittlebug, Leaf roller, apply approximately 1<sup>2</sup>/<sub>3</sub> lbs./1000 sq. ft. DO NOT apply within 1 day of harvest.
- ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS: For Japanese beetles, Periodical cicada, Aphid, Tent caterpillars, Lace bugs, Boxwood leaf miner and certain other insects, apply approximately 2-4 lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: May be harmfut if swallowed. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash thoroughly after handling.

#### STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Induce vomiting and seek medical attention immediately.
- IF IN EYES OR ON SKIN: Flush eyes with plenty of water. Wash skin thoroughly with soap and water.
- NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.
- ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.
- STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

#### STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhone Poulenc Ag Company.

Manufactured by DRAGON CORPORATION Roanoke, VA 24019