[FRONT PANEL]

DRAGON® 1³/4% SEVIN® brand carbaryl insecticide DUST

Nothing Beats The Dragon!

Kills Japanese Beetles, Flea Beetles, Mexican Bean Beetles, Tomato Hornworm, Corn Earworm, and certain other listed insects attacking vegetables, ornamentals and shrubs.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects controlled - Japanese beetle, Bean beetle, Tomato hornworm.]

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)

1.75%

INERT INGREDIENTS:

1.75 %

TOTAL:

Keep Out of Reach of Children
CAUTION
See back/side/other panel(s) for additional cautions.

NET WT. 4 LB

FEB 2 4 1997
Under the Federal Insecticide.
Fungleide, and Rodenticlide Act.
as amended, by the pesticide
registered under
EPA Reg. No. 11-2-5

[BACK/SIDE/OTHER PANEL(S)]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

FIRST AID

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes with plenty of water. Wash skin thoroughly with soap and water.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

EPA Reg. No. 16-28 November 12, 1996

EPA Est. 16-VA-1

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOMEOWNER USE ONLY. DRAGON® 1³/4% SEVIN® DUST controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using DRAGON® SEVIN® Liquid. Do not repeat application more than once a week. Do not use with lime or other alkaline materials. DRAGON® 1³/4% SEVIN® Dust may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

- BEANS: For Mexican bean beetles, Leafhoppers, Flea beetles, apply approximately 1½ lbs./1000 sq. ft. DO NOT feed treated bean vines to domestic animals.
- LETTUCE: For Lygus and Stink bugs, apply approximately 2-2½ lbs./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.
- CORN: For Corn earworm, Fall armyworm, Flea beetles, Japanese beetles, Sap beetles, Leafhoppers and European corn borer apply approximately 2-2½ lbs./1000 sq. ft.For corn borers and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.
- POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For Colorado potato beetles, Leafhoppers and Flea beetles, apply approximately ³/₄-1¹/₄ lbs./1000 sq. ft. For Tomato fruitworm, Fall armyworm, Tomato hornworm, European corn borer, Stink bug and Lygus bugs, apply approximately 2-2¹/₂ lbs./1000 sq. ft.
- CUCUMBERS, SUMMER SQUASH: For Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetles, apply approximately 11/4 lbs./1000 sq. ft.
- GRAPES: For Leafhoppers, Japanese beetle, Grape berry moth, apply approximately 2³/₄ lbs./1000 sq. ft.
- STRAWBERRIES: For Meadow spittlebug, Leaf roller, apply approximately 1²/₃ lbs./1000 sq. ft. DO NOT apply within 1 day of harvest.
- ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS: For Japanese beetles, Periodical cicada, Aphid, Tent caterpillars, Lace bugs, Boxwood leaf miner and certain other insects, apply approximately 2-4 lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. DO NOT use on Boston Ivy.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by DRAGON CORPORATION Roanoke, VA 24019