1 9 3

DRAGON® MIRACLE WEED PREVENTER

Prevents weeds in flower and vegetable gardens and around trees and shrubs.

READY TO USE DACTHAL GRANULES
THIS BOX TREATS 600 SQUARE FEET

CAUTION
SEE BACK AND SIDE PANELS
FOR ADDITIONAL CAUTIONS.
NET WEIGHT 3 LB.

ACTIVE INGREDIENT:	By Wt.	
ACTIVE INGREDIENT: DCPA (dimethyl tetrachloroterephthalate)	5%	V .
INERT INGREDIENTS:		
TOTAL	100%	

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

Environmental Hazards: Collect and incorporate granules spilled on soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes.

Groundwater Advisory: Tetrachloroterephthalic acid, a breakdown product of Dacthal, is known to leach through soil as a result of agricultural and turf uses and has been found in groundwater which may be used for drinking water. Users are advised not to apply Dacthal to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater used for drinking water.

MIRACLE WEED PREVENTER is approved for use on the following:

Abelia	Coneflower, purple	Gaillardia	Magnolia	Privet
Ageratum	Coreopsis	Geranium	Maple	Redbud
Alyssum	Coralbells	Gladiolus	Marguerite, golden	Rhododendron
Andromeda, Jap	Cosmos	Golddust	Marigold	Russian Olive
Arabovitae	Contonester	Goldentuft	Mockorange	Sage, scarlet
Ash	Crabapple	Gum	Morningglory	Snapdragon
Aster	Cuphea	Hawthorn	Moss rose	Spiderwort, Va
Azalea	Cypress	Heath, pink	Mother-of-Thyme	Spirea
Babysbreath	Dahlia	Hally	Mountain Laurel	Spruce
Barberry	Delphinium	Honeysuckle	Mourning Bride	Stonecrop
Bellflower	Deutzia	Iris	Nasturtium	Strawilower
Birch	Dogwood	Ivy, Baltic	Oak	Sundrops
Bleedingheart	Elaeagnus	Ivy, English	Orpine	Sunflower
Bloodleaf	Elm	lvy, Wilson	Pachistima	Swee: Pea
Boxwood	Euonymus	Juniper	Pachysandra	Sycamore
Bugloss	Evening primrose	Lantana	Peony	Juse heony
Camellia	Fevertew	Larkspur, candle	Petunia	Tulip tree
Candytufts	Fir	Lavendercotton	Pine	Viburnum
Chestnut	Forget-me-not	Lilac	Pittosporum	Violet
Chysanthemum	Forsythia	Lily	Podocarpus	Walnut
Cinquefoil	Four O'Glock	Locust	Poker Plant	Weigela
Coleur	CEPPED	Lupine	Poplar	Worrnwood
Columbine			-	Yarrow, ferleaf
	mild interest			Now

ACCEPTEL

Page 1 of 3 Pages

Under the Federal Intended in Pungionse, and Rodenburde Action amended, for the periode

Weeds Controlled by Miracle Weed Preventer

Carpetweed	Lovegrass	Purslane	Jonnsongrass
Yellow Foxtail	Fall Panicum	Witch grass	Smooth Crabgrass
Green Foxtail	Nodding Spurge	Florida Pussley	Common Chickweed
Texas Millet	Lambsquarter	Hairy Crabgrass	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use only when wearing the following protective clothing:

Wear long-sleeved shirt and long legged pants and chemical resistant gloves.

IMPORTANT! If the pesticide comes in contact with skin wash off with soap and water. Always wash hruids, face, and arms with soap and water before smoking, eating, drinking, or toileting.

AFTER USE: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during use must be laundered separately from household articles.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

- APPLICATION RATE: 2 1/2 Cups (1 lb.) spread evenly over 200 square feet of weed free soil. Water in thoroughly.
- DRAGON MIRACLE WEED PREVENTER is a 5% granular DACTHAL formulation for use in preemergence (before weeds appear) weed control of crabgrass and other annual grasses and broadleaf weeds. (See side panel for list of weeds and grasses controlled.) It is recommended for use in home vegetable and flower gardens, and around trees and shrubs as listed on this label.
- DIRECTIONS: For best results, MIRACLE WEED PREVENTER should be applied to a well-prepared soil which is free of weeds and PRIOR to weed seed germination. For garden soils, apply to a uniform weed free soil within 2 to 3 days following cultivation to prevent new weed seed from germinating. Disturbing the soil surface after application will reduce effectiveness. Measure the area to be treated and divide into small plots. Sprinkle the granules evenly over the soil and water in thoroughly. FOUR MONTH control can be expected when applied properly.
- FLOWER GARDENS: Apply to the weed-free soil around the listed plants after they are established or after transplanting. Apply MIRACLE WEED PREVENTER at the rate of 2 1/2 cups (1 lb.) per 200 sq. it. Water in thoroughly, rinsing all granules off plant foliage. Additional applications, when needed, can be made at any time provided beds have been cultivated and are weed free. Do not use on Phlox, Bugle Weed, Button Pink, Carnation, Geum, Germander, Pansy, Sweet William, Telanthera, Masambtyanthem.
- ORNAMENTALS, SHRUBS AND TREES (As listed on side panel): Apply 2 1/2 cups (1 lb.) MIRACLE WEED PREVENTER per 200 sq. ft. uniformly to soil around the base of plants in late fall or early spring. May be applied following transplanting into well prepared soil. For established plantings apply proper cultivation to remove existing weeds. SEE SIDE PANEL FOR L'ST OF TREES AND SHRUBS THIS PRODUCT MAY BE APPLIED AROUND.

- **VEGETABLE CROPS:** Any application of **MIRACLE WEED PREVENTER** should be made prior to weed seed germination. Apply at seeding to snap beans, broccoli, brussels sprouts, cabbage, cauliflower, collards, garlic, kale, mustard greens, onions, potatoes, (Southern peas) turnip greens, turnips.
- Apply to seeded cucumbers, rielons (cantaloupe, watermelons, and honeydew), squash and head lettuce about 4 weeks after seeding, but no later than 6 weeks. For leaf lettuce, make a single application 1-3 weeks after emergence. When transplanting broccoli, brussels sprouts, cabbage, cauliflower, eggplant, peppers, onions, and sweet potatoes, apply immediately after transplanting.
- **TOMATOES:** Apply only to well established plants, 4 to 6 weeks after transplanting, when conditions are favorable.
- STRAWBERRIES: Apply to strawberries following transplanting or on established strawberry plants prior to weed seed germination or after clean cultivation. On established plants, do not apply after first bloom.
- **REPLANTING:** Replanting crops other than those included on the label in DACTHAL treated soil within 8 months of application may result in crop injury. If replanting is required because of early crop failure, the planting of onions, seeded melons, potatoes, tomatoes, eggplants, or peppers at this time may result in crop injury. However, all crops listed on this label may be planted following harvest of a crop treated with this product.
- STORAGE: C not store near food or foodstuffs. Store in the tightly closed original container in a cool, dry preferacity locked storage area, out of reach of children.
- PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
- CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Distributed By DRAGON CORPORATION

Roanoke, Va 24019-0311

EPA Reg. No. 16-42

EPA Esí. 16-VA-1

Form #L-42

Α