10-11-Please read instructions on reverse before completing form Form Approved. OMB No. 2070-0060. Approval expires 11-30-93 OPP Identifier Number United States Environmental Protection Agency Registration Office of Pesticide Programs (H7505C) **Amendment** Washington, DC 20460 196863 Application for Pesticide: Other Section I 3. Proposed Classification 2. EPA Product Manager 1. Company/Product Number 16-42 Miller Restricted PMs 4. Company/Product (Name) DRAGON Miracle Weed Preventer 23 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) 5 Name and Address of Applicant (Include ZIP Code) (b)(i), my product is similar or identical in composition and labeling Dragon Corporation to: P. O. Box 7311 Roanoke, VA 24019 EPA Reg. No._ Check if this is a new address **Product Name** Section I I Final printed labels in response to Amendment - Explain below Agency letter dated_ Resubmission in response to Agency letter dated "Me Too" Application Notification - Explain below. Other - explain below Explanation: Use additional page(s) if necessary (For section I and Section II) Compliance with PR Notice 93-3 -- Statement Prohibiting Application to Water Section III 1. Material This Product Will Be Packaged in: Child Resistant Packaging **Unit Fackaging** Water Soluble Packaging Type of Container Yes Metal Yes Plastic No Glass No Paper If "Yes." If "Yes." No per No per Other (Specify) Unit Package wgt container Package wgt. container * Certification must be submitted. Location of Label Directions 3 Location of Net Contents Information Size(s) of Retail Container ▼ On Label Container 3LBOn Labeling accompanying product 6 Manner In Which Label Is Affixed To Product Lithograph Other (Paper glued Stenaled ide)

âme	Title	Telephone Nov (Include Area Co (703) 362-3657	
Rebecca M. Horton	Executive Assistant		
Certify that the statements I have made on this form a lacknowledge that any knowingly false or misleading both under applicable law			6 Date Application Recurred (Stainped)
Relucca M. Hoston	3 Time Executive Assistant		••••
Typed Name Rebecca M. Horton	5 Date 6/22/93	<u>-</u>	

DRAGON³ MIRACLE WEED PREVENTER

Prevents weeds in flower and vegetable gardens, and around trees and shrubs.

READY TO USE DACTHAL GRANULES THIS BOX TREATS 600 SQUARE FEET

١.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., vegetable or flower garden.]

ACTIVE INGREDIENT:	BY	Y WT.
DCPA (dimethyl tetrachloroterephthalate)		. 5%
INERT INGREDIENTS:		95%
TOTAL		100%

Keep Out of Reach of Children CAUTION See side/back/other panel(s) for additional cautions.

NET WT. 3 LB

NCTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 84-0
7/12/93

DATE 7/12/93

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZ.\RDS: Collect and incorporate granules spilled on soil surface. For terrestrial uses, do not apply directly to watch, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Groundwater Advisory: Tetrachloroterephthalic acid, a breakdown product of Dacthal, is known to leach through soil as a result of agricultural and turf uses and has been found in groundwater which may be used for drinking water. Users are advised not to apply Dacthal to sand and loamy sand soils where the water table (groundwater) is close to the surface and where diese soils are very permeable, i.e., well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater used for drinking water.

MIRACLE WEED PREVENTER is approved for use on the following:

Abelia	Barberry	Chrysanthemum	Crahapple	Evening primrose
Ageratum	Bellflower	Cinquefoil	Cuphea	Feverfew
Alyssum	Birch	Coleus	Cypress	Fir
Andromeda,	Bleedingheart	Columbine	Dahlia	Forget-me-nor
Japanese	Bloouleaf	Coneflower,	Delphinium	Forsythia
Arborvitae	Boxwood	Purple	Deutzia	Four-O'clock
Ash	Bugloss	Coreopsis	Dogwood	Fox love
Aster	Camellia	Coral bells	Elaegnus	Gaillardia
Azales	Candytufts	Cosmos	Elm	Geranium
Fabysbreath	Chestnut	Cotoneaster	Euonymus	· G!a Jiolus

EPA Reg. No. 16-42

EPA Est. 16-VA-1

June 22, 1993

Page 1

Golddust	Larkspur, candle	Mother-of-Thyme	Poker plant	Sundrops
Goldentuft	Lavendercotton	Mountain laurel	Poplar	Sunflower
Gum	Lilac	Mourning bride	Privet	Sweetpea
Hawthorn	Lily	Nasturtium	Redhud	Sycamore
Heath, pink	Locust	Oak	Rhododendron	ree peony
Holly	Lupine	Orpine	Russian olive	Tulip tree
Honeysuckle	Magnolia	Pachistima .	Sage, scarlet	Vibumum
Iris	iviaple	Pachysandra	Snapdragon	Violet
Ivy, Baltic	Marguerite, golden	Peony	Spiderwort, Virginia	Walnut
Ivy, English	Marigold	Petunia	Spirea	Weigela
Ivy, Wilson	Mockorange	Pine	Spruce	Wormwood
Juniper	Morningglory	Pittosporum	Stonecrop	Yarrow, femleaf
Lantana	Moss rose	Podocarous	Strawflower	Yew

Weeds Controlled by Miracle Weed Preventer

	Weens Controlled by	y Minacle Weed Fictence	
Carpetweed	Lovegrass	Purslane	Johnsongrass
Yellow Foxtail	Fall Panicum	Witch grass	Smooth crabgrass
Green Foxtail	Nodding Spurge	Florida Pussley	Common chickweed
Texas Millet	Lambsquarter	Hairy Crabgrass	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use only when wearing the following protective clothing: Wear long-sleeved shirt and long-legged pants and chemical resistant gloves.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face and arms with soap and water before smoking, eating, drinking or toileting.

AFTER USE: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during use must be laundered separately from household articles.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

APPLICATION RATE: 2½ cups (1 lb.) spread evenly over 200 square feet of weed-free soil. Water in thoroughly.

DRAGON® MIRACLE WEED PREVENTER is a 5% granular DACTHAL formulation for use in preemergency (before weeds appear) weed control of crabgrass and other annual grasses and broadleaf weeds. (See side panel for list of weeds and grasses controlled.) It is recommended for use in home vegetable and flow in gardens, and around trees and shrubs as listed on this label.

DIRECTIONS: For best results, MIRACLE WEED PREVENTER should be applied to a well-prepared soil which is free of weeds and PRIOR to weed seed germination. For garden soils, apply to a uniform well-free soil within 2 to 3 days following cultivation to prevent new weed seed from germinating. Disturbing the soil surface after application will reduce effectiveness. Measure the area to be treated and divide into small plots

EPA Reg. No. 16-42 June 22, 1993

o. 16-42 EPA Est. 16-VA-1

Sprinkle the granules evenly over the soil and water in thoroughly. FOUK MONTH control can be expected when applied properly.

FLOWERING GARDENS: Apply to the weed-free soil around the listed plants after they are established or after transplanting. Apply MIRACLE WEED PREVENTER at the rate of 2½ cups (1 lb.) per 200 sq. ft. Water in thoroughly, rinsing all granules off plant foliage. Additional applications, when needed, can be made at any time provided beds have been cultivated and are weed free. Do not use on Phlox, Bugle weed, Button pink, Carnation, Geum, Germander, Pansy, Sweet William, Telanthera, Mesembryantheum.

ORNAMENTALS, SHRUBS and TREES (as listed on side panel): Apply 2½ cups (1 lb.) MIRACLE WEED PREVENTER per 200 sq. ft. uniformly to soil around the base of plants in late fall or early spring. May be applied following transplanting into well prepared soil. For established plantings, apply proper cultivation to remove existing weeds. SEE SIDE PANEL FOR LIST OF TREES AND SHRUBS THIS PRODUCT MAY BE APPLIED AROUND.

VFGETABLE CROPS: Any application of MIRACLE WEED PREVENTER should be made prior to weed seed germination. Apply at seeding to snap beans, broccoli, Brussels sprouts, cabbage, cauliflower, collards, garlic, kale, mustard greens, onions, potatoes, southern peas, turnip greens, turnips.

Apply to seeded cucumbers, melons (cantaloupe, watermelons and honeydew), squash and head lettuce about 4 weeks after seeding, but no later than 5 weeks. For leaf lettuce, make a single application 1-3 weeks after emergence. When transplanting broccoli, Brussels sprouts, cabbage, cauliflower, eggplant, peppers, onions and sweet potatoes, apply immediately after transplanting.

Tomatoes: Apply only to well-established plants, 4 to 6 weeks after transplanting, when conditions are favorable.

STRAWBERRIES: Apply to strawberries following transplanting or on established strawberry plants prior to weed seed germination or after clean cultivation. On established plants, do not apply after first bloom.


REPLANTING: Replanting crops other than those included on the label in DACTHAL treated soil within 8 months of application may result in crop injury. If replanting is required because of early crop failure, the planting of onions, seeded melons, potatoes, tomatoes, eggplants or peppers at this time may result in crop injury. However, all crops listed on this label may be planted following harvest of a crop t eated with this product.

STORAGE: Do not store near food or feedstuffs. Store in the tightly closed original container in a cool, dry preferably locked storage area, out of reach of children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Distributed by DRAGON CORPORATION Roanoke, VA 24019



DRAGON® MIRACLE WEED PREVENTER

Prevents weeds in flower and vegetable gardens and around trees and shrubs.

READY TO USE DACTHAL GRANULES THIS BOX TREATS 600 SQUARE FEET

Keep Out of Reach of Children CAUTION
SEE BACK AND SIDE PANELS
FOR ADDITIONAL CAUTIONS.
NET WEIGHT 3 LB.

ACTIVE INGREDIENT: DCPA (dimethyl tetrachloroterephthalate) S% INERT INGREDIENTS: TOTAL 100%

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water u after handling.

Environmental Hazards: Collect and incorporate granules spilled on soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes.

Groundwater Advisory: Tetrachloroterephthalic acid, a breakdown product of Dacthal, is known to leach through soil as a result of agricultural and turf uses and has been found in groundwater which may be used for drinking water. Users are advised not to apply Dacthal to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater used for drinking water.

MIRACLE WEED PREVENTER is approved for use on the following:

Abelia	Coneflower, purple	Gaillardia	Magnolia	Privet
Ageralum	Coreopsis	Geranium	Maple	Redbud
Alyssum	Coralbells	Gladiolus	Marguerite, golden	Rhododendron
Andromeda, Jap	Cosmos	Golddust	Marigold	Russian Olive
Arabovitae	Contonester	Goldentuft	Mockorange	Sage, scarlet
A sh	Crabapple	Gum	Morningglory	Snapdragori
Aster	Cuphea	Hawthorn	Moss rose	Spiderwort, Va
Azalea	Cypress	Heath, pink	 Mother-of-Thyme 	Spirea
Babysbreath	Dahlia	Holly	Mountain Laurel	Spruce
Barberry	Delphinium	Honeysuckle	Mourning Bride	Stonecrop
Bellflower	Deutzia	lris	Nasturtium	Strawflower
Birch	Dogwood	lvy, Baltic	Oak	Sundrops
Bleedingheart	Elaeagnus	lvy, Englisn	Orpine	Sunflower
Bloodleaf	Elm	lvy, Wilson	Pachistima	Sweet Pea
Boxwood	Euonymus	Juniper	Pachysandra	Sycamore
Bugloss	Evening primrose	Lantana	Peony	Jusa l la-duð
Camellia	Feverfew	Larkspur, candle	Petunia	Tulip*ti ee *
Candytuits	Fir	Lavendercotton	Pine	Viburnum
Chestnut	Forget-me-not	Lilac	Pittosporum	Violet . •
Chysanthemum	Forsythia	Lily	Podocarpus	Walnut,
Cinquigor	TE FRUTTO EL GENT	Locust	Poker Plant	Wa çela '
Coeus	11 IN F99992	Lupine	Poplar	Wormwood
Columbine3 ()	JUN 1302	·		Yarrow, Jelical
1 0 4				Yew y
Uniter the 1	Federal Insection in			- (: /:
For p. 1. S.				
147	11/2.42	Page 1 of 3 Pages		
FLY Hall				

Weeds Controlled by Miracle Weed Preventer

Carpetiveed
Yellow Foxtail
Green Foxtail
Texas Millet

Lovegrass Fall Panicum Nodding Spurge Lambsquarter

Purslane Witch grass Florida Pussley Hairy Crabgrass Johnsongrass Smooth Crabgrass Common Chickweed

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use only when wearing the following protective clothing:

Wear long-sleeved shirt and long legged pants and chemical resistant gloves.

IMPORTANT! If the pesticide comes in contact with skin wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or toileting.

AFTER USE: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during use must be laundered separately from household articles.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

APPLICATION RATE: 2 1/2 Cups (1 lb.) spread evenly clier 200 square feet of weed free soil. Water in thoroughly.

DRAGON MIRACLE WEED PREVENTER is a 5% granular DACTHAL formulation for use in preemergence (before weeds appear) weed control of crabgrass and other annual grasses and broadleaf weeds. (See side panel for list of weeds and grasses controlled.) It is recommended for use in home vegetable and flower gardens, and around trees and shrubs as listed on this label.

DIRECTIONS: For best results, MIRACLE WEED PREVENTER should be applied to a well-prepared soil which is free of weeds and PRIOR to weed seed germination. For garden soils, apply to a uniform weed-free soil within 2 to 3 days following cultivation to prevent new weed seed from germinating. Disturbing the soil surface after application will reduce effectiveness. Measure the area to be treated and divide into small plots. Sprinkle the granules evenly over the soil and water in thoroughly. FOUR MONTH control can be expected when applied properly.

FLOWER GARDENS: Apply to the weed-free soil around the listed plants after they are established or after transplanting. Apply MIRACLE WEED PREVENTER at the rate of 2 1/2 cups (1 lb.) per 200 sq. ft. Water in thoroughly, rinsing all granules off plant foliage. Additional applications, when needed, can be made at any time provided beds have been cultivated and are weed free. Do not use on Phlox, Bugle Weed, Button Pink, Carnation, Geum, Germander, Pansy, Sweet William, Telanth-rera, Masambtyanthem.

ORNAMENTALS, SHRUBS AND TREES (As listed on side panel): Apply 2 1/2 cups (1 lb.) MEACLE WEED PREVENTER per 200 sq. ft. uniformly to soil around the base of plants in late full or early spring. May be applied following transplanting into well prepared soil. For established plantings apply proper cultivation to remove existing weeds. SEE SIDE PANEL FOR LIST OF TREES AND SHRUBS THIS PRODUCT MAY BE APPLIED AROUND.

1/1/2/1!

VEGETABLE CROPS: Any application of MIRACLE WEED PREVENTER should be made prior to weed seed germination. Apply at seeding to snap beans, broccoli, brussels sprouts, cabbage, caulillower, collards, garlic, kale, mustard greens, onions, potatoes, (Southern peas) turnip greens, turnips.

Apply to seeded cucumbers, melons (cantaloupe, watermelons, and honeydew), squash and head lettuce about 4 weeks after seeding, but no later than 6 weeks. For leaf lettuce, make a single application 1-3 weeks after emergence. When transplanting broccoli, brussels sprouts, cabbage, cauliflower, eggplant, peppers, onions, and sweet potatoes, apply immediately after transplanting.

TOMATOES: Apply only to well established plants, 4 to 6 weeks after transplanting, when conditions are favorable.

STRAWBERRIES: Apply to strawberries following transplanting or on established strawberry plants prior to weed seed germination or after clean cultivation. On established plants, do not apply after first bloom.

REPLANTING: Replanting crops other than those included on the label in DACTHAL treated soil within 8 months of application may result in crop injury. If replanting is required because of early crop failure, the planting of onions, seeded melons, potatoes, tomatoes, eggplants, or peppers at this time may result in crop injury. However, all crops listed on this label may be planted following harvest of a crop treated with this product.

STORAGE: Do not store near food or foodstuffs. Store in the tightly closed original container in a cool, dry preferably locked storage area, out of reach of children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of rspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Distributed By DRAGON CORPORATION Roanoke, Va 24019-0311

EPA Reg. No. 16-42

EPA Esí. 16-VA-1

Form #L-42

Α

