

1042

FOR USE
 as, ...
 se Leafhoppers, Spider
 Flies, Mealybugs, etc.
 For Japanese Beetle
 Four-lined Leaf Bugs,
 s, Scale Crawlers, and
 gallon of water.
 per gallon of water on
 s and Mealybugs. Apply
 for the following vege-
 (Aphids and Leafhoppers),
 (Aphids), Beans (Aphids,
 Japanese Beetles), Cu-
 and Spider Mites). IM-
 Beans, Cucumbers and
 Do not apply to Head
 Kale within 7 days of
 lettuce within 14 days of
 within 3 days of harvest
 treated crops are to be

BAT NO



CYTHON
PREMIUM GRADE
MALATHION
50% EMULSIFIABLE LIQUID
 *Trademark of American Cyanamid Company

KILLS:

- APHIDS
- RED SPIDERS
- MEALYBUGS
- BAGWORMS
- WHITE FLIES
- FLEAS
- AND MANY OTHER PESTS

ACTIVE INGREDIENTS:		
Malathion*	50.0%	NET WEIGHT ONE LIQUID GALLON
Heavy Aromatic Naphtha	45.0%	
INERT INGREDIENTS	5.0%	
*O, O-dimethyl phosphorodithioate of diethyl mercaptosuccinate		

...
 ...
 body surfaces including the inside of the ears. S
 treatment may be necessary in 10 days
 Hogs should be kept out of sun and wind for
 hours after treatment.
Sheep and Goats:
 Lice, Ticks and Keds: Dilute 4 ounces of Kill-Ko
 Cythion with 3 gallons of water. Treat animals
 oughly. Repeat application after 2 or 3 weeks if ne
 Do not apply to milk goats.
FLEAS IN DOG KENNELS, YARDS, AND ON DOGS:
 infested areas with 10 tablespoons (5 ounces) Co
 trate in 1 gallon water at rate of 1 gallon mixtu
 1000 square feet. Remove manure and debris &
 Spraying. Wet dogs with 2 tablespoons Concentra
 1 gallon water. Animals should be closely examine
 fleas before retreating. Repeat treatment not more
 than once every 2 or 3 weeks. Do not treat dogs
 one month after

CAUTION: Avoid breathing any mist. Avoid oc
 contact with eyes. Wash thoroughly with water after use. Harm
 followed by a liberal amount of food and food
 kept out of reach of animals. Do not use in
 to avoid contact with animals. Do not use in
 animals. Do not use in areas where animals
 with fresh water. Do not use in areas where
 animals. Do not use in areas where animals
 animals. Do not use in areas where animals
 animals. Do not use in areas where animals
 animals. Do not use in areas where animals

5% MALATHION DUST

READ ENTIRE LABEL CAREFULLY
CAUTION: WASH HANDS THOROUGHLY AFTER USE
WITH FEDERAL AND STATE REGULATIONS

DIRECTIONS FOR USE

For control of hornflies on beef and dairy cattle, apply 3 tablespoons of this form of dust per animal. Apply the dust on the back and neck of the animal at least 10-14 days intervals. Do not use on dairy cattle later than 7 hours before milking and do not use on water troughs.

Begin dusting when plants are small. Make application every 7-10 days. Keep all parts of plants thoroughly and evenly dusted, especially new growth. Quantity of dust to use will vary with size of plant, dusting apparatus, etc. For crops listed below approximate application rates are: 1/2 lb. per 1000 sq. ft. is usually sufficient, except on beans, where 1 lb. per 1000 sq. ft. is usually sufficient, and on peas where 1/2 lb. per 1000 sq. ft. is usually sufficient. Remove dust deposits from food by washing before consuming.

BEANS: For control of aphids, Japanese Beetle, leafhopper, Mexican Bean Beetle. Make no application to beans later than 7 days before harvest.

POTATOES: For control of aphids. Make no application on potatoes later than 3 days prior to harvest.

COLE CROPS: For control of aphids, imported cabbage worm, and cabbage looper on cabbage, broccoli, kale, mustard and turnip. Make no application to cabbage, kale and mustard within 7 days of harvest or to broccoli and turnip within 3 days of harvest.

PEAS: For control of pea aphid. Make no application later than 7 days before harvest. Peas should not be treated within 7 days of harvest if vines are to be fed to livestock.