

NOK^{*}OUT

5% SEVIN^{*} DUST

FOR CONTROL OF JAPANESE BEETLE, FLEA BEETLE, MEXICAN BEAN BEETLE,
TOMATO HORNWORM, CORN EARWORM AND CERTAIN OTHER LISTED
INSECTS ATTACKING VEGETABLES, GRAPES, ORNAMENTALS AND SHRUBS.

ACTIVE INGREDIENT	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate)	5%
INERT INGREDIENTS	95%
	TOTAL 100%

*Registered Trademark of Union Carbide Corporation

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK/SIDE PANELS FOR ADDITIONAL CAUTIONS.

4 LBS. NET WT.

ACCEPTED

JUL 11 1986

U.S. Environmental Protection Agency,
Federal Insecticide, Fungicide, and Rodenticide Act,
registered under the pesticide
label under EPA Reg. No. 16-69

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. **DO NOT** take internally. Wear regular long sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

GENERAL INFORMATION

MOR-OUT 5% SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. **DO NOT** use with lime or other alkaline material. May be applied up to harvest except where noted. **DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS — For control of Mexican bean beetles, leafhoppers, flea beetles, apply approximately ½ lb./1000 sq. ft. **DO NOT** feed treated bean vines to dairy or meat animals.

LETTUCE — For control of lygus and stink bugs, apply approximately 1 lb./1000 sq. ft. **DO NOT** apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN — Use approximately ¾ lb./1000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS — For control of Colorado potato beetles, leafhoppers, and flea beetles, apply approximately ½ lb./1000 sq. ft. For control of tomato fruitworms, fall armyworm, tomato hornworm, European corn borer, stink bug and lygus bugs, apply approximately ¾ lb./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH — For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles, apply approximately ½ lb./1000 sq. ft.

GRAPES — For control of leafhoppers, Japanese beetle, grape berry moth, apply approximately 1 lb./1000 sq. ft.

STRAWBERRIES — For control of meadow spittlebug, leaf roller, apply approximately 1 lb./1000 sq. ft. **DO NOT** apply within 1 day of harvest.

ROSES, AZALEAS, SHADE TREES AND OTHER ORNAMENTALS — For control of Japanese beetles, periodical cicada, aphid, tent caterpillars, lace bugs, boxwood leaf miner, apply approximately 1-1½ lbs./1000 sq. ft. Injury to tender foliage may occur if plants are wet when dusted or in the presence of high humidity. **DO NOT** use on Boston Ivy.

DOGS, CATS — For control of fleas and brown dog ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. **DO NOT** treat kittens or puppies under 4 weeks old. **DO NOT** apply more than once weekly. Do not use on pregnant dogs.

STORAGE AND DISPOSAL

Storage: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs. **Pesticide disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container disposal:** Do not reuse empty container. Discard container in trash.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN is a registered trademark of Union Carbide Corporation for carbaryl insecticide.

EPA Reg. No. 16-66-

EPA Est. 16-1A-1

Manufactured by

DRAGON CHEMICAL CORPORATION
Roanoke, Virginia 24019

