

FRONT PANEL

DRAGON
LIME SULPHUR
SOLUTION

NET CONTENTS

TEST: 52° Baume at 50° .

Controls: Overwintering Scale And Diseases

ACTIVE INGREDIENT	BY WT.
Calcium Polysulphide.....	29%
INERT INGREDIENTS.....	71%
Total	100%

DO NOT STORE BELOW 5° F.

CAUTION: Keep out of Reach of Children. See side/back panels
for additional cautions. (1)

DRAGON CHEMICAL CORPORATION
ROANOKE, VA.
MAKERS OF BLUE DRAGON
USDA Reg. No. 16-X

Inert Ingredients: To be supplied by Miller Chemical & Fer-
tilizer Co.

BACK PANEL

DIRECTIONS FOR USE

PEACHES AND PLUMS: For control of LEAF CURCER and other leaf-eating insects. Apply when peaches and plums are dormant. Use at rate of 2½ gallons per 100 or 1 pint per gallon of water.

FRUIT: Pre-harvest spray: For control and prevention of BROWN ROT. Use 2 to 2½ quarts per 100 gallons of water. Suggested for use after hail damage or close to harvest and has the advantage over Wettable Sulfur in that spray residue is much less noticeable. Some leaf injury may result during hot weather. Can be used within 1 day of harvest.

APPLES: For control of SCALE, LEAF SPOT, POWDERY MILDEW: Apply when blossoms are breaking and first showing pink. Use at rate of 2½ gallons per 100 or 1 pint to 5 gallons, or 6 tablespoonfuls per gallon of water. Additional cover sprays, pink, petal fall and first cover should be applied at 2 gallons per 100 or 5 tablespoonfuls per gallon.

FRUIT AND ORNAMENTALS: For control of HARD or ARMORED SCALE, Scurfy scale, San Jose scale, Euonymus scale, Pine needle scale, Juniper scale on most Ornamental and Fruit trees such as Apple, Peach, Lilac, Ash, Poplar, Dogwood, Elm, Birch, Willow, Euonymus, Pines and Junipers: Spray when trees are fully dormant and use at rate of 10 to 12 gallons per 100 or 1 pint to each gallon of water.

CURRENTS AND GOOSEBERRIES: For control of ANTHRACNOSE: Use at rate of 1 gallon to 40 to 50 gallons of water. Make the first application when buds are breaking. Repeat in 10 to 15 days. Two to three additional sprays may be necessary. Use 2 to 2½ qts. per 100 gallons of water applied at 10 to 15 day intervals, if necessary, for control.

RASPBERRIES: For control of RUST, ANTHRACNOSE, CANE AND SPUR BLIGHT: Spray as delayed dormant when buds begin to break and show silver. Use at rate of 10 to 12 gallons per 100 or 1 pint per gallon of water.

ROSES: For control of ANTHRACNOSE: Apply when roses are dormant. Use 10 to 12 gallons per 100 or 1 pint per gallon of water. For control of POWDERY MILDEW: First cover spray should be applied when buds break in spring. Use at rate of 2 gallons to 100 or 1 pint to 6 gallons of water. Two to three applications at 7 to 10 day intervals are desirable.

BOXWOOD: For control of CANKER: Remove infected branches down to healthy wood. Use Lime Sulfur Solution 2 gallons to 100 or 1 pint to 6 gallons of water. Make applications as follows: 1st, as soon as dead leaves have been cleaned up in the spring; 2nd, when new growth is about half completed; 3rd, after spring growth is completed; 4th, after fall growth is completed.

CAUTION

Wash hands after use. DO NOT spray on painted surfaces to avoid staining. Keep bottle tightly closed when not in use. When container empty, immediately wash thoroughly and destroy. Never re-use. ②

For best results, Lime sulfur Solution should be used alone. Use in cool weather, early spring and fall sprays. Lime Sulfur Solution should not be used within 10 to 14 days of an oil spray or when temperatures are high.

NOTICE

Buyer assumes all risks for personal injury or property damage.

③

CORRECTIONS TO BE INSERTED ON ATTACHED LABEL COPY

DRAGON LIME SULFUR SOLUTION

1. SEE ATTACHED TYPE SIZE SHEET FOR REQUIREMENTS.
2. INSERT the word "is" as indicated.
3. INSERT the following copy: WARRANTY

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.

GTA/vt
3/28/67