

**DRAGON LIME SULFUR SOLUTION**

**FUNGICIDE-INSECTICIDE**

**A Dormant Clean Up For Fruit Trees, Ornamentals, Roses, Shrubs  
Controls Overwintering Insects and Diseases As Listed.**

<b>ACTIVE INGREDIENT:</b>	<b>BY WT.</b>
Calcium Polysulfide.....	29.0%
<b>INERT INGREDIENTS:</b> .....	<b>71.0%</b>
<b>TOTAL:</b> .....	<b>100.0%</b>

Test 32° Baume at 60° F.

**Keep Out of Reach of Children**

**DANGER**



See back panel for additional precautions

NET CONTENTS

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**DANGER:** Fatal if swallowed, inhaled or absorbed through skin. This product is corrosive to flesh because of its caustic alkaline nature and it causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking and using tobacco. Remove contaminated clothing and wash before reuse.



**FIRST AID**

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:  
CHEMTREC AT: 1-800-424-9300

**IF INHALED:**

Move person to fresh air.  
If person is not breathing, call 911, or an ambulance, then give artificial respiration, preferably by mouth to mouth, if possible.  
Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING**

Take off contaminated clothing.  
Rinse skin immediately with plenty of water for 15 to 20 minutes.  
Call a poison control center or doctor for further treatment advice.

**IF IN EYES**

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.  
Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.  
Call a poison control center or doctor for further treatment advice.

**IF SWALLOWED**

Call Poison Control Center or Doctor immediately for treatment advice.  
Have person sip a glass of water if able to swallow.  
Do not induce vomiting unless told to do so by the poison control center or doctor.  
Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. *You may also contact 1 - 800 - 222- 1222 for emergency medical treatment information.*

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

**PHYSICAL OR CHEMICAL HAZARDS**

MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.


**STOP!** Read the entire label first, observe all precautions and follow directions carefully.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE & DISPOSAL**

**Storage:** Store only in original container, in a cool dry place inaccessible to children and pets and away from feed and foodstuffs.  
**Disposal: If Empty:** Do Not reuse this container. Place in trash or offer for recycling if available.  
**If partly filled:** Call your local solid waste agency or call CHEMTREC AT 1-800-424-9300 for disposal instructions. Never place unused product down any indoor or outdoor drain.



**GENERAL INFORMATION: FOR OUTDOOR HOMEOWNER USE ONLY.** For best results, Lime Sulfur Solution should be used alone. Use in cool weather - early Spring and Fall sprays. Apply sufficient spray to uniformly wet all plant surfaces. Lime Sulfur Solution should not be used within 10 to 14 days of an oil spray, nor when temperatures are high.

**APPLES:** For control of Leaf Spot, Powdery Mildew - Apply when blossoms are breaking and first showing pink. Use at rate of 1 pint to 5 gallons, or 6 Tbs. per gallon of water. Additional cover sprays, pink, petal fall and first cover should be applied at 5 Tbs. per gallon. For control of Hard and Armored Scales, Scurly Scale and San Jose Scale - Spray when trees are fully dormant and use at rate of 1 pint to each gallon of water.

**PEACHES:** Pre-harvest spray: For control or prevention of Brown Rot - Use 1 1/4 Tbs. per gallon of water. Suggested for use after hail damage or close to harvest, and has the advantage over wettable sulfur in that spray residue is much less noticeable. Some leaf injury may result during hot weather. Can be used within 1 day of harvest. For control of Leaf Curl, Hard and Armored Scales, Scurly Scale and San Jose Scale - Spray when trees are fully dormant and use at rate of 1 pint to each gallon of water.

**PLUMS:** For control of Leaf Curl and San Jose Scale - Apply when plums are dormant. Use 1 pint per gallon of water.

**CURRENTS, GOOSEBERRIES:** For control of Anthraxnose - Make the first application when buds are breaking. Repeat in 10 to 15 days. Two to three additional sprays may be necessary applied at 10 to 15 day intervals, if necessary, for control.

**RASPBERRIES:** For control to Rose Scale, Anthracnose, Cane and Spur Blight - Spray as delayed dormant when buds begin to break and show silver. Use at rate of 1 pint per gallon of water.

**ORNAMENTALS, SHRUBS, TREES:** For control of Hard and Armored Scales, Scurly Scale, San Jose Scale, Euonymus Scale, Pine Needle Scale and Juniper Scale on Lilac, Ash, Poplar, Dogwood, Elm, Birch, Willow, Euonymus, Pines and Junipers - Spray when trees are fully dormant and use at rate of 1 pint to each gallon of water.

**ROSES:** For control of Anthraxnose - Apply when roses are dormant. For control of Powdery Mildew - First cover spray should be applied when buds break in Spring. Two to three applications at 7 to 10 day intervals are desirable.

**BOXWOOD:** For control of Canker - Remove infected branches down to healthy wood. Make applications as follows: 1st, as soon as dead leaves have been cleaned up in the Spring; 2nd, when new growth is about half completed; 3rd, after Spring growth is completed; 4th, after Fall growth is completed.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

EPA Reg. No. 16-72

Distributed by  
DRAGON CHEMICAL CORPORATION  
P O Box 7311, Roanoke, Virginia 24019  
www.dragonchemical.com

