

16-710

08/04/1971

1/2

Thrips (exposed): Apply 4 standard teaspoonfuls of concentrate per gallon of water when insects or their damage appears. Re-apply as necessary when dried powder no longer is apparent. Completely cover all parts of plants, including underside of leaves and branches. Repeat application in 7 to 10 days, as necessary.

VEGETABLES: Beans, Corn, Pepper, Tomato. To control Potato Beetle, Mexican Bean Beetle, Flea Beetles, Japanese Beetle, Army Worms, Tomato Hornworm and Tomato Fruit Worm: Mix 4 to 6 standard teaspoonfuls in each gal. of water. Use 1 gal. of diluted spray to 430 sq. ft. Apply when insects or damage appears and repeat in 7 to 10 days, as necessary. Do not apply to Beans, Peppers, Potatoes and Tomatoes within 1 day of harvest. Do not apply to Corn within 7 days of harvest. Do not feed treated corn forage to Dairy Cattle or Livestock.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do not breathe spray. Do NOT take internally. Avoid prolonged or repeated breathing of spray. Skin contact may be harmful. Avoid repeated or prolonged contact. Wash hands before eating. Wash thoroughly after using. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOODSTUFFS: Avoid contamination of feed, feeding troughs and watering receptacles.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS AND CAUTIONS stated on this label.

SMALL MEASUREMENTS

One oz. 2 standard tbsp. or 6 standard tsp.
One cup 8 oz. or 16 standard tbsp.

\$2.69



NET CONTENTS
1 PINT

LIQUID SEVIN® SPRAY

CONTROLS CERTAIN INSECTS ON
ORNAMENTALS, VEGETABLES

| | |
|--|--------|
| ACTIVE INGREDIENTS | By Wt. |
| Carbaryl (1-naphthyl N-methylcarbamate)..... | 21.5% |
| INERT INGREDIENTS | 78.5% |

USDA REG. NO. 16-76

Trade Mark of UCC for 1-naphthyl N-methylcarbamate

CAUTION: Keep out of Reach of Children.
See other panels for additional cautions.

Manufactured for

DRAGON CHEMICAL CORP., ROANOKE, VIRGINIA

LIQUID SEVIN SPRAY

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON LIQUID SEVIN SPRAY mixes with water and is ideal for use in home garden sprayers. Will not clog spray nozzles. LIQUID SEVIN controls most insects that damage VEGETABLES, ORNAMENTAL TREES and SHRUBS such as COLORADO POTATO BEETLE, FLEA BEETLE, MEXICAN BEAN BEETLE, JAPANESE BEETLE and LEAFHOPPER. LIQUID SEVIN can be used on many Vegetable Crops close to harvest.

COMPATIBILITY

DRAGON LIQUID SEVIN can be mixed with most of the wettable powder insecticides and fungicides. It can also be mixed with many other liquid insecticides; however, before using with other liquid materials, make a small batch and test to make sure they are physically compatible. LIQUID SEVIN does not control mites; use with a suitable miticide such as Kelthane. This product is unstable when mixed with lime, Bordeaux, Lime and Sulphur, Casein-lime stickers or other alkaline materials. Some injury may occur when used on tender foliage when prolonged wet periods or during periods of high humidity. Do not use on Boston Ivy.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USING. CLOSE CONTAINER IMMEDIATELY AFTER USE TO PREVENT MATERIAL FROM EVAPORATION OR DRYING. PROTECT FROM TEMPERATURES BELOW 32° F. Pour recommended amount of this material into nearly filled spray tank, or pre-mix in a bucket. Agitate during filling and spraying.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: Rose, Chrysanthemum, Arbor Vitae, Azalea, Juniper, Elm, Maple, Oak, Dogwood and Pines. To control Japanese Beetle, Rose Aphid, Periodical Cicada, Mealy Bugs, Leafhoppers, Lace Bugs, Flea Beetles, Aphid, Bagworm, Roseslug, Scale Insects,

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

ACCEPTED
AUG 4 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. *16-76* SUBJECT
TO ATTACHED COMMENTS.

16-76

16-76

2/2