

November 18, 1971

EPA REG. NO. 16-76

DRAGON
LIQUID
SEVIN SPRAY

ACCEPTED
16 76

FRONT PANEL

DRAGON
LIQUID
SEVIN*SPRAY

NET CONTENTS
1 Pint

CONTROLS CERTAIN INSECTS ON ORNAMENTALS, VEGETABLES, LAWNS, STRAWBERRIES.

ACTIVE INGREDIENTS	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate).....	21.5%
INERT INGREDIENTS.....	78.5%
	<u>Total 100.0%</u>

*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See other panels for additional cautions.

DRAGON CHEMICAL CORPORATION

Roanoke, Virginia 24019

EPA REG. NO. 16-76

BACK PANEL

LIQUID SEVIN SPRAY

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON LIQUID SEVIN SPRAY mixes with water and is ideal for use in home garden sprayers. Will not clog spray nozzles. LIQUID SEVIN controls most insects that damage TURF, VEGETABLES, STRAWBERRIES, ORNAMENTAL TREES and SHRUBS such as COLORADO POTATO BEETLE, FLEA BEETLE, MEXICAN BEAN BEETLE, JAPANESE BEETLE and LEAFHOPPER.

COMPATIBILITY

DRAGON LIQUID SEVIN can be mixed with most of the wettable powder insecticides and fungicides. It can be mixed with many other liquid insecticides; however, before using with other liquid materials, make a small batch and test to make sure they are physically compatible. LIQUID SEVIN does not control mites; use with a suitable miticide such as Kelthane. This product is unstable when mixed with lime, Bordeaux, Lime and Sulphur, Casein-lime stickers or other alkaline materials. Some injury may occur when used on tender foliage when prolonged wet periods or during periods of high humidity. Do not use on Boston Ivy.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USING. CLOSE CONTAINER IMMEDIATELY AFTER USE TO PREVENT MATERIAL FROM EVAPORATION OR DRYING. PROTECT FROM TEMPERATURES BELOW 32°F. Pour recommended amount of this material into nearly filled spray tank, or pre-mix in a bucket. Agitate during filling and spraying.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: Rose, Chrysanthemum, Arbor Vitae, Azalea, Juniper, Elm, Maple, Oak, Dogwood and Pines. To control Japanese Beetle, Rose Aphid, Periodical Cicada, Mealy Bugs, Leafhoppers, Lace Bugs, Flea Beetles, Aphid, Bagworm, Roseslug, Scale Insects, Thrips (exposed): Apply 4 standard teaspoonfuls of concentrate per gallon of water when insects or their damage appears. Re-apply as necessary when dried powder no longer is apparent. Completely cover all parts of plants, including underside of leaves and branches. Repeat application in 7 to 10 days, as necessary.

VEGETABLES: Beans, Corn, Pepper, Tomato. To control Potato Beetle, Mexican Bean Beetle, Flea Beetles, Japanese Beetle, Army Worms, Tomato Hornworm and Tomato Fruit Worm: Mix 4 to 6 standard teaspoonfuls in each gal. of water. Use 1 gal. of diluted spray to 430 sq. ft. Apply when insects or damage appears and repeat in 7 to 10 days, as necessary. Do not apply to Beans, Peppers, Potatoes and Tomatoes within 1 day of harvest. Do not apply to Corn within 7 days of harvest. Do not feed treated corn forage to Dairy Cattle or Livestock.

A

LAWNS: To control Earwigs, Mosquitoes, Leafhoppers, Ants: It is necessary to put the chemical where the insects live. Mow lawn, remove cuttings, and apply immediately after rain or good watering 2/3 pt. in 15 to 20 gallons of water for each 500 sq.ft. of established lawn area. Use hose-on sprayer or knapsack sprayer and apply sufficient water for good penetration into turf. Wash in with additional water to wet the soil to a depth of 1 inch. Repeat in 2 to 3 weeks, if necessary.

STRAWBERRIES: For control of Meadow Spittlebug, Strawberry Leaf Roller and Strawberry Weevil: Use 2 pts. per 30 gallons of water to spray 1/3 acre or 2 standard tablespoonfuls per gallon of water and apply to 430 sq. ft. Apply early spring for Spittlebug and repeat in 5 to 7 days, as needed. Do not apply within 1 day of harvest.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do not breathe spray. Avoid skin contact. Skin contact may be harmful. Wash thoroughly after using. Take shower or bath after work. Wear regular longsleeved work clothing. Change to clean clothing daily.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOODSTUFFS: Avoid contamination of feed, feeding troughs and watering receptacles.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SMALL MEASUREMENTS

One oz. = 2 standard tbsp. or 6 standard tsp.
One cup = 8 oz. or 16 standard tbsp.