

MAY 29, 1972

EPA REG. NO. 16-79

ACCEPTED  
JUN 30 1972  
16 79

DRAGON  
WEED BLASTER

FRONT PANEL

DRAGON

WEED BLASTER

NET CONTENTS  
Quarts & Gallons

For General Weed Control Along Sidewalks, Driveways, Fences, Around  
Buildings, And On Similar Non-Crop Areas.

ACTIVE INGREDIENTS	% By Wt.
Sodium Cacodylate.....	6.7%
Dimethylarsinic Acid (Cacodylic Acid).....	1.1%
INERT INGREDIENTS.....	92.2%
Total	<u>100.0%</u>

Total Arsenic (as elemental) all in water soluble form 3.78%.  
Product contains 0.62 lbs. Cacodylic Acid equivalent per gallon.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back panel for additional  
cautions.

INERT INGREDIENTS: See attached letter from the Ansul Company.

DRAGON CHEMICAL CORPORATION

Roanoke, Virginia 24019

EPA REG. NO. 16-79

## BACK PANEL

### DRAGON

### WEED BLASTER

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

#### GENERAL INFORMATION

DRAGON WEED BLASTER is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It produces top-kill only so repeat applications are required for season long weed control of perennials.

#### MIXING INSTRUCTIONS

DRAGON WEED BLASTER must be thoroughly dissolved. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

#### CONTAINER DISPOSAL

Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

#### DIRECTIONS FOR USE

##### NON-CROP

DRAGON WEED BLASTER is useful for general weed control on drainage ditchbanks and rights-of-way, along sidewalks, driveways and fences, around buildings and ornamentals, and on similar non-crop areas. Apply with either tank type or hose-end sprayer, adjusting for coarse spray to prevent drift which could damage nearby desirable plants. Follow directions in the table below. Spray unwanted vegetation thoroughly to just short of run-off. If regrowth occurs, reapply as required.

Mixing Directions for Both Tank Type and Hose-End Sprayers:

Non-Crop:	<u>Amount of DRAGON WEED BLASTER</u>	<u>AMOUNT of Water</u>
	5 to 10 fluid ounces	1 gallon
	10 to 20 fluid ounces	2 gallons
	1½ to 3 pints	5 gallons
	1 to 2 quarts	6 gallons
	1½ to 3 quarts	10 gallons
	2 to 4 quarts	15 gallons
	1 to 2 gallons	25 gallons

DO NOT APPLY DIRECTLY, OR ALLOW SPRAY TO COME IN CONTACT WITH FOLIAGE, GREEN BARK, GRAFTED UNIONS, OR SCUFFED, DAMAGED OR BROKEN BARK WHEN SPRAYING AROUND TREES AND ORNAMENTALS.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES. DO NOT GRAZE TREATED AREAS TO LIVESTOCK.

#### LAWN RENOVATION

Mow lawn to be renovated about 1 inch high before treatment. Apply with either tank type or hose-end sprayer. Follow directions in the table below. Cover foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or uncovered foliage, may require a second treatment. Reapply five days later if green areas remain.

Mixing Directions for Both Tank Type and Hose-End Sprayers:

<u>AREAS TO BE TREATED</u>	<u>AMOUNT OF DRAGON WEED BLASTER</u>	<u>AMOUNT OF WATER</u>
1000 sq. ft.	2½ pints	4 gallons
1600 sq. ft.	2 quarts	6 gallons
2400 sq. ft.	3 quarts	10 gallons
3200 sq. ft.	1 gallon	15 gallons

When foliage is all brown, matted areas should be raked and bare areas scarified. About five days after last treatment, water thoroughly and then seed at rate recommended for new lawns. Seed must be raked into soil. Keep seeded areas well watered.

KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL AFTER RAIN OR SPRINKLING FOLLOWING TREATMENT. DO NOT TRACK FROM TREATED TO UNTREATED AREAS.

CAUTION

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE

If taken internally, induce vomiting and call physician at once.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.