

16-91  
5/6/71

16-91 5/6/71

pg 182

acceptance stamp on  
p.2

4 LBS. NET WT.



**DRAGON**

**1%**  
**ROTENONE**  
**DUST**

CONTROLS CERTAIN INSECTS ON VEGETABLES and FLOWERS.

ACTIVE INGREDIENTS	BY WT.
Rotenone .....	1.00%
Other Cube Resins .....	1.53%
INERT INGREDIENTS .....	97.47%
	<b>TOTAL 100.00%</b>

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

SEE BACK SIDE PANELS FOR ADDITIONAL CAUTIONS.

USDA Reg. No. 16-91

MANUFACTURED BY  
**DRAGON CHEMICAL CORPORATION**  
ROANOKE, VIRGINIA 24019,

45-PA

4-1071

ACCEPTED

MAY 6 1971

UNDER THE FEDERAL  
FUNGICIDE AND RO  
FOR ECONOMIC PE  
BY (UNIT NO. ...)  
TO ATTACHED CERTIFICATE

16-91

262



**DRAGON**

## 1% ROTENONE DUST OR SPRAY

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS, AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

### DIRECTIONS FOR USE

Do not mix the contents of this package with any other material but apply to the plant with a dust gun just as it comes from the package. For average dusting, one pound will treat approximately 3630 sq. ft. of row. If a liquid spray is desired, add one pound of Dragon 1% Rotenone Dust to four gallons of water and apply to the same area. Add the dust gradually to a small amount of water to form a thin paste and then slowly add the paste to the rest of the water. Spray with a good liquid spray gun which will force the mist on the underside of the leaves. Shake sprayer during application to keep mixture suspended.

**MEXICAN BEAN BEETLE**—Dust or spray leaves as soon as the adult beetles appear. Several hours are required for control. Repeat the application when necessary.

**BEAN LEAF BEETLE**—Apply same as for MEXICAN BEAN BEETLE.

**CUCUMBER BEETLES**—On BEANS, CUCUMBERS, PUMPKINS, SQUASH—Use 1 lb. to treat approximately 2900-3600 sq. ft.—begin early and repeat as necessary.

**VEGETABLE WEEVIL**—On BROCCOLI, CABBAGE, CAULIFLOWER, CARROTS, COLLARDS, KALE, TURNIPS, MUSTARD GREENS—apply to foliage as needed.

**ASPARAGUS BEETLE**—Dust as soon as larvae appear.

**CABBAGE WORMS**—Also on BROCCOLI and CAULIFLOWER—apply at the rate of 1 lb. to approximately 2200 sq. ft. as soon as worms appear—repeat in about two weeks.

**POTATO BEETLE**—Use 1 lb. to treat approximately 2900-3600 sq. ft. as soon as larvae begin feeding—repeat when needed.

**TOBACCO FLEA BEETLE**—In plant beds—dust through cheese cloth—8 ounces of DRAGON 1% ROTENONE DUST per 100 yards.

**PEPPER MAGGOT**—Apply on foliage as needed.

**FLEA BEETLES**—On OKRA, SWISS CHARD, TURNIPS—apply on foliage as needed.

**SQUASH VINE BORER**—On PUMPKINS, SQUASH—apply once or twice a week to stems and vines at base of plants.