

5399633

JUL 18 1989

Mr. James C. Wilson
Dragon Corporation
P.O. Box 7311
Roanoke, VA 24019

Dear Mr. Wilson:

Subject: 1% Rotenone Dust
EPA Registration No. 16-91
Confidential Statement of Formula and Label Dated June 1,
1989 Submitted in Response to Registration Standard
Requirements

The ingredient statement on the front panel should include another
line showing the total active and inert ingredients.

The statement "Natural Organic" is an implied safety claim and must
be removed from the label at the next printing of labels--no more than six
(6) months from the date stamped on this letter.

The Confidential Statement of Formula dated June 1, 1989 will be
kept in the file for this product. Further revisions will be required if
changes are made in labels for Manufacturing Use Concentrates supplying
the active ingredients for this product.

Sincerely yours,



Phil Hutton
Product Manager (18)
Insecticide-Rodenticide Branch
Registration Division (47505C)

BEST AVAILABLE COPY

52342:1:Schroeder;C.Disk;KEVRIC;06/11/91;CL:WO:CL

CONCURRENCES

SYMBOL	SURNAME	DATE					

Front panel



DRAGON

1% ROTENONE
DUST

ORGANIC INSECT CONTROL.

(illustrations of Mexican bean beetle, cucumber beetle,
cabbageworm, squash vine borer)

ACTIVE INGREDIENTS

Rotenone 1.00%
Other Associated
Resins 1.53%
INERT INGREDIENTS 97.47%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side/back panels for
additional cautions.
NET WT 1 LB , 4 LB

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL 18 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

16-91

[Faint rectangular stamp or mark]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL & STATE REGULATIONS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid inhalation of dust. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. Wear long-sleeved work clothing. Take shower or bath after work. Wash hands before eating or smoking.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not mix the contents of this package with water. Apply to the plant with a garden duster or other appropriate applicator.

RATE OF APPLICATION Apply at a rate to give a thin, uniform covering to the upper and lower leaf surfaces. One pound treats approximately 400 linear feet of row.

BEANS To control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Flea Beetle, Thrips, Twelve-spotted Beetle, dust as the adult beetles appear. Repeat the application as necessary.

CUCUMBERS, PUMPKINS, SQUASH Apply as directed above to control Cucumber Beetle. Begin early & repeat as necessary.

BROCCOLI, CABBAGE, CAULIFLOWER, CARROTS, COLLARDS, KALE, TURNIPS, MUSTARD GREENS Apply as directed above to control Vegetable Weevil, Imported Cabbageworm.

ASPARAGUS Apply as directed above, as soon as larvae appear, to control Asparagus Beetle.

SWEET CORN To control European Corn Borer, make the first application as soon as young borers appear on leaves. Start applications in late May or early June for first brood, and in late July or early August for second brood. Make four applications at exactly five day intervals. Direct the dust downward into the whorls or crowns and leaf axils.

POTATOES Apply as directed above to control Potato Beetle, as soon as larvae begin feeding.

PEPPERS Apply as directed above to control Pepper Maggots

FLOWERS & SHRUBS To control many external leaf-chewing insects, apply as soon as the insects appear, and repeat at weekly

BEST AVAILABLE COPY

intervals as long as the insects continue to appear. Examples of the types of insects that may be controlled are: cabbage looper on carnation, chrysanthemum and geranium; flea beetle on petunia, primrose; Imported Cabbage Worm on nasturtium, mignonette and sweet alyssum; rose chafer on rose and peony.; Asiatic garden Beetle on petunia, phlox and rose.

FOR DOGS & CATS To control lice and fleas, apply to all parts of the animal, rubbing it thoroughly down to the skin. Special attention should be given to the neck and base of the tail. Repeat in 10 days. Do not use on puppies or kittens less than 4 weeks old.

Storage Store only in original container, in a cool, dry place inaccessible to children and pets. Keep away from feed & foodstuffs. **Pesticide disposal** Securely wrap original container in several layers of newspaper and discard in trash. **Container disposal** Do not re-use empty container; discard in trash.

NOTICE Buyer assumes all risks of use, storage or handling of this material not in accordance with **DIRECTION & CAUTIONS** stated on this label.

EPA Reg. No. 16-91

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

EPA Est 16-VA - 1

rotenone 9/89

POST OFFICE