


NET CONTENTS
1 GALLON



72% CHLORDANE SPRAY

CONTROLS: ANTS, GRASSHOPPERS, BEETLE GRUBS, NIGHT CRAWLERS, TERMITES, ROACHES, AND CERTAIN OTHER INSECTS.

ACTIVE INGREDIENTS	By Wt.
Technical Chlordane*	72%
Petroleum Distillate	23%
INERT INGREDIENTS	5%
	100%

*Equivalent to 43.2% Octachloro-4,7-Methanotetrahydroindane and 28.8% related compounds

WARNING: EPA REG. NO. 16-96
KEEP OUT OF REACH OF CHILDREN.
for additional warnings.

See back panel

DRAGON CHEMICAL CORP., ROANOKE, VIRGINIA 24019
Packed For

72% CHLORDANE SPRAY

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

DIRECTIONS FOR USE

DO NOT store diluted spray. Make new dilution for each use.

LAWNS

INSECTS: JUNE BEETLE GRUBS, WHITE GRUBS, JAPANESE BEETLE GRUBS, SOD WEBWORMS, ANTS, MOLE CRICKETS, CUTWORMS, GRASSHOPPERS, CHIGGERS — Spray thoroughly at 1 Tablespoonful in 1 gallon water per 150 sq. ft. area (16 Tablespoonfuls in 15 gals. water per 2400 sq. ft.)

NIGHT CRAWLERS: Apply 4 ounces (8 Tablespoonfuls) in 7½ to 15 gals. water to 1000 sq. ft. of lawn area. Water thoroughly after application.

MOLES: The above treatment for insects may be used to reduce damage caused by moles which feed on such insects.

GARDENS

CUTWORMS, STRAWBERRY ROOT WEEVIL (BFACHYRHINUS), CAMELIAS, AZALEAS, RHODODENDRONS, PRIMROSES AND MANY OTHER ORNAMENTALS — Spray generously at 1 teaspoonful per gallon water around base of plants, working into top 4 inches of soil. Repeat as necessary.

HOUSEHOLD

ANTS, BOX ELDER BUGS, (OUTSIDE BUILDINGS) — 5-7½ Tablespoonfuls per 1 gallon water. Apply along runways, in cracks in cement walks and patios, around garbage containers, under porches, around outside house foundations. Spot treat nests in lawns or other infested areas.

ANTS, ROACHES, SPIDERS, (FOR SPOT APPLICATION ONLY INSIDE BUILDINGS) — 3-7½ Tablespoonfuls per 1 gallon water. Spray thoroughly only trails around door sills, window frames, openings around water pipes, heat ducts, electrical outlets and baseboards, where these pests may enter. Treat cracks behind baseboards and areas behind shelves and other hiding places. DO NOT contaminate food, utensils, dishes or drinking water. Use a coarse wet spray. Repeat as necessary. May cause staining of finished surfaces.

TERMITES

FOR ALL TERMITE APPLICATIONS USE A 1.0% CONCENTRATION (1 PINT OF 72% CHLORDANE SPRAY DILUTED WITH 12 GALS. WATER). CONTROL IN EXISTING BUILDINGS, (BUILDINGS HAVING CRAWL

SPACES OR BASEMENTS) — Dig trenches adjacent to and around all piers and pipes and along the sides of foundation walls. Around solid concrete foundations without cracks the trenches should be 6 to 8 inches deep and wide. The emulsion should be poured in at the rate of 2 gallons per 10 linear feet of trench. Then as the excavated soil is put back into the trench, it also should be treated at the rate of 10 linear feet of trench. In brick or hollow-block foundations, or concrete foundations that have cracked, dig the trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. For example, a trench 3 feet deep would require 12 gallons per 10 linear feet. Treat voids in hollow-block foundations at the rate of 2 gallons per 10 linear feet of wall, at or near the footing.

(SLAB-ON-GROUND CONSTRUCTION) — Infestations in this type of construction are difficult to control. One method consists of drilling holes about a foot apart through the concrete slab, adjacent to all cracks and expansion joints, and injecting the chemical into the soil beneath the slab. Another method is to drill through the foundation walls from the outside and force the chemical just beneath the slab along the inside of the foundation and along all cracks and expansion joints. The emulsion should be applied at the rate of at least 4 gallons per 10 linear feet of foundation or expansion joint.

PRECONSTRUCTION TREATMENT, (SLAB-ON-GROUND CONSTRUCTION) — Apply an over-all treatment under entire surface of floor slab including porch floors and entrance platforms. Apply at rate of 1 gallon per 10 square feet, except that if fill under slab is gravel or other coarse absorbent material, apply at rate of 1½ gallons per 10 square feet.

Apply to critical areas to a depth of 1 foot, along both sides of foundation wall, along interior foundation walls, and around plumbing at the rate of 4 gallons per 10 linear feet. Chemical shall be mixed with the soil as it is being replaced in the trench.

Apply to voids of unit masonry foundation walls at or near bottom of foundation at rate of 2 gallons per 10 linear feet.

Under slab-on-ground porch floors and entrance platforms, apply over-all treatment at rate of 1 gallon per 10 square feet.

Where finished rooms are provided in basements or below grade areas, apply an over-all treatment to basement floor at rate of 1 gallon per 10 square feet.

GENERAL: Treatment shall not be made when the soil or fill is excessively wet or immediately after heavy rains, to avoid surface flow of the toxicant from application site. Unless the treated areas are to be immediately covered, precautions shall be taken to prevent disturbance of the treatment by human or animal contact with the treated soil. This chemical is toxic to animal and

plant life, and should be applied only with caution by an experienced person. Where soil treatment is used, individual water supply systems are not acceptable.

WARNING

- Harmful if swallowed. Avoid breathing spray mist. Contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. For eyes — flush with water for 15 minutes, get medical attention. Wash hands after use.
- Keep out of reach of children. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. DO NOT use on household pets or humans.
- DO NOT use or store near heat or open flames. DO NOT apply on plants in bloom as Chlordane is toxic to pollinizing insects. DO NOT apply to plants or lawns that are not well irrigated or when air temperature exceeds 90°F.
- DO NOT use in dairy barns or poultry houses. Avoid contamination of feed and foodstuffs. DO NOT feed forage treated with this material to dairy animals or to animals being finished for slaughter.
- This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. DO NOT apply when weather conditions favor drift from areas treated. DO NOT apply where runoff is likely to occur. DO NOT contaminate water by cleaning of equipment or disposal of wastes.
- Apply this product only as specified on this label. When container is empty, immediately wash thoroughly and destroy. Never re-use.
- DO NOT use on cats or humans.
- This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS AND WARNINGS STATED ON THIS LABEL.