

DRAGON

DRAGON

DRAGON

READ ENTIRE LABEL
WARNINGS AND
STATE REGULATIONS

DRAGON 10% S
of insect infestation,
oughly to upper and
days. DO NOT use
harvest except when

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near food or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

10%

SEVIN* DUST

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near food or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

BEANS: For Mexican Beans (10-15 sq. ft.). For Bean Leafhoppers, Velvet Beans and Tarnished Beetles (approximately 10-15 sq. ft.). Use up to the animals.

CORN: For Corn Earworm, Flea Beetle (10-15 lbs./acre (approximately 10-15 lbs./acre) as soon as they appear immediately after emergence and to entire extent of infestation agricultural experience be fed to livestock, product to the tassel and bee populations.

CABBAGE, BROCCOLI, CUMBERS, MELONS, TOMATOES, EGGS: Beetles and Harlequin Beetles (10-15 sq. ft.). For Cucumber Beetles, and Corn Earworms, and Corn Earworms (10-15 sq. ft.). Avoid excess made when tender. DO NOT apply to lettuce within 3 days.

For Chinch Bugs, Fire Ants, Earwigs, Flea Beetles and Bugs in Florida use 1

FOR CONTROL OF CERTAIN INSECTS ON VEGETABLES AND LAWNS

CAUTION.

EPA REG. NO. 16-98

DRAGON CHEMICAL CORPORATION

ROADSIDE VIRGINIA

DRAGON

DRAGON

DRAGON

READ ENTIRE LABEL
WARNINGS AND
STATE REGULATION

DRAGON 10% SE
of insect infestation,
oughly to upper and
days. DO NOT use
harvest except when

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.**

This product is highly toxic to bees exposed to direct treatment residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

For use instructions, use storage instructions of this product, consult the label with DIRECTIONS and PRECAUTIONS notes on this label.

10%

SEVIN* DUST

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.**

This product is highly toxic to bees exposed to direct treatment residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

For use instructions, use storage instructions of this product, consult the label with DIRECTIONS and PRECAUTIONS notes on this label.

BEANS: For Mexican Beans (sq. ft.). For Bean Leafhoppers, Velvet Beans (approximately 1 acre (approximately 1 sq. ft.). Use up to the animals.

CORN: For Corn Earworm, Flea Beetle (lb. acre (approximately 1/2 acre as they apply to the entire extent of infestation). For experiment, use 1/2 lb. per acre. For leafhoppers, use 1/2 lb. per acre.

CABBAGE, BROCCOLI, CUCUMBERS, MELONS, TOMATOES, EGGPLANT: For leafhoppers, use 1/2 lb. per acre. For Flea Beetles, use 1/2 lb. per acre. For leafhoppers, use 1/2 lb. per acre.

For Control of Flea Ants, Flea Beetles, Flea Bugs in Florida use 1/2

FOR CONTROL OF CERTAIN INSECTS ON VEGETABLES AND LAWNS.

ACTIVE INGREDIENTS	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate)	10%
INERT INGREDIENTS	90%
TOTAL	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK SIDE PANELS FOR ADDITIONAL CAUTIONS

*Trade Mark of UCC for 1-naphthyl N-methylcarbamate

EPA REG. NO. 16-98

MANUFACTURED BY
DRAGON CHEMICAL CORPORATION
ROANOKE, VIRGINIA

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

DRAGON 10% SEVIN[®] DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

DIRECTIONS

VEGETABLES

BEANS: For Mexican Bean Beetle use 5 lbs./acre (approximately 0.1 lb./1000 sq. ft.). For Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvet Bean Caterpillar and Western Bean Cutworm use 10 lbs./acre (approximately ¼ lb./1000 sq. ft.). For Armyworms, Corn Earworms, Stink Bugs and Tarnished Plant Bug use 10-15 lbs./acre (approximately ¼-½ lb./1000 sq. ft.). Use up to the day of harvest. DO NOT feed treated vines to dairy or meat animals.

CORN: For Corn Earworm, Corn Rootworm, Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetles and Leafhoppers use 10-20 lbs./acre (approximately ¼-½ lb./1000 sq. ft.). For Corn Earworm, apply to silks as soon as they appear. Make 3-4 applications at 2-3 day intervals. Re-apply immediately after rain. For Corn Borer, make one or more applications into the whorls and to entire plant beginning when borer damage threatens depending on the extent of infestation and degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. When forage is to be fed to livestock, DO NOT apply within 7 days of harvest. Application of this product to the tassel of corn during the pollen shed period will seriously reduce bee populations.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATOES, TOMATOES, EGGPLANT, PEPPER, AND OKRA: For Pickleworm, Melonworm, Flea Beetles and Harlequin Bug use 5-10 lbs./acre (approximately 1¼ lb./1000 sq. ft.). For Cucumber Beetles, Leafhoppers, Squash Bugs, Cabbage Caterpillars, Armyworms, and Corn Earworms use 10-15 lbs./acre (approximately ¼-½ lb./1000 sq. ft.). Avoid excessive applications. Some leaf injury may occur, if treatments are made when tender foliage is wet or during periods of prolonged high humidity. DO NOT apply to cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, or head lettuce within 3 days of harvest.

LAWNS

For Chinch Bugs, Fall Armyworms, Leafhoppers, Sod Webworms (Lawn Moths), Ants, Earwigs, Fleas, Millipedes, Mosquitoes, use 2 lbs. per 1000 sq. ft. Chinch Bugs in Florida use 12.5 lbs. per 5000 sq. ft.