



United States Environmental Protection Agency
Office of Pesticide Programs (H7505C)
Washington, DC 20460

Application for Pesticide:

Registration
 Amendment
 Other

OPP Identifier Number

188529

Section I

1. Company/Product Number 16-98	2. EPA Product Manager Edwards	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DRAGON ^(R) 10% SEVIN ^(R) Dust	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Dragon Corporation P. O. Box 7311 Roanoke, VA 24019 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below

Explanation: Use additional page(s) if necessary (For section I and Section II)

Compliance with PR Notice 93-3 -- Labeling Statement Prohibiting Application to Water

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3

Section III

DATE: 7/12/93

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
	If "Yes," Unit Package wgt	No. per container	If "Yes," Package wgt	<input checked="" type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
				Other (Specify) _____	
* Certification must be submitted.					
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container 4LB, 25LB		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed To Product		<input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other (_____)	

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name Rebecca M. Horton	Title Executive Assistant	Telephone No (Include Area Code) (703) 362-3617
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Rebecca M Horton</i>	3. Title Executive Assistant	
4. Typed Name Rebecca M. Horton	5. Date 6/21/93	

DRAGON®
10% SEVIN®
brand carbaryl insecticide
DUST

For control of certain insects on vegetables and lawns.

[NOTE TO EPA: Front panel of label may include graphics representing use, e.g., listed insects
Cucumber beetle, Stink bug, Corn borer, Tarnished plant bug.]

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)	10%
INERT INGREDIENTS:	<u>90%</u>
TOTAL	100%

Keep Out of Reach of Children
CAUTION

See back/side/other panel(s) for additional cautions.

NET WEIGHT 4 LB/25 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

GENERAL INFORMATION: DRAGON® 10% SEVIN® DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

VEGETABLES

BEANS: For Mexican bean beetle use 5 lbs./acre (approximately 0.1 lb./1000 sq. ft.). For Bean leaf beetle, Cucumber beetles, Flea beetles, Japanese beetles, Leafhoppers, Velvet bean caterpillar and Western bean cutworm, use 10 lbs./acre (approximately ¼ lb./1000 sq. ft.). For Armyworms, Corn earworms, Stink bugs and Tarnished plant bug, use 10-15 lbs./acre (approximately ¼ - ½ lb./1000 sq. ft.). Use up to the day of harvest. DO NOT feed treated vines to dairy or meat animals.

CORN: For Corn earworm, Corn rootworm adults, European corn borer, Fall armyworm, Flea beetles, Japanese beetle, Sap beetles and Leafhoppers, use 10-20 lbs./acre (approximately ¼ - ½ lb./1000 sq. ft.). For Corn earworm, apply to silks as soon as they appear. Make 3-4 applications at 2-3 day intervals. Re-apply immediately after rain. For Corn Borer, make one or more applications into the whorls and to entire plant beginning when borer damage threatens depending on the extent of infestation and degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. When forage is to be fed to livestock DO NOT apply within 7 days of harvest. Application of this product to the tassel of corn during the pollen shed period will seriously reduce bee populations.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATOES, TOMATOES, EGGPLANT, PEPPER AND OKRA: For Pickleworm, Melonworm, Flea beetles and Harlequin bug, use 5-10 lbs./acre (approximately 0.1-¼ lb./1000 sq. ft.). For Cucumber beetles, Leafhoppers, Squash bugs, Cabbage caterpillars, Armyworms and Corn earworms, use 10-15 lbs./acre (approximately ¼ - ½ lb./1000 sq. ft.). Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during periods of prolonged high humidity. DO NOT apply to cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi or head lettuce within 3 days of harvest.

LAWNS

For Chinch bugs, Fall armyworms, Leafhoppers, Sod webworms (lawn moths), Ants, Earwigs, Fleas, Millipedes, Mosquitoes, use 2 lbs. per 1000 sq. ft. Chinch bugs in Florida use 12½ lbs. per 5000 sq. ft.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

Label of Record - as submitted under 8/25/92, 1144
Notification

DRAGON[®]
10% SEVIN[®]
brand carbaryl insecticide
DUST

EPA Reg. No. 16-98

EPA Est. 16-VA-1

For control of certain insects on vegetables and lawns.

[NOTE TO EPA: Front panel of label may include graphics representing use, e.g., listed insects - Cucumber beetle, Stink bug, Corn borer, Tarnished plant bug.]

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 10%

INERT INGREDIENTS: 90%

TOTAL 100%

Keep Out of Reach of Children
CAUTION

See back/side/other panel(s) for additional cautions.

NET WEIGHT 4 LB/25 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, areas where surface water is present, or areas below the mean high tidal mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

GENERAL INFORMATION

DRAGON[®] 10% SEVIN[®] DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

VEGETABLES

BEANS: For Mexican bean beetle use 5 lbs./acre (approximately 0.1 lb./1000 sq. ft.). For Bean leaf beetle, Cucumber beetles, Flea beetles, Japanese beetles, Leafhoppers, Velvet bean caterpillar and Western bean cutworm, use 10 lbs./acre (approximately 1/4 lb./1000 sq. ft.). For Armyworms, Corn earworms, Stink bugs and Tarnished plant bug, use 10-15 lbs./acre (approximately 1/4 - 1/3 lb./1000 sq. ft.). Use up to the day of harvest. DO NOT feed treated vines to dairy or meat animals.

CORN: For Corn earworm, Corn rootworm adults, European corn borer, Fall armyworm, Flea beetles, Japanese beetle, Sap beetles and Leafhoppers, use 10-20 lbs./acre (approximately 1/4 - 1/2 lb./1000 sq. ft.). For Corn earworm, apply to silks as soon as they appear. Make 3-4 applications at 2-3 day intervals. Re-apply immediately after rain. For Corn Borer, make one or more applications into the whorls and to entire plant beginning when borer damage threatens depending on the extent of infestation and degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. When forage is to be fed to livestock DO NOT apply within 7 days of harvest. Application of this product to the tassel of corn during tassel pollen shed period will seriously reduce bee populations.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATOES, TOMATOES, EGGPLANT, PEPPER AND OKRA: For Pickleworm, Melonworm, Flea beetles and Harlequin bug, use 5-10 lbs./acre (approximately 0.1-1/4 lb./1000 sq. ft.). For Cucumber beetles, Leafhoppers, Squash bugs, Cabbage caterpillars, Armyworms and Corn earworms, use 10-15 lbs./acre (approximately 1/4 - 1/3 lb./1000 sq. ft.). Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during periods of prolonged high humidity. DO NOT apply to cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi or head lettuce within 3 days of harvest.

LAWNS

For Chinch bugs, Fall armyworms, Leafhoppers, Sod webworms (lawn moths), Ants, Earwigs, Fleas, Millipedes, Mosquitoes, use 2 lbs. per 1000 sq. ft. Chinch bugs in Florida use 12 1/2 lbs. per 5000 sq. ft.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

12/2/6

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS** and **CAUTIONS** stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019