


UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

FEB 10 1994

Dragon Corporation P.O. Box 7311 Roanoke, VA 24019



OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Subject: Label Amendment Submission of 7/29/93 in Response to PR Notice 93-7 EPA Reg. No. 16-98 Dragon 10% Sevin Dust

Dear Registrant:

ř

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed. The comments follow.

# WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

2/2/94

BEST AVAILABLE COP

Send to EPA one (1) copy of the final printed labeling:

BEFORE selling or distributing any product bearing the final printed labeling
AND

WITHIN one year from date of this acceptance.



Page 2

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL) Office of Pesticide Programs (7505C) U.S. Environmental Protection Agency 401 M Street, SW Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL) Office of Pesticide Programs Room 266A, Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202

Sincerely,

Jomes a. Downing

Registration Support Branch Registration Division (7505W)

Attachment

### DRAGON® 10% SEVIN® brand carbaryl insecticide DIST

vith COMMENTS In EPA Letter Dated FEB 10 1994 Under the Federal Insecticide,

Fundicide, and Rodenticide Act

as amended, for the posticide registered under EPA Rog. No. 16-98

ACCEPTED

2 ' L+

DUST

For control of certain insects on vegetables and lawns.

[NOTE TO EPA: Front panel of label may include graphics representing use, e.g., listed insects -Cucumber beetle, Stink bug, Corn borer, Tarnished plant bug.]

## ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)	10%
INERT INGREDIENTS:	<u>90%</u>
ΤΟΤΛΙ	00%

### Keep Out of Reach of Children CAUTION

See back/side/other panel(s) for additional cautions.

### NET WEIGHT 4 LB/25 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

HARMFUL IF INHALED OR SWALLOWED. DO NOT oreathe dust. DO NOT take internally. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

GENERAL INFORMATION: FOR HOMEOWNER USE ONLY. DRAGON<sup>®</sup> 10% SEVIN<sup>®</sup> DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat application every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

EPA Reg. No. 16-98 July 21, 1993



EPA Est, 16-VA-1 Page 1

#### DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

### VEGETABLES

- BEANS: For Mexica's bean heetle use '4 ib. per 2000 sq. ft. For Bean leaf beetle, Cucumber beetles, Flea beetles, Japanese beetles, Leafhoppers, Velvet bean caterpillar and Western bean cutworm, use '4 lb. per 1000 sq. ft. For Armyworms, Corn carworms, Stink bugs and Tarnished plant bug, use '4 - '/, ib. per 1000 sq. ft. Use up to the day of harvest. DO NOT feed treated vines to domestic animals.
- CORN: For Corn earworm, Corn rootworm adults, European corn borer, Fall armyworm, Flea beetles, Japanese beetle, Sap beetles and Leafhoppers, use  $\frac{1}{2} - \frac{1}{2}$  lb. per 1000 sq. ft. For Corn earworm, apply to silks as soon as they appear. Make 3-4 applications at 2-3 day intervals. Reapply immediately after rain. For Corn Borer and larvae in whorls (budworm damage) and for foliage feeders, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. Allow 7 days before harvest, or before allowing domestic animals to feed in treated areas. Application of this product to the tassel of corn during the pollen shed period will seriously reduce bee populations.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI, CUCUMBERS, MELONS, PUMPKIN, SQUASH, HEAD LETTUCE, CARROTS, POTATOES, TOMATOES, EGGPLANT, PEPPER AND OKRA: For Pickleworm, Melonworm, Flea beetles and

Harlequin bug, use  $\frac{1}{-1}$  lb. per 2000 sq. ft. For Cucumber beetles, Leafhoppers, Squash bugs, Cabbage caterpillars, Armyworms and Corn earworms, use  $\frac{1}{-1}$  lb. per 1000 sq. ft. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or during periods of prolonged high humidity. DO NOT apply to cabbage, broccoli, brussels sprouts, caulitlower, kohlrabi or head lettuce within 3 days of harvest.

#### LAWNS

For Chinch bugs, Fall armyworms, Leafhoppers, Sod webworms (lawn moths), Ants, Earwigs, Fleas, Millipedes, Mosquitoes, use 2 lbs. per 1000 sq. ft. Chinch bugs in Florida use  $12\frac{1}{2}$  lbs. per 5000 sq. ft.

#### STORAGE & DISPOSAL

- **STORAGE:** Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.
- **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.
- CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN<sup>®</sup> is a registered trademark of Rhóne Poulenc Ag Company.

Manufactured by DRAGON CORPORATION Roanoke, VA 24019 BEST AVAILABLE COPT

EPA Est. 16-VA-1

Page 2

EPA Reg. No. 16-98 July 21, 1993