


16-99

07/23/1973

1/4

FRONT PANEL



DRAGON

SEVIN*
50 WETTABLE

NET CONTENTS
2 pounds

A WETTABLE POWDER FOR CONTROL OF CERTAIN LISTED INSECT PESTS ON
FRUIT, VEGETABLES, ORNAMENTALS, LAWNS, TOBACCO AND ALFALFA.


ACTIVE INGREDIENT		By Wt.
Carbaryl (1-naphthyl N-methylcarbamate).....		50%
INERT INGREDIENT		50%
	Total	<u>100%</u>

*Trade Mark for UCC for 1-naphthyl N-methylcarbamate.

EPA REG. NO. 16-99

CAUTION: KEEP OUT OF REACH OF CHILDREN. See side/back panels
for additional cautions.

MANUFACTURED FOR
 DRAGON CHEMICAL CORPORATION
 Roanoke, Virginia 24019



GENERAL INFORMATION

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

Keep spray mixtures well agitated. DRAGON SEVIN 50 WETTABLE may be applied within 1 day of harvest on all food crops listed unless otherwise noted. Apply when insects or damage appear, repeat at 7 to 10 day intervals or as necessary.

DIRECTIONS

APPLES--PEARS--(Eastern States)--For control of green apple aphid, white apple leafhopper, apple mealybug, use 1 lb. per 100 gals. water (2 Tbs./gal.). To control apple maggot, codling moth, oriental fruit moth, and plum curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved. For fruit tree leaf roller, periodical cicada (locusts), eastern tent caterpillar, Japanese beetle, San Jose scale, bagworm, use 2 lbs. per 100 gals. water (4 Tbs./gal.). Avoid thinning by delaying use until 30 days after petal fall. For maximum scale control apply when crawlers are present.

GRAPES--Use 1-2 lbs. per 100 gals. water (2-4 Tbs./gal.) for European Fruit Lecanium, grape leaf folder and grape leafhopper; apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. Use 2 lbs. per 100 gals. water (4 Tbs./gal.) for grape berry moth, Japanese beetle and June beetle. Dilute spray of 200 gal./acre (4½ gal./1000 sq. ft.) is suggested.

PEACHES--For oriental fruit moth, peach twig borer, lesser peach tree borer, plum curculio, cat facing bug, periodical cicada (locusts), Japanese beetle, San Jose scale, use 2 lbs. per 100 gals. water (4 Tbs./gal.). For best scale control apply when crawlers are present.

STRAWBERRIES--Meadow Spittlebug, Strawberry Leaf Roller, use 2 lbs. per 100 gals. water (4 Tbs./gal.) (200 gals. per acre), (4½ gal./1000 sq. ft.). Make full coverage application. Apply in summer sprays. Repeat 3 to 5 times when insects appear.

BEANS--For control of Mexican bean beetle, use 1 lb. per 100 gals. water per acre (2 Tbs./gal./1000 sq. ft.). for leafhoppers, bean leaf beetle, flea beetle, Japanese beetle, use 2 lbs. per 100 gals. water per acre (4 Tbs./2½ gals./1000 sq. ft.). For Lygus, stink bugs, lima bean pod borer, use 4 lbs. per 100 gal. water per acre (8 Tbs./2½ gal./1000 sq. ft.). DO NOT feed treated bean vines to dairy or meat animals.

CORN--For control of corn earworm, European corn borer, fall armyworm, Japanese beetle, flea beetle, leafhoppers, use 2-4 lbs. per 100 gals. water per acre (4-8 Tbs./2½ gal./1000 sq. ft.). Make applications into the whorls and to the entire plant. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silk begins to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. When forage is to be fed to livestock, DO NOT apply within 7 days of harvest. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

3/4

CUCUMBERS, MELONS, PUMPKINS, SQUASH--For pickleworm, melonworm, apply 1 lb. per 100 gals. water per acre (2 Tbs./2½ gals./1000 sq. ft.). For cucumber beetles, squash bugs, flea beetles, leafhoppers, use 2 lbs. per 100 gals. per acre (4 Tbs./2½ gal./1000 sq. ft.).

LETTUCE, COLLARDS, BALE, TURNIP GREENS, MUSTARD GREENS, SPINACH--For control of flea beetles, leafhoppers, harlequin bug, apply 1 to 2 lbs. per 100 gals. water per acre (2-4 Tbs./2½ gal./1000 sq. ft.). For six-spotted leafhopper, corn earworm, armyworm, imported cabbageworm, spittlebugs, lygus and stink bugs, apply 2 to 4 lbs. per 100 gals. water per acre (4-8 Tbs./2½ gals./1000 sq. ft.). DO NOT apply within 14 days of harvest. DO NOT apply to leaf lettuce within 14 days of harvest or within 3 days to head lettuce.

TOMATO, EGGPLANT, PEPPERS--Use 1-2 lbs. per 100 gals. water per acre (2-4 Tbs./2½ gals./1000 sq. ft.) for control of Colorado Potato Beetle, flea beetles, leafhoppers. Use 2-4 lbs. per 100 gals. water per acre (4-8 Tbs./2½ gals./1000 sq. ft.) for tomato fruitworm, tomato hornworm, European Corn Borer, tarnished plant bugs, lace bugs.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS--For control of flea beetles, harlequin bug, apply 1 to 2 lbs. per 100 gals. water per acre (2-4 Tbs./2½ gals./1000 sq. ft.). For corn earworm, stink bug, armyworm, imported cabbageworm, apply 2 to 4 lbs. per 100 gals. water per acre (4-8 Tbs./2½ gals./1000 sq. ft.). DO NOT apply within 3 days of harvest.

TOBACCO--(in fields)--For budworm, flea beetles, Japanese beetle, and Hornworm, use 2-4 lbs. per 100 gals. water per acre. Avoid excessive application. If later season applications are necessary, treat immediately after priming rather than before priming.

ALFALFA--For Spring application control of Alfalfa weevil larvae, use 3 lbs. in 15-20 gals. water per acre. For leafhoppers, thrips, three-cornered alfalfa hopper, alfalfa caterpillar, green cloverworm, blister beetle, use 2 lbs. in 15-20 gals. water per acre. For armyworm, webworms, bollworm or corn earworm, stink bugs, use 2-3 lbs. in 15-20 gals. water per acre. For grasshoppers use 1-3 lbs. per 100 gals. water per acre. May be applied up to and including the day of harvest. DO NOT apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

ROSES, AZALEAS, OTHER ORNAMENTALS, SHRUBS, SHADE TREES--For control of certain aphids, bagworm, blister beetle, boxelder bug, boxwood leaf miner, elm leaf beetle, flea beetles, Japanese beetle, lace bugs, leafhoppers, leaf rollers, oak leaf miner, periodical cicada (locusts), rose aphid, rose slug, scale insects, tent caterpillar, gypsy moth, use 2 lbs. per 100 gals. water (4 Tbs. per gal.).

LAWNS--For control of chinch bugs, leafhopper, fall armyworm, sod webworm, lawn moths, ants, earwigs, fleas, millipeds, mosquitoes, use 2 lbs. per 150-200 gals. water per 5000 square feet of lawn.

4/4

SIDE PANEL

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or feedstuffs.**

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.