

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Avoid contact with skin. Skin contact may be harmful. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs. This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. DO NOT re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.



✓ 50 WETT

GENERAL INFORMATION

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS. Keep spray mixtures well agitated. DRAGON SEVIN 50 WETTABLE material on all food crops listed unless otherwise noted. Apply when insects are active. Repeat applications at 10 day intervals or as necessary.

DIRECTIONS

APPLES-PEARS—(Eastern States)—For control of codling moth, green hopper, apple mealybug, use 1 lb. per 100 gals. water (2 Tbs./gal.) fruit tree leaf roller, pear psylla, periodical cicada (locusts), pear beetle, San Jose scale, bagworm, use 2 lbs. per 100 gals. water delaying use until 30 days after petal fall. For maximum scale control present. Apply at 10 day intervals for optimum apple maggot control. For codling moth and plum curculio, apply every 10 to 14 days until