

16-106

# DIPEL® 150 DUST\*

(Bacillus thuringiensis)

4 LBS.  
NET WT.

# DIPEL® 150 DUST\*

(Bacillus thuringiensis)

**ACTIVE INGREDIENT:**

<u>Bacillus thuringiensis</u> , Berliner Potency of 320 International	
Units per mg. (at least 0.5 billion viable spores	
per Gm.)*	0.064%
INERT INGREDIENTS	99.936%
Total	100.000%

\*Equivalent to potency of 0.15 billion International Units per pound of this product.

**NOTICE**

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

REASONABLE CARE SHOULD BE EXERCISED TO AVOID INHALATION OR CONTACT WITH OPEN WOUNDS. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.



# DIPEL® 150 DUST

(Bacillus thuringiensis)

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

**DIRECTIONS FOR USE**

DIPEL 150 Dust is a biological insecticide intended for use against chewing larvae (caterpillars) of certain lepidopterous species.

DIPEL 150 Dust is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

**APPLICATION RATE**

For the control of Cabbage Looper on Broccoli, Cabbage, Cauliflower, Collards, Kale, Mustard and Turnip Greens, Celery, Lettuce, Melons, Potatoes, Tomatoes and Cucumbers (Southeastern U. S. only) apply 25 to 50 lbs. per acre (approximately 1/2 to 1 lb./1000 sq. ft.)

For the control of Imported Cabbage Worm on Broccoli, Cabbage, Cauliflower, Collards, Kale and Mustard and Turnip Greens, apply 15 to 25 lbs. per acre (approximately 1/2 to 1 lb./1000 sq. ft.)

For the control of Grape Leaf Folder on Grapes, apply 25 to 50 lbs. per acre (approximately 1/2 to 1 lb./1000 sq. ft.)

For the control of Hornworm and Budworm on Tobacco and Tomatoes, apply 12 to 25 lbs. per acre (approximately 1/4 to 1/2 lb./1000 sq. ft.)