

16-119

12/12/2002

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Thomas McEntee
Dragon Chemical Corporation
71 Carolyn Blvd.
Farmingdale, NY 11735

DEC 12 2002

Dear Mr. McEntee:

Subject: Amendment - Comply with 2000 Diazinon MOA;
Delete Celery;
Update First Aid per PR Notice 2001-1; and
Update CSF/replacement diazinon source
Dragon 5% Diazinon Granules
EPA Registration Number 16-119
Your submissions dated October 25 and December 10, 2002

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. On the front panel, slightly revise the site limitation so that it reads "FOR OUTDOOR USE ON THE HOME LAWN AND THE HOME GARDEN ONLY".
 - b. On the front panel claims, revise "and many others" to read "and many others as listed in Directions For Use".
 - c. Under the heading Directions For Use, after the sentence beginning "It is a violation of Federal law..." slightly revise the prohibition so that it reads "Reformulation or repackaging of this product is prohibited."
 - d. In the first sentence of the directions for Fire Ants, add "home" before "lawns".
 - e. Under the heading VEGETABLE GARDENS, after the sentence beginning "Applications for control of pests listed below..." add "Do not apply this product in greenhouses."

- f. In the entry Wireworms, Flea Beetles - Sweet Potatoes, delete the pest Flea Beetles since soil application will not control this foliar pest. Since the only remaining pest is wireworms, you may choose to just include Sweet Potatoes in the list of plants under the heading Wireworms rather than have a separate entry for this crop with this same pest.
 - g. In the Hazards to Humans and Domestic Animals, after "Do not store near food or feed products." add "Do not use inside the home or in greenhouses."
 - h. In the FIRST AID section add a fourth item to the If swallowed advice to read " Do not induce vomiting or give anything by mouth if person is unconscious or convulsing."
3. Submit three (3) copies of your final printed labeling before you release the product for shipment.
 4. Comply with all applicable provisions of the 2000 Memorandum of Agreement Regarding Pesticide Products Containing Diazinon (i.e., the Diazinon MOA). In accordance with the Diazinon MOA you have requested the following revisions to the conditions of registration for this diazinon product with outdoor non-agricultural (i.e., home and garden) uses:
 - a. You have requested under section 6(f) of FIFRA that this registration be voluntarily canceled on a date not later than December 31, 2004.
 - b. You may not manufacture, label, or release for shipment this product after June 30, 2003.
 - c. You may not sell or distribute this product after August 31, 2003.
 - d. The registration for this product shall expire and be canceled on December 31, 2004.
 - e. You will send to every person who directly purchases this product (i.e., to every seller) a notification of Product Recovery Program to be conducted by Syngenta and Makhteshim after December 31, 2004.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The Basic Confidential Statement of Formula (CSF) dated November 5, 2002 which identifies a replacement diazinon source Manufacturing Use Product is acceptable and has been added to this product's registration file.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

DRAFT 4/13

ACCEPTED
with COMMENTS
In EPA Letter Dated:

DEC 12 2002


Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for insecticide
registered under EPA Reg. No.
16-119

DRAGON[®]

5% DIAZINON GRANULES

FOR OUTDOOR USE ON THE HOME LAWN
AND IN THE HOME GARDEN ONLY

This product must not be used on golf courses and sod farms.



Kills: Ants, Armyworms, Bermudagrass mites, Billbugs, Brown dog ticks, Chiggers, Chinch bugs, Crickets, Cutworms, Earwigs, Fleas, Root maggots, Sod webworms, Wireworms, and White grubs of Japanese beetle, European chafer, and Southern chafer, and many others.

TREATS UP TO 5,000 SQUARE FEET

ACTIVE INGREDIENT:	BYWT:
Diazinon [0,0-Diethyl 0-(2isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	5%
INERT INGREDIENTS:	95%
TOTAL:	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for additional
precautionary statements.

NET WEIGHT 10 LB

5/13

RE-CUMULATING OR RE-APPLYING THIS PRODUCT IS PROHIBITED

DRAGON[®] 5% DIAZINON GRANULES

IN LAWNS, DRAGON[®] 5% DIAZINON GRANULES KILLS:

Ants	Crickets	Millipedes
Armyworm	Cutworms	Sod webworms
Bermudagrass mites	Digger Wasps	Sowbugs (pillbugs)
Billbugs	Earwigs	Springtails
Brown dog ticks	Fire Ants*	White Grubs of:
Chiggers	Fleas	European chafar
Chinch bugs	Lawn Moths	Japanese beetle
Clover mites	Leafhoppers	Southern chafar
		*Aids in control

IN GARDENS, DRAGON[®] 5% DIAZINON GRANULES KILLS:

Cutworms	Mole crickets	Root maggots
Flea beetles	Onion maggots	Wireworms

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

SPECIAL PRECAUTIONS TO REDUCE RISK TO BIRDS: Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds: (1) Do not treat areas larger than 15,000 sq. ft. (2) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. (3) Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. (4) Avoid overwatering treated areas. (5) Follow all watering instructions in the directions for use, however, stop watering before puddling occurs. (6) Do not apply this product to areas occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds. (7) If it is necessary to repeat applications, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

DRAGON[®] 5% DIAZINON GRANULES KILLS the following lawn insects on both grass and dichondra lawns. Apply this product with a fertilizer spreader. Refer to the following table for spreader settings. Apply at the rate of 2 lbs. per 1,000 sq. ft. Note: 1,000 sq. ft. = an area 20' x 50'; 100 sq. ft. = an area 10' x 10'; 2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (pillbugs), Springtails (collembola). - Apply when grass is dry. Water lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. Billbugs, White grub of Japanese beetle, European chafar, Southern chafar, Aeneas spruceus (dung beetle): Water grass with at least 1/4 inch of water after application. However, stop watering before puddling occurs. For **billbug control**, apply when activity is first observed or when chewed indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For **grub control**, except dung beetle, treat anytime between late July and early October. For **dung beetle control**, apply in Spring or Summer when damage is first noticed. Water grass with at least 1/4 inch of water after application; however, stop watering before puddling occurs. Treat again in late Summer or early Fall if a second generation occurs. Hyperodone weevils: Water grass with at least 1/4 inch of water after application, however stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

Spreader Make and Model	Spreader Setting No.
Central Imperial Mark #1983-G	6 1/2
Central Saburo Lawn Spreader #1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 106	6 1/2
ORTHO Drop Spreader #3000	4 1/2
ORTHO Broadcast Spreader #4000	4 1/2
J. C. Parry Model 6005	7 1/4
J. C. Parry Rotary Spreader	5
Scott's Precision Flow Spreader PF-4	4
Scam Craftsman #871-10198	6

*For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

LAWN SPOT TREATMENT

ANTS - Apply 1 1/2 teaspoons of DRAGON[®] 5% DIAZINON GRANULES over and around each ant hill. **FIRE ANTS** - To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply 1/2 cup over, and 2 ft. around each fire ant mound immediately after application, water-in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gal. of water should be applied to each mound area. Reapply as necessary. Note: For best results, apply in cool weather, 55° - 80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger wasps - Apply 1 1/2 teaspoons of DRAGON[®] 5% DIAZINON GRANULES in and around each nest opening. Apply at dark when wasps are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION: The following pests are commonly found around or near house foundations, and may enter the house. Ants, Brown dog ticks, Clover mites, Crickets, Earwig, Flea, Millipede, Sowbug, (pillbug), Springtails (collembola). Apply a 2 foot band of DRAGON[®] 5% DIAZINON GRANULES next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (2' by 50') treated. If necessary, repeat application at 3-week intervals.

VEGETABLE GARDENS

SPECIAL PRECAUTIONS TO REDUCE RISK TO BIRDS: Birds feeding on treated areas may be killed. To reduce risk to birds: (1) Thoroughly mix DRAGON[®] 5% DIAZINON GRANULES into the soil during or immediately after application. Refer to the following directions for depth of soil incorporation. (2) Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

DRAGON[®] 5% DIAZINON GRANULES kills: Cutworms, Mole Crickets, Root Maggots, Wireworms, Flea Beetles, and Seed Corn Maggots on crops listed below. Apply uniformly as directed for each crop. When a range of rates is given, use the higher only for heavier infestations.

Applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

Cutworms: - (Surface and Subterranean), Beans^{*}, (pole, snap or lima) (succulent only), Beets, (red table), Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Collards, Cucumber, Endive (escarole), Kale, Lettuce, Melons, (cantaloupe, casaba, crenshaw, honeydew, muskmelon, Persians & hybrids of these), Watermelons, Parsley, Peas^{*}, (succulent only), Peppers, Potatoes, (surface and subterranean), Spinach, Squash, (summer & winter), Sweet Corn, Swiss Chard, Tomatoes. Using 7 - 14 oz. per 500 sq. ft., broadcast and mix into soil just before planting. For surface cutworms, mix into soil 2 - 3 inches deep. For subterranean cutworms, mix into soil 3 - 6 inches deep. If type of cutworm cannot be identified, mix into soil 4 - 6 inches deep. *Do not use on dried bean varieties such as pintos, dried limas, navy beans, etc. *Do not use on dried pea varieties such as dried blackeyed peas.

Wireworms - Beans^{*} (pole, snap or lima), (succulent only), Beets, (red, table), Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, Celery, Collards, Cucumber, Endive (escarole), Kale, Lettuce, Melons (cantaloupe, casaba, crenshaw, honeydew, muskmelon, Persians & hybrids of these), Watermelons, Onions (green & bulb), Parsley, Peas^{*} (succulent only), Peppers, Potatoes, Spinach, Summer Squash, Winter squash, Sweet corn, Swiss Chard, tomatoes, Turnips. Using 10 1/2 to 14 oz. per 500 sq. ft., broadcast and mix thoroughly into soil 4 - 8 inches deep just before planting. *Do not use on dried bean varieties such as pintos, dried limas, navy beans, etc. *Do not use on dried pea varieties such as dried blackeyed peas.

Wireworms, Flea beetles - Sweet Potatoes. Using 10 1/2 oz. per 500 sq. ft., broadcast and mix into soil 4 - 8 inches just before planting.

Mole Crickets - Beets, (Red, table), Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, Turnips. Using 3 1/2 oz. per 500 sq. ft. broadcast and mix into soil 1 - 2 inches deep, one to two days before planting.

Root Maggots - Broccoli, Brussels Sprouts, Cabbage, Cauliflower. Using 7 - 10 1/2 oz. per 500 sq. ft. broadcast and mix into soil 3 - 4 inches deep just before planting or transplanting.

Seed Corn Maggots - Sweet Corn. Using 7 - 14 oz. per 500 sq. ft. broadcast and mix into soil 2 inches deep just before planting.

Onion Maggots - Onions (bulb & green). Using 7 - 14 oz. per 500 sq. ft., broadcast just before planting and mix into the top 3 - 4 inches of soil. Note: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. **IF INHALED:** Remove victim to fresh air and apply artificial respiration, if indicated. **IF ON SKIN:** Wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention. **IF IN EYES:** Rinse eyes with plenty of water and call a physician immediately. **NOTE TO PHYSICIAN:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS: This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overapplying granules. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

For lawns, follow all watering instruction in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to areas occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes.

Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

STORAGE AND DISPOSAL


STORAGE: Store in original container in a cool, dry, well ventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food or feed. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty container. Discard bag in container. For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

EPA Reg. No 16-119

EPA Est. 16-VA-1 (1); 44616-MO-01 (2); 10370-TX-1 (3); 829-NC-2 (4); 32802-MO-1B (5); 18713-GA-1 (7) 9188-OH-2 (8); 44615-GA-1 (9); 769-GA-1 (10); 788-GA-1 (11); 3342-NC-1 (12). Number following Est. No. corresponds to first digit in Lot No.

D-1



DISTRIBUTED BY
DRAGON[®] CHEMICAL CORPORATION
P. O. Box 7311
ROANOKE, VA. 24019

REFER TO APPENDIX A

12/17/2002

READING COPY OF 6/13

[FRONT PANEL]

**DRAGON®
5% DIAZINON GRANULES**

Nothing Beats The Dragon!

This product must not be used on golf courses and sod farms.

Kills: Ants, Armyworms, Bermudagrass mites, Billbugs, Brown dog ticks, Chiggers, Chinch bugs, Crickets, Cutworms, Earwigs, Fleas, Root maggots, Sod webworms, Wireworms, and White grubs of Japanese beetle, European chafer and Southern chafer, and many others.

READY TO USE

TREATS UP TO 5,000 SQUARE FEET

FOR OUTDOOR USE ON THE HOME LAWN AND IN THE HOME GARDEN ONLY.

[Note to EPA: Label front panel may include graphics illustrating product use; e.g., Ticks, Fleas, White grubs, etc.]

ACTIVE INGREDIENT:	BY WT.
Diazinon [O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	5%
INERT INGREDIENTS:	<u>95%</u>
TOTAL:	100%

Keep Out of Reach of Children
CAUTION

See additional precautionary statements on back/side/other panel(s).

NET WT. 10 LB

[BACK/SIDE/OTHER PANEL(S)]

IN LAWNS, DRAGON® 5% DIAZINON GRANULES KILLS:

- | | | |
|--------------------|--------------|--------------------|
| Ants | Crickets | Millipedes |
| Armyworms | Cutworms | Sod webworms |
| Bermudagrass mites | Digger wasps | Sowbugs (pillbugs) |
| Billbugs | Earwigs | Springtails |
| Brown dog ticks | Fire ants* | White grubs of: |
| Chiggers | Fleas | European chafer |
| Chinch bugs | Lawn moths | Japanese beetle |
| Clover mites | Leafhoppers | Southern chafer |

*Aids in control

IN GARDENS, DRAGON® 5% DIAZINON GRANULES KILLS:

Cutworms
Mole crickets

Onion maggots
Root maggots

Wireworms
Flea beetles

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

REFORMULATING OR REPACKAGING THIS PRODUCT IS PROHIBITED

OVERALL LAWN APPLICATION

SPECIAL PRECAUTIONS TO REDUCE RISK TO BIRDS: Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds: (1) Do not treat areas larger than 15,000 sq. ft. (2) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. (3) Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. (4) Avoid overlapping treated areas. (5) Follow all watering instructions in the directions for use, however, stop watering before puddling occurs. (6) Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds. (7) If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

DRAGON® 5% DIAZINON GRANULES kills the following lawn insects on both grass and dichondra lawns. Apply this product with a fertilizer spreader. Refer to the following table for spreader settings. Apply at the rate of 2 lb. per 1,000 sq. ft. Note: 1,000 sq. ft. = an area 20'x50'; 100 sq. ft. = an area 10'x10'; 2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (pillbugs), Springtails (collembola) — Apply when grass is dry. Water lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.

Billbugs, White grubs of Japanese beetle, European chafer, Southern chafer, *Ataenius spretulus* (dung beetle) — Water grass with at least ¼ inch of water after application, however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle control, apply in Spring or Summer when damage is first noticed. Water grass with at least ¼ inch of water after application, however, stop watering before puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.

Hyperodes weevils — Water grass with at least ¼ inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

LAWN SPREADER SETTINGS (to deliver 2 lbs. per 1,000 sq. ft.)**
 Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application.

Spreader Make and Model	Spreader Setting No.
Central Imperial Mark 1662-0	6 1/2
Central Suburbia Lawn Spreader #1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader #3000	4 1/2
ORTHO Broadcast Spreader #4000	4 1/2
J.C. Penney Model 6005	7 1/4
J.C. Penney Rotary Spreader	5
Scott's Precision Flow Spreader PF-4	4
Sears Craftsman #671.19198	6

**For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

LAWN SPOT TREATMENT

Ants — Apply 1 1/2 teaspoons of DRAGON® 5% Diazinon Granules over and around each ant hill.

Fire Ants — To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water-in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gal. of water should be applied to each mound area. Reapply as necessary.

Note: For best results, apply in cool weather, 65°-80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger wasps — Apply 1 1/2 teaspoons of DRAGON® 5% Diazinon Granules in and around each nest opening. Apply at dusk when wasps are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION

The following pests are commonly found around or near house foundations, and may enter the house. Ants, Brown dog ticks, Clover mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs (pillbugs), Springtails (collembola). Apply a 2 foot band of DRAGON® 5% Diazinon Granules next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (2' x 50') treated. If necessary, repeat application at 3-week intervals.

VEGETABLE GARDENS

SPECIAL PRECAUTIONS TO REDUCE RISK TO BIRDS: Birds feeding on treated areas may be killed. To reduce risk to birds: (1) Thoroughly mix DRAGON® 5% Diazinon Granules into the soil during or immediately after application. Refer to the following directions for depth of soil incorporation. (2) Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

DRAGON® 5% Diazinon Granules kills: Cutworms, Mole crickets, Root maggots, Wireworms, Flea beetles and Seed Corn maggots on crops listed below. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

Cutworms — (Surface and Subterranean). Beans¹ (pole, snap or lima) (succulent only), Beets (red, table), Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, ~~Celery~~, Collards, Cucumber, Endive (escarole), Kale, Lettuce, Melons (cantaloupe, casabas, crenshaws, honeydew, muskmelon, Persians and hybrids of these), Watermelons, Parsley, Peas² (succulent only), Peppers, Potatoes (surface and subterranean), Spinach, Squash (summer & winter), Sweet corn, Swiss chard, Tomatoes. Using 7-14 oz. per 500 sq. ft., broadcast and mix into soil just before planting. For surface cutworms, mix into soil 2-3 inches deep. For subterranean cutworms, mix into soil 3-6 inches deep. If type of cutworm cannot be identified, mix into soil 4-6 inches deep.



¹Do not use on dried bean varieties such as pintos, dried limas, navy beans, etc. ²Do not use on dried pea varieties such as dried blackeyed peas.

Wireworms — Beans¹ (pole, snap or lima) (succulent only), Beets (red, table), Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, ~~Celery~~, Collards, Cucumber, Endive (escarole), Kale, Lettuce, Melons (cantaloupes, casabas, crenshaws, honeydew, muskmelons, Persians & hybrids of these), Watermelons, Onions (green & bulb), Parsley, Peas² (succulent only), Peppers, Potatoes, Spinach, Summer squash, Winter squash, Sweet corn, Swiss chard, tomatoes, Turnips. Using 10½-14 oz. per 500 sq. ft., broadcast and mix thoroughly into soil 4-8 inches deep just before planting.



¹Do not use on dried bean varieties such as pintos, dried limas, navy beans, etc. ²Do not use on dried pea varieties such as dried blackeyed peas.


Wireworms, Flea beetles — Sweet potatoes. Using 10½ oz. per 500 sq. ft., broadcast and mix into soil 4-8 inches just before planting

Mole Crickets — Beets (red, table), Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, Turnips. Using 3½ oz. per 500 sq. ft., broadcast and mix into soil 1-2 inches deep, one to two days before planting.

Root Maggots — Broccoli, Brussels sprouts, Cabbage, Cauliflower. Using 7-10½ oz. per 500 sq. ft., broadcast and mix into soil 3-4 inches deep just before planting or transplanting.

Seed Corn Maggots — Sweet corn. Using 7-14 oz. per 500 sq. ft., broadcast and mix into soil 2 inches deep just before planting.

Onion Maggots — Onions (bulb & green). Using 7-14 oz. per 500 sq. ft., broadcast just before planting and mix into the top 3-4 inches of soil. Note: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

REFER TO APPENDIX A

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

IF INHALED: Remove victim to fresh air and apply artificial respiration, if indicated.

IF ON SKIN: Wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention.

IF IN EYES: Rinse eyes with plenty of water and call a physician immediately.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

For lawns, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes.

Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

STORAGE: Store in original container in a cool, dry, well ventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food or feed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

11/13

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS** and **CAUTIONS** stated on this label.

Roanoke, VA 24019

Appendix A

12/13

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed	<ul style="list-style-type: none">•Call a poison control center, doctor, or 1-800-xxx-xxxx¹ immediately for treatment advice.•Have person sip a glass of water if able to swallow.•Do not induce vomiting unless told to do so by the poison control center or doctor.²
If on skin or clothing	<ul style="list-style-type: none">•Take off contaminated clothing.•Rinse skin immediately with plenty of water for 15-20 minutes.•Call a poison control center, doctor, or 1-800-xxx-xxxx immediately for treatment advice.
If inhaled	<ul style="list-style-type: none">•Move person to fresh air.•If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.•Call a poison control center, doctor, or 1-800-xxx-xxxx immediately for treatment advice.
If in eyes	<ul style="list-style-type: none">•Hold eye open and rinse slowly and gently with water for 15-20 minutes.•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.•Call a poison control center, doctor, or 1-800-xxx-xxxx immediately for treatment advice.

C:\dpeacock\Dan Peacock\First Aid\Format for First Aid Statements2.wpdAugust 1, 2001

¹Toll free telephone number is optional throughout.

²Use all statements for each route of exposure if Toxicity Category is 1, 2, or 3. Statements are optional if Toxicity Category is 4.

**Format Labeling for Homeowner Disposal Statements
Non-Pressurized Container**

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or [toll free number which meets the criteria in paragraph II.E. of PR Notice 2002-6] for disposal instructions. Never place unused product down any indoor or outdoor drain.

