2 LBS.
NET WT.

DRAGON®

SEVIN* 50 WETTABLE

(for agricultural use only)

KILLS JAPANESE BEETLES

on certain listed crops

A wettable powder for the control of Mexican bean beetle, corn earworm, Colorado potato beetle and many other listed insects on listed crops.

*Trademark of UCC for 1-napthyl methylcarbamate

ACTIVE INGREDIENT	BY WT.
Carbaryl (1-napthyl methylcarbamate)	50%
INERT INGREDIENTS	50%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautionary statements

Manufactured for

DRAGON CHEMICAL CORPORATION

Roanoke, Virginia 24019

EPA REG. NO. 16-120 EPA Est. 16-VA-1

PRECAUTIONARY STATEMENTS

CAUTION

KEEP OUT OF REACH OF CHILDREN.

STATEMENT OF PRACTICAL TREATMENT

SEE RIGHT SIDE PANEL FOR ADDITIONAL CAUTIONS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

ENVIRONMENTAL HAZARDS

Avoid contamination of food, feed, water supplies, streams and ponds during application or when cleaning equipment. This product is highly toxic to bees exposed to direct treatment or residues. [illegible] Protective information may be obtained from your local Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Inform the nearby beekeepers [illegible] Application should be made when bees are [illegible] 2 days.

PHYSICAL OR CHEMICAL HAZARDS

[illegible] possible injury to tender foliage. [illegible] 2 days.

NOTICE

Buyer assumes all risks [illegible]

SEE LEFT SIDE PANEL FOR ADDITIONAL CAUTIONS

SEVIN 50 WETTABLE

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Do not store near feed or foodstuffs. DISPOSAL: Unused spray mixture or rinse should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited. Do not reuse empty containers. Destroy by burying in a landfill approved for pesticide containers or other safe place. Consult Federal, State or Local disposal authorities for approved alternative procedures such as controlled burning.

VEGETABLE CROP CONTROL

Recommended dosages refer to pounds of SEVIN 50 W per acre. Where a dosage range is indicated, use lower rate for young plants, higher rate on mature plants. Use sufficient spray gallonage to obtain full coverage. This will usually vary from 1 to 20 gallons per acre depending on equipment, degree of insect infestation and stage of crop growth. Keep spray mixture well agitated. Apply when insects or their damage appear and repeat at 7 to 14 days intervals as necessary unless shorter interval is specified below.

ASPARAGUS – Use 1 to 2 pounds for asparagus beetle on seedlings or spears; use 2 pounds for asparagus beetle and Asparagus Aphid in fern or for fern growth in the pre-harvest period. Do not apply more than [illegible] every 3 days. Do not apply within 1 day of harvest.

BEANS – Such as lima beans, green beans, snap beans, navy beans, pinto beans, snap beans and [illegible] including [illegible] and black eyed peas. Use [illegible] pound for Mexican bean beetle, 1 pound for bean leaf beetle, [illegible] bean beetles, flea beetles, [illegible] beetle, [illegible] cutworms, cowpea curculio and western flower thrips. [illegible] and alternaria leaf blight.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI – [illegible] pounds for flea beetles and [illegible] bugs, [illegible] pounds for harlequin bug. [illegible] dosages [illegible] 3 days of harvest.

CHINESE CABBAGE, COLLARDS, HORSERADISH, KALE, MUSTARD GREENS, RADISHES, RUTABAGAS AND TURNIPS – [illegible] pounds for flea beetles, [illegible] bugs, [illegible] pounds for harlequin bug. [illegible] dosages [illegible] 3 days of harvest.

CARROTS, PARSNIPS AND PARSLEY – [illegible] pound for flea beetles and leafhoppers. Use 1 to 2 pounds for [illegible] 5 to 7 day schedule if [illegible] pounds for harlequin bug. [illegible] dosages [illegible] 3 days of harvest. [illegible] 14 days of harvest. [illegible]

CORN – Sweet, seed and popcorn. Use 1 to 2 pounds for corn earworm, corn earworm adults, European corn borer, armyworm, flea beetles, Japanese beetle, leafhoppers, stalk borers, [illegible] and wireworms. Repeat as necessary. [illegible] 3 days. [illegible] 8 days. [illegible]

CUCUMBER, MELONS, PUMPKIN AND SQUASH – [illegible] pounds for pickleworm and [illegible] pounds for squash beetles, flea beetles, leafhoppers and squash bug. Avoid excessive applications. [illegible] 3 days.

ENDIVE (ESCAROLE), LETTUCE AND SALSIFY – [illegible] pounds for flea beetles, [illegible] bugs, [illegible] pounds for harlequin bug. [illegible] 3 days. [illegible]

GARDEN BEET, SPINACH AND SWISS CHARD – [illegible] pounds for 5 to 7 day schedule if [illegible] 3 days of harvest. [illegible] 14 days of harvest. [illegible]

OKRA – [illegible] 5 to 7 day [illegible]

PEAS – [illegible]

POTATO, TOMATO, EGGPLANT AND PEPPER – [illegible] 2 pounds for [illegible] beetles and leafhoppers. [illegible] 2 pounds for [illegible] bugs. [illegible]

SMALL FRUIT INSECT CONTROL

[illegible]

BLACKBERRIES, RASPBERRIES AND DEWBERRIES – [illegible]

BLUEBERRIES – [illegible]

GRAPES – [illegible]

STRAWBERRIES – [illegible]