

Ms. Rebecca M. Horton  
 Regulatory & Graphics Coordinator  
 Dragon Corporation  
 P.O. Box 7311  
 Roanoke, VA 24019

Dear Ms. Horton:

Subject: Labelling Submitted 2/23/90 in Compliance with the  
Bacillus thuringiensis Registration Standard  
 Dragon Sevin with Dipel Insect & Worm Dust  
 EPA Registration No. 16-121

The labelling to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable subject to the following comments.

1. Replace "Bacillus thuringiensis subspecies kurstaki Lepidopteran active toxin ..... 0.056%" with "Bacillus thuringiensis subspecies kurstaki Potency of 240 International Units per mg of Product (0.1089 IU/Pound of Product).... 0.048%" When your supplier of Bacillus thuringiensis modifies their label to list the percent lepidopteran toxin, you must modify your label to list the toxin percentage and submit a label amendment within 60 days.

2. Add the statement "Potency units should not be used to adjust use rates beyond those specified in the Directions for Use section." directly below the ingredient statement.

3. Replace the statement "Do not apply directly to water or wetlands..." with "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."

4. Replace "Sutabaga" with "Rutabaga."

CONCURRENCES

SYMBOL	H 205C	H7505C					
SURNAME	Morales	Kumar					
DATE	6/5/92	6/5/92					

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

2046

-2-

A stamped copy of the label is enclosed for your files. Submit five (5) final printed labels. Release of the revised labeling approved in this letter into commerce constitutes your acceptance of the conditions outlined in this letter.

Sincerely,

DHE

Dennis H. Edwards, Jr.  
Product Manager 19  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

*Handwritten notes:*  
✓ 1/11/92  
✓ 2/25/92  
✓ 4/11/92




# SEVIN®

brand carbaryl insecticide

## WITH DIPEL®

### INSECT & WORM DUST

Gives the broad spectrum insect control of SEVIN® brand carbaryl insecticide and the leaf chewing worm control of DIPEL®



**ACTIVE INGREDIENTS:**

Bacillus thuringiensis subsp. kurstaki*	
Lepidopteran active toxin	0.056%
Carbaryl (1-naphthyl-N-methyl carbamate)	5.000%

<b>INERT INGREDIENTS:</b>	94.944%
<b>TOTAL</b>	<u>100.000%</u>

\*Potency: 240 International Units per mg (1089 Billion International Units per pound of product)

**Keep Out of Reach of Children CAUTION**

See back/side panels for additional cautions.

**NET WEIGHT 4 LB**

JUN 11 1992

*Handwritten notes:*  
100-500-000  
100-500-000  
100-500-000

# DRAGON<sup>®</sup>

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

## STORAGE AND DISPOSAL

**Storage:** Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

**Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.

**Container Disposal:** Do not reuse empty container. Discard container in trash.

## THIS PRODUCT WILL CONTROL THE FOLLOWING PESTS ON HOME GARDEN VEGETABLES

Armyworms	Flea Beetle	Pickleworm
Bean Leaf Beetle	Marbled Bug	Sap Beetle
Cabbage Looper	Imported Cabbageworm	Spittlebug
Colorado Potato Beetle	Japanese Beetle	Squash Bug
Corn Earworm	Leafhoppers	Stink Bug
Cucumber Beetle	Lima Bean Pod Borer	Tarnished Plant Bug
Diamond Back Moth Larvae	Lygus Bug	Tomato Fruitworm
European Corn Borer	Melonworms	Tomato Hornworm
Fall Armyworm	Mexican Bean Beetle	

## FOLIAGE APPLICATION

Apply dust at the rate of 2 oz. per 50 feet of row (40 lbs. per acre) at the first sign of insects, using a good mechanical or handpowered duster. Repeat as necessary for control. For best results, dust early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. The number in ( ) following each crop indicates days required between last application and harvest; (0) indicates it may be used up to and including day of harvest.

### This product may be used on the following vegetables:

Beans (0)	Cucumbers (0)	Peppers (0)
Broccoli (0)	Eggplants (0)	Potatoes (0)
Bush Beans (root 3)	Garden Beans (root 3, top 14)	Pumpkins (0)
Cabbage (0)	Kale (14)	<del>Spinach (14)</del>
Cantaloupe (0)	Kohlrabi (3)	Swiss Chard (3)
Carrots (0)	Lettuce (head 3, leaf 14)	Squash (0)
Cauliflower (3)	Melons (0)	Tomato (0)
Celery (14)	Mustard Greens (14)	Turnips (root 3, top 14)
Corn (0)	Onion (0)	

To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

DRAGON is a registered trademark of Dragon Corporation, Roanoke, Virginia.  
EPA REG. NO. 16-121

Distributed By  
**DRAGON CORPORATION**  
Roanoke, VA 24019-0311

DRAGON is a registered trademark of Dragon Corporation, Roanoke, Virginia.

A EPA REG. NO. 16-VA-1

BEST AVAILABLE COPY

# DRAGON

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domes : Animals

**CAUTION:** Harmful if inhaled or swallowed. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin and eyes. Wear regular long sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

SEE LEFT SIDE PANEL FOR  
ADDITIONAL CAUTIONS

# DRAGON

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**IF INHALED:** Remove victim to fresh air. Apply respiration if indicated.

**IF ON SKIN:** Wash contaminated skin with soap and water.

**IF IN EYES:** Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

### NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEE RIGHT SIDE PANEL FOR  
ADDITIONAL CAUTIONS

# DRAGON

**SEVIN**<sup>®</sup>

brand carbaryl insecticide

WITH **DIPEL**<sup>®</sup>

INSECT & WORM DUST

NET WEIGHT 4 LB

UPC Code No.  
22001-08604