11-30-93

Please read Instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 11:30:93					
(A) SEPA United States Environment Office of Pesticide Pro Washington, D Application f	ograms (H7505C) Amendment				
	or Pesticide: Other 196857				
Se	ction I				
1. Company/Product Number 16-121	2. EPA Product Manager Edwards 3. Proposed Classification				
4. Company/Product (Name) DRAGON (R) SEVIN (R) with DIPEL (R) Insect and Worm Dust	PM# Restricted				
5. Name and Address of Applicant (Include ZIP Code)	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)				
Dragon Corporation	(b)(i), my product is similar or identical in composition and labeling to:				
P. O. Box 7311	io.				
Roanoke, VA 24019	EPA Reg. No				
Check if this is a new address	Product Name				
Sec	tion I i				
Amendment - Explain below	Final printed labels in response to Agency letter dated				
Resubmission in response to Agency letter dated	"Me Too" Application.				
XX Notification - Explain below.	Other - explain below.				
Explanation: Use additional page(s) if necessary. (For section	n I and Section II.)				
Compliance with R Notice 93-3	3 Labeling Statement Prohibiting Application to Water				
	MOTIFICATION				
	ARIA NOT PEVEN				
	PER PR MOTICE 7/12/93				
0	DATE.				
Section 1. Material This Product Will Be Packaged In:	QN III				
Child-Resistant Packaging Unit Packaging	Water Soluble Packaging 2. Type of Container				
☐ Yes* ☐ Yes	Yes Metal				
	- Plastic				
LXI No XI No Der	X Paper				
* Certification must be Unit Package wgt. Container	If "Yes," No. per Package wgt. Container Other (Specify)				
submitted.					
3. Location of Net Contents Information 4. Size(s) of	Retail Container 5. Location of Label Directions 7 On Label				
★ Label	On Labeling accompanying product				
6. Manner In Which Label Is Affixed To Product Lithog	graph Other ()				
Stendied					
Section IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
	Title Telephone No. (Include Area Code)				
Rebecca M. Horton	Executive Assistant (703)362-3657				
Certification 6. Date Application					
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or the statement of th					
2 Signature	3. Title				
Rebecca M. Hoston	Executive Assistant				
1 · ·	5. Date				
Rebecca M. Horton	6/21/93				

DRAGON® SEVIN®

brand carbaryl insecticide with DIPEL® INSECT & WORM DUST

NOTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 80-6
DATE. 7/12/93

Gives the broad spectrum insect control of SEVIN® brand carbaryl insecticide and the leaf chewing worm control of DIPEL®.

[NOTE TO EPA: Front panel of label may include graphics representing use, e.g., insects listed - Cabbage looper, Bean leaf beetle, Armyworms, Potato flea beetle.]

ACTIVE INGREDIENTS:

Keep Out of Reach of Children CAUTION

See back/side/other panel(s) for additional cautions.

NET WEIGHT 4 LB

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled or swallowed. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or, allow it to drift to blooming crops or weeds if bees are visiting the treatment.

EPA Reg. No. 16-121

June 4, 1993

EPA Est. 16-VA-1

Page 1

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash contaminated skin with soap and water.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

THIS PRODUCT WILL CONTROL THE FOLLOWING PESTS ON HOME GARDEN **VEGETABLES:**

Armyworms	Fall Armyworm	Mexican Bean Beetle
Bean Leaf Beetle	Flea Beetle	Pickleworm
Cabbage Looper	Harlequin Bug	Sap Beetle
Colorado Potato Beetle	Imported Cabbageworm	Spittlebug
Corn Earworm	Japanese Beetle	Squash Bug
Cucumber Beetle	Leafhoppers	Stink Bug
Diamond-back Moth	Lima Bean Pod Borer	Tarnished Plant Bug
Larvae	Lygus bug	Tomato Fruitworm
European Corn Borer	Melonworms	Tomato Hornworm

FOLIAGE APPLICATION

Apply dust at the rate of 2 oz. per 50 feet of row (40 lbs. per acre) at the first sign of insects, using a good mechanical or hand-powered duster. Repeat as necessary for control. For best results, dust early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. The number in () following each crop indicates days required between last application and harvest; (0) indicates it may be used up to and including day of harvest.

This product may be used on the following vegetables:

Beans (0)	Eggplants (0)	Peppers (0)
Broccoli (3)	Garden Beets	Potatoes (0)
Brussels Sprouts (3)	(roots 3, tops 14)	Pumpkins (0)
Cabbage (3)	Kale (14)	Radishes (3)
Cantaloupe (0)	Kohlrabi (3)	Rutabaga (3)
Carrots (0)	Lettuce	Spinach (14)
Cauliflower (3)	(head 3, leaf 14)	Squash (0)
Collards (14)	Melons (0)	Tomato (0)
Corn (0)	Mustard Greens (14)	Turnips
Cucumbers (0)	Okra (0)	(roots 3, tops 14)

To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN[®] is a registered trademark of Rhóne Poulenc Ag Company. DIPEL[®] is a registered trademark of Abbott Laboratories.

Distributed by DRAGON CORPORATION Roanoke, VA 24019

Label of Record - as submitted under 6/25/92*

notification

DRAGON® SEVIN®

brand carbaryl insecticide with DIPEL® INSECT & WORM DUST

EPA Reg. No. 16-121

EPA Est. No. 16-VA-1

Gives the broad spectrum insect control of SEVIN® brand carbaryl insecticide and the leaf chewing worm control of DIPEL®.

[NOTE · TO EPA: Front panel of label may include graphics representing use, e.g., insects listed - Cabbage looper, Bean leaf beetle, Armyworms, Potato flea beetle.]

ACTIVE INGREDIENTS:

Bacillus thuringiensis subspecies <u>kurstaki</u>
Potency of 240 International Units per
mg. of Product (0.1089 Billion IU/Pound of

Potency units should not be used to adjust use rates beyond those specified in the Directions for Use section.

Keep Out of Reach of Children CAUTION

See back/side/other panel(s) for additional cautions.

NET WEIGHT 4 LB

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled or swallowed. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment.

EPA Reg. No. 16-121 June 25, 1992

10041

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
- IF INHALED: Remove victim to fresh air. Apply respiration if indicated.
- IF ON SKIN: Wash contaminated skin with soap and water.
- IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

NOTE \cdot TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

THIS PRODUCT WILL CONTROL THE FOLLOWING PESTS ON HOME GARDEN VEGETABLES:

Armyworms
Bean Leaf Beetle
Cabbage Looper
Colorado Potato Beetle
Corn Earworm
Cucumber Beetle
Diamond-back Moth
Larvae

European Corn Borer

Fall Armyworm
Flea Beetle
Harlequin Bug
Imported Cabbageworm
Japanese Beetle
Leafhoppers
Lima Bean Pod Borer
Lygus bug
Helonworms

Mexican Bean Beetle
Pickleworm
Sap Beetle
Spittlebug
Squash Bug
Stink Bug
Tarnished Plant Bug
Tomato Fruitworm
Tomato Hornworm

EPA Reg. No. 16-121 June 25, 1992 EPA Est. No. 16-VA-1 Page 2

FOLIAGE APPLICATION

Apply dust at the rate of 2 oz. per 50 feet of row (40 lbs. per acre) at the first sign of insects, using a good mechanical or hand-powered duster. Repeat as necessary for control. For best results, dust early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. The number in () following each crop indicates days required between last application and harvest; (0) indicates it may be used up to and including day of harvest.

This product may be used on the following vegetables:

Beans (0)	Eggplants (0)	Peppers (0)
Broccoli (3)	Garden Beets	Potatoes (0)
Brussels Sprouts (3)	(roots 3, tops 14)	Pumpkins (0)
Cabbage (3)	Kale (14)	Radishes (3)
Cantaloupe (0)	Kohlrabi (3)	Rutabaga (3)
Carrots (0)	Lettuce	Spinach (14)
Cauliflower (3)	(head 3, leaf 14)	Squash (0)
Collards (14)	Melons (0)	Tomato (0)
Corn (0)	Mustard Greens (14)	Turnips
Cucumbers (0)	Okra (0)	(roots 3, tops 14)

To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhone Poulenc Ag Company. DIPEL® is a registered trademark of Abbott Laboratories.

Distributed by DRAGON CORPORATION Roanoke, VA 24019

