

[FRONT PANEL]

**DRAGON®
SEVIN®**
brand carbaryl insecticide
with **DIPEL® INSECT & WORM DUST**

Nothing Beats The Dragon!

Gives the broad spectrum insect control of SEVIN® brand carbaryl insecticide and the leaf chewing worm control of DIPEL®.

[NOTE TO EPA: Front panel of label may include graphics representing use, e.g., insects listed - Cabbage looper, Bean leaf beetle, Armyworms, Potato flea beetle.]

ACCEPTED
FEB 24 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 16-121

ACTIVE INGREDIENTS:

Bacillus thuringiensis subspecies kurstaki
Potency of 240 International Units per mg. of Product (0.1089 Billion IU/Pound of Product) 0.048%
Carbaryl (1-naphthyl N-methylcarbamate) 5.000%

INERT INGREDIENTS: 94.952%

TOTAL 100.000%

Potency units should not be used to adjust use rates beyond those specified in the Directions for Use section.

Keep Out of Reach of Children

CAUTION

See back/side/other panel(s) for additional cautions.

NET WEIGHT 4 LB

[BACK/SIDE/OTHER PANEL(S)]

FIRST AID

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash contaminated skin with soap and water.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

REC'D
NOV 20 P 3:23
PPD1

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled or swallowed. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin and eyes.

USER SAFETY REQUIREMENTS
<ul style="list-style-type: none"> • Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. • Change clothing as soon as possible after use. • Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. • Do not allow children or pets to contact treated area until dust has settled.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment.

THIS PRODUCT WILL CONTROL THE FOLLOWING PESTS ON HOME GARDEN VEGETABLES:

Armyworms	Fall Armyworm	Mexican Bean Beetle
Bean Leaf Beetle	Flea Beetle	Pickleworm
Cabbage Looper	Harlequin Bug	Sap Beetle
Colorado Potato Beetle	Imported Cabbageworm	Spittlebug
Corn Earworm	Japanese Beetle	Squash Bug
Cucumber Beetle	Leafhoppers	Stink Bug
Diamond-back Moth	Lima Bean Pod Borer	Tarnished Plant Bug
Larvae	Lygus bug	Tomato Fruitworm
European Corn Borer	Melonworms	Tomato Hornworm

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply at the rate of **2 oz. per 50 feet of row** when insects first appear before damage from feeding occurs. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid insecticide. Do not repeat application more than once a week. Do not use with lime or other alkaline materials. This product may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. The number in () following each crop

indicates days required between last application and harvest; (0) indicates it may be used up to and including day of harvest.

This product may be used on the following vegetables:

Beans (0)	Eggplants (0)	Peppers (0)
Broccoli (3)	Garden Beets	Potatoes (0)
Brussels Sprouts (3)	(roots 3, tops 14)	Pumpkins (0)
Cabbage (3)	Kale (14)	Radishes (3)
Cantaloupe (0)	Kohlrabi (3)	Rutabaga (3)
Carrots (0)	Lettuce	Spinach (14)
Cauliflower (3)	(head 3, leaf 14)	Squash (0)
Collards (14)	Melons (0)	Tomato (0)
Corn (0)	Mustard Greens (14)	Turnips
Cucumbers (0)	Okra (0)	(roots 3, tops 14)

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

DIPEL® is a registered trademark of Abbott Laboratories.

Distributed by
DRAGON CORPORATION
Roanoke, VA 24019