

Copper Dragon

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS
CAUTION


Skin contact may be harmful.
Wear long sleeved work clothing.

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

NOTICE

Copper Dragon



BUG AND BLIGHT DUST

Protects your garden from many of the most common pests AND diseases.

KILLS Japanese Beetles, Fleas, Beetles, Tomato Hornworm, flea beetle, and many other insects.

CONTROLS many of the most common blights on Tomatoes and Potatoes - White Mold, Black Mold, Anthracnose, Bacterial Wilt, and many other plant diseases.

ACTIVE INGREDIENTS

Copper, Zinc, and other active ingredients.

INERT INGREDIENTS

CAUTION

KEEP OUT OF REACH OF CHILDREN

See left side panel for additional cautions

Coppe Drago




NET WT. 4 LB.

Copper Dragon

SEP 8 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. 160.110



BRIGHT DUST

any of the most common pests AND

Bean beetle, tomato hornworm, flea

light, Leaf Spot, in Tomatoes and Spot, Anthracnose, Bacterial Wilt on

carbamate)

UTION

EACH OF CHILDREN

for additional cautions

WT. 4 LB.

Copper Dragon



Copper Dragon

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL: Store in a cool, dry place away from feed or foodstuffs. Do not reuse empty container - wrap in trash collection.

COPPER DRAGON Bug & Bright Dust is an effective combination insecticide fungicide dust that controls many insects and diseases at the same time. Apply when insects first appear. For diseases apply when plants are young and BEFORE damage occurs. Dust thoroughly to upper and lower surfaces. If needed repeat every 7-10 days. Do not use with lime or other alkaline material. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. **COPPER DRAGON** may be applied up to harvest except when noted.

BEANS: For control of Mexican bean beetle, leafhopper, flea beetles, Japanese beetles, velvet bean caterpillar, apply 1 1/2 lb. per 1,000 sq. ft. For control of virus, stink bug, horn worm, tarnished plant bug, apply 1 1/2 to 2 lb. per 1,000 sq. ft.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For control of Colorado potato beetle, flea beetles, leafhoppers, Leaf Spot, Early and Late Blight on potatoes and tomatoes and Cercospora (Leaf Spot), Anthracnose on eggplant and peppers, apply 1 1/2 to 2 lb. per 1,000 sq. ft. For control of Colorado potato beetle, armyworm, tomato hornworm, European corn borer, stink bugs, leaf bugs, virus bugs, apply 1 1/2 to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: Control of watermelon, squash, melon, cucumber, pumpkin, melonworm, cucumber beetle, squash bug, flea beetle, leafhoppers, Anthracnose, Bacterial Wilt, Downy Mildew, Virus. After harvest apply 1 1/2 to 2 lb. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH: For control of flea beetles, leafhoppers, harlequin bug, Downy mildew, leaf spot, apply 1 1/2 to 2 lb. per 1,000 sq. ft. For six spotted leafhoppers, corn earworm, armyworm, cabbage caterpillar, spotted burr, lygus and stink bugs, apply 1 1/2 to 2 lb. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce).

ASPARAGUS: For control of asparagus beetle, snipe, apply 1 1/2 to 2 lb. per 1,000 sq. ft. Apply during cutting season in 2 to 3 equal applications within three days of cutting.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS: For control of flea beetle, harlequin bug, leafhopper, European corn borer, apply 1 1/2 to 2 lb. per 1,000 sq. ft. For control of armyworm, spotted burr, apply 1 1/2 to 2 lb. per 1,000 sq. ft.

ROSES, AZALEA, AND OTHER ORNAMENTALS: For control of Japanese beetle, European corn borer, flea beetle, leafhopper, apply 1 1/2 to 2 lb. per 1,000 sq. ft.

GRAPES: For control of grape leafhopper, apply 1 1/2 to 2 lb. per 1,000 sq. ft.

STRAWBERRIES: For control of strawberry root weevil, apply 1 1/2 to 2 lb. per 1,000 sq. ft.

For additional information, contact your local extension agent or write to: Dragon Chemical Corporation, Roanoke, Virginia 24019.

Manufactured By

DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24019