

173052
 17/03
 300

Mr. J.C. Wilson
 Dragon Chemical Corporation
 P.O. Box 7311
 Roanoke, VA 24019

JUL 11 1986

Dear Mr. Wilson

Subject: General Revision of Label
 Copper Dragon Bug & Blight Dust
 EPA Registration Number 16-127
 Your Application Dated April 11, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At your next printing of labeling for this product, submit five (5) copies of revised draft labeling incorporating the comments listed below.

1. Our June 9, 1986 letter was incorrect regarding the substatement required for copper sulfate. The substatement should read:

Derived from Basic Copper Sulfate

2. Add to your Environmental Hazards section the paragraph:

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Sincerely yours,

LJS
 Lawrence J. Schnaubelt
 Acting Product Manager (12)
 Insecticide-Rodenticide Branch
 Registration Division (TS-767C)

		CONCURRENCES					
INBOLE	Enclosure						
SURNAME	92967:Schnaubelt:E-11:KENCO: 6/18/86:						
DATE							



COPPER DRAGON®



GRAPES — For control of leathoppers, Japanese beetles, grape berry moth, June beetles, Black rot, Downy and Powdery mildew, apply $\frac{3}{4}$ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES — For control of meadow spittle bug, leaf roller, strawberry weevil, leaf spot, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

EPA Reg. No. 16 127

EPA EST. 16 VA 1

*SEVIN is a registered trademark of Union Carbide Corporation

COPPER DRAGON

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store only in original container, in a cool dry place inaccessible to children and pets and away from feed and foodstuffs.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash. **Container Disposal:** Do not reuse empty bag. Discard bag in trash.

COPPER DRAGON Bug & Blight Dust is an effective combination insecticide-fungicide dust that controls many insects and diseases at the same time. Apply when insects first appear. For diseases apply when plants are young and **BEFORE** damage occurs. Dust thoroughly to upper and lower surfaces; if needed, repeat every 7-10 days. Do not use with lime or other alkaline material. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. **COPPER DRAGON** may be applied up to harvest except when noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS — For control of Mexican bean beetle, leathopper, flea beetles, Japanese Beetles, velvet bean caterpillar, apply $\frac{3}{4}$ lb. per 1,000 sq. ft. For control of lygus, stink bug, corn earworm, tarnished plant bug, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS — For control of Colorado potato beetle, flea beetles, leathoppers, Leaf Spot, Early and Late Blight on potatoes and tomatoes and Cercospora (Leaf Spot), Anthracnose on eggplant and peppers, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs, lygus bugs, apply $1\frac{1}{2}$ lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, watermelon, honey dew) — For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetle, leathoppers, Anthracnose, Bacterial Wilt, Downy Mildew, Scab, Alternate Leaf Spot, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH — For control of flea beetles, leathoppers, harlequin bug, Downy mildew, leaf spot, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For six-spotted leathoppers, corn earworm, armyworm, cabbage caterpillar, spittlebugs, lygus and stink bugs, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce — 3 days).

ASPARGUS — For control of asparagus beetle on spears, apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft. Apply during cutting season; do not repeat application within three days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS — For control of flea beetles, harlequin bugs, leaf spot, Downy Mildew, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per sq. ft. For corn earworm, stink bug, armyworm, cabbage caterpillar, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEA AND OTHER ORNAMENTALS — For control of Japanese beetle, bagworm, flea beetles, leathoppers, scale insects, tent caterpillars and many others, apply 2 to $2\frac{1}{4}$ lbs. per 1,000 sq. ft. SEE RIGHT SIDE PANEL FOR ADDITIONAL DIRECTIONS.

Manufactured By

DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 240



COPPER DRAGON

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if inhaled or swallowed. Do NOT breathe dust. Do NOT take internally. Skin contact may be harmful. Wear gloves when applying this product. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing after use. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, promptly drink a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Call a physician immediately. If on skin, wash thoroughly with soap and water. If in eyes, immediately flush eyes with plenty of water for at least 15 minutes; call a physician.


ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds if bees are visiting the treatment area. This product is toxic to fish. Keep out of lakes, streams, or ponds.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

COPPER DRAGON



BUG AND BLIGHT DUST

Protects your garden from many of the most common pests AND diseases

EXCELLENT TOMATO PROTECTION

KILLS Japanese beetle, Mexican Bean beetle, tomato hornworm, flea beetle and other listed insects

CONTAINS: **SEVIN**®*

CONTROLS Early and Late Blight, Leaf Spot on Tomatoes and Potatoes, Downy Mildew, Black Spot, Anthracnose, Bacterial Wilt on cucumbers, squash and melons

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl N-methylcarbamate)	2%
Copper Expressed as Metallic (Derived from Copper Sulfate)	7%
INERT INGREDIENTS	91%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See left side panel for additional cautions

NET WEIGHT 4 LB.