

PM 19 16-127 127

Please read instructions on reverse before completing form.

Form Approved OMB No 2070-0060 Approval expires 11-30-93

	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number 188524
	Application for Pesticide:		

Section I

1. Company/Product Number 16-127	2. EPA Product Manager Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Copper Dragon (R) Bug & Blight Dust	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Dragon Corporation P. O. Box 7311 Roanoke, VA 24019 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Compliance with PR Notice 93-3 -- Labeling Statement ~~for Registration~~
 Prohibiting Application to Water

**NOTIFICATION
 LABEL NOT REVIEWED
 PER PR NOTICE 93-3**

DATE: 7/17/93

Section III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container		
* Certification must be submitted.			<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container 14oz / 4LB / 25LB		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed To Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other (_____)		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Rebecca M. Horton	Title Executive Assistant	Telephone No. (include Area Code) (703) 362-3657	
2. Signature 			6. Date Application Received (Stamped)
3. Title Executive Assistant		4. Typed Name Rebecca M. Horton	
5. Date 6/21/93			

2071

**DRAGON®
COPPER DRAGON® BUG & BLIGHT DUST**

Protects your garden from many of the most common pests and diseases.

**NOTIFICATION
LABEL NOT REVISION
PER PR NOTICE 02-1**

EXCELLENT TOMATO PROTECTION.

DATE 7/12/93

Contains SEVIN® brand carbaryl insecticide.

KILLS: Japanese Beetle, Mexican Bean Beetle, Tomato Hornworm, Flea Beetle and other listed insects.

CONTROLS: Early and Late Blight, Leaf Spot on tomatoes and potatoes, Downy Mildew, Black Spot, Anthracnose, Bacterial Wilt on cucumbers, squash and melons.

Active Ingredients:

Carbaryl (1-Naphthyl N-methylcarbamate)	2%
Copper Expressed as Metallic	7%
(Derived from Basic Copper Sulfate)	

Inert Ingredients:	<u>91%</u>
TOTAL	100%

**Keep Out of Reach of Children
CAUTION
See back/side panels for additional cautions.**

Net WEIGHT 4 LB / 25 LB

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

COPPER DRAGON® Bug & Blight Dust is an effective combination insecticide-fungicide dust that controls many insects and diseases at the same time. Apply when insects first appear. For diseases apply when plants are young and **BEFORE** damage occurs. Dust thoroughly to upper and lower surfaces. If needed, repeat every 7-10 days. Do not use with lime or other alkaline material. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. **COPPER DRAGON®** may be applied up to harvest except when noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: For control of Mexican bean beetle, Leafhopper, Flea beetles, Japanese beetles, Velvet bean caterpillar, apply 3/4 lb. per 1,000 sq. ft. For control of Lygus, Stink bug, Corn ear worm, Tarnished plant bug, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For control of Colorado potato beetle, Flea beetles, Leafhoppers, Leaf spot, early and late Blight on potatoes and tomatoes and Cercospora (leaf spot), Anthracnose on eggplant and peppers, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For control of Tomato fruitworm, Fall armyworm, Tomato hornworm, European corn borer, Stink bugs, Lace bugs, Lygus bugs, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, watermelon, honeydew): For control of Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetle, Leafhoppers, Anthracnose, Bacterial wilt, Downy mildew, Scab, Alternate leaf spot, apply $\frac{1}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH: For control of Flea beetles, Leafhoppers, Harlequin bug, Downy mildew, Leaf spot, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For Six-spotted Leafhoppers, Corn earworm, Armyworm, Cabbage caterpillar, Spittlebugs, Lygus and Stink bugs, apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce — 3 days).

ASPARAGUS: For control of Asparagus beetle on spears - Apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft. Apply during cutting season; do not repeat application within three days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS: For control of Flea beetles, Harlequin bugs, Leaf spot, Downy mildew - Apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per sq. ft. For Corn earworm, Stink bug, Armyworm, Cabbage caterpillar - Apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEA AND OTHER ORNAMENTALS: For control of Japanese beetle, Bagworm, Flea beetles, Leafhoppers, Scale insects, Tent caterpillars and many others - Apply 2 to $2\frac{1}{4}$ lbs. per 1,000 sq. ft.

GRAPES: For control of Leafhoppers, Japanese beetles, Grape berry moth, June beetles, Black rot, Downy and Powdery mildew - Apply $\frac{3}{4}$ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES: For control of Meadow spittle bug, Leaf roller, Strawberry weevil, Leaf spot - Apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Wear gloves when applying this product. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing after use. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

10/27/93

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Promptly drink a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Call a physician immediately.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes, call a physician.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to fish. Keep out of lakes, streams or ponds. This product is extremely toxic to aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE: Store only in original container, in a cool dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

Label of Record - as submitted under 9-9-92. 5071
Notification

**DRAGON® COPPER DRAGON®
BUG & BLIGHT DUST**

EPA Reg. No. 16-127 EPA Est. No. 16-VA-1

Protects your garden from many of the most common pests and diseases.

EXCELLENT TOMATO PROTECTION.

Contains SEVIN® brand carbaryl insecticide.

KILLS: Japanese Beetle, Mexican Bean Beetle, Tomato Hornworm, Flea Beetle and other listed insects.

CONTROLS: Early and Late Blight, Leaf Spot on tomatoes and potatoes, Downy Mildew, Black Spot, Anthracnose, Bacterial Wilt on cucumbers, squash and melons.

Active Ingredients:

Carbaryl (1-Naphthyl N-methylcarbamate)	2%
Copper Expressed as Metallic	7%
(Derived from Basic Copper Sulfate)	

Inert Ingredients:	<u>91%</u>
TOTAL	100%

**Keep Out of Reach of Children
CAUTION
Net WEIGHT 4 LBS / 25 LBS**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. DO NOT breathe dust. DO NOT take internally. Skin contact may be harmful. Wear gloves when applying this product. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing after use. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

STATEMENT OF PRACTICAL TREATMENT

If swallowed, promptly drink a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Call a physician immediately.
If on skin, wash thoroughly with soap and water.
If in eyes, immediately flush eyes with plenty of water for at least 15 minutes, call a physician.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds if bees are visiting the treatment area. This product is toxic to fish. Keep out of lakes, streams or ponds. This product is extremely toxic to aquatic invertebrates. Do not apply directly to water, areas where surface water is present, or areas below the mean high tidal mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

6071

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store only in original container, in a cool dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

COPPER DRAGON Bug & Blight Dust is an effective combination insecticide-fungicide dust that controls many insects and diseases at the same time. Apply when insects first appear. For diseases apply when plants are young and **BEFORE** damage occurs. Dust thoroughly to upper and lower surfaces. If needed, repeat every 7-10 days. Do not use with lime or other alkaline material. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. **COPPER DRAGON** may be applied up to harvest except when noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

BEANS: For control of Mexican bean beetle, Leafhopper, Flea beetles, Japanese beetles, Velvet bean caterpillar, apply 3/4 lb. per 1,000 sq. ft. For control of Lygus, Stink bug, Corn earworm, Tarnished plant bug, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS: For control of Colorado potato beetle, Flea beetles, Leafhoppers, Leaf spot, early and late Blight on potatoes and tomatoes and Cercospora (leaf spot), Anthracnose on eggplant and peppers, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft. For control of Tomato fruitworm, Fall armyworm, Tomato hornworm, European corn borer, Stink bugs, Lace bugs, Lygus bugs, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, watermelon, honeydew): For control of Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetle, Leafhoppers, Anthracnose, Bacterial wilt, Downy mildew, Scab, Alternate leaf spot, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH: For control of Flea beetles, Leafhoppers, Harlequin bug, Downy mildew, Leaf spot, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft. For Six-spotted Leafhoppers, Corn earworm, Armyworm, Cabbage caterpillar, Spittlebugs, Lygus and stink bugs, apply 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce — 3 days).

ASPARAGUS: For control of Asparagus beetle on spears - Apply 1 1/4 to 2 1/2 lbs. per 1,000 sq. ft. Apply during cutting season; do not repeat application within three days. Do not apply within 1 day of harvest.

7/10/77

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS: For control of Flea beetles, Harlequin bugs, Leaf spot, Downy mildew - Apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per sq. ft. For Corn earworm, Stink bug, Armyworm, Cabbage caterpillar - Apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEA AND OTHER ORNAMENTALS: For control of Japanese beetle, Bagworm, Flea beetles, Leafhoppers, Scale insects, Tent caterpillars and many others - Apply 2 to $2\frac{1}{4}$ lbs. per 1,000 sq. ft.

GRAPES: For control of Leafhoppers, Japanese beetles, Grape berry moth, June beetles, Black rot, Downy and Powdery mildew - Apply $\frac{3}{4}$ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES: For control of Meadow spittle bug, Leaf roller, Strawberry weevil, Leaf spot - Apply $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019