

110 13  
14  
SEP 27 1993

Ms. Rebecca M. Horton  
Dragon Corporation  
P.O. Box 7311  
Roanoke, VA 24019

Dear Ms. Horton:

Subject: Copper Dragon Bug & Blight Bust  
EPA Registration No. 16-127  
Your submission dated 9/9/93  
General Label Revisions

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Standard (WPS). Under the WPS labeling regulations at 40 CFR part 156, subpart K, §156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.  
Product Manager (19)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

[FRONT PANEL]

**DRAGON®  
COPPER DRAGON® BUG & BLIGHT DUST**

[Additional brand names: Copper Dragon Tomato & Vegetable Dust, Dragon Tomato & Vegetable Dust II]

*Nothing Beats The Dragon!*

Ready-to-Use!

Protects your garden from many of the most common pests and diseases.

**EXCELLENT TOMATO PROTECTION.**

Contains SEVIN® brand carbaryl insecticide.


**KILLS:** Japanese Beetle, Mexican Bean Beetle, Tomato Hornworm, Flea Beetle and other listed insects.

**CONTROLS:** Early and Late Blight, Leaf Spot on tomatoes and potatoes, Downy Mildew, Black Spot, Anthracnose, Bacterial Wilt on cucumbers, squash and melons.

**Active Ingredients:**

Carbaryl (1-Naphthyl N-methylcarbamate) . . . . .	2%
Copper Expressed as Metallic . . . . .	7%
(Derived from Basic Copper Sulfate)	

<b>Inert Ingredients:</b> . . . . .	91%
<b>TOTAL</b> . . . . .	100%



**Keep Out of Reach of Children**

**CAUTION**

**See back/side panels for additional cautions.**

**NET WEIGHT 14 OZ / 4 LB / 25 LB**

[BACK/SIDE/OTHER PANEL(S)]

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**COPPER DRAGON® Bug & Blight Dust** is an effective combination insecticide-fungicide dust that controls many insects and diseases at the same time. Apply when insects first appear. For diseases apply when plants are young and **BEFORE** damage occurs. Dust thoroughly to upper and lower surfaces. If needed, repeat every 7-10 days. Do not use with lime or other alkaline material. To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. **COPPER DRAGON®** may be applied up to harvest except when noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.



**BEANS:** For control of Mexican bean beetle, Leafhopper, Flea beetles, Japanese beetles, Velvet bean caterpillar, apply  $\frac{3}{4}$  lb. per 1,000 sq. ft. For control of Lygus, Stink bug, Corn earworm, Tarnished plant bug, apply  $1\frac{1}{4}$  to 2 lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to 1 lb. per 500 sq. ft.).

**POTATOES, TOMATOES, EGGPLANT AND PEPPERS:** For control of Colorado potato beetle, Flea beetles, Leafhoppers, Leaf spot, early and late Blight on potatoes and tomatoes and Cercospora (leaf spot), Anthracnose on eggplant and peppers, apply  $\frac{3}{4}$  to  $1\frac{1}{4}$  lbs. per 1,000 sq. ft. For control of Tomato fruitworm, Fall armyworm, Tomato hornworm, European corn borer, Stink bugs, Lace bugs, Lygus bugs, apply  $1\frac{1}{4}$  to 2 lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to 1 lb. per 500 sq. ft.).

**CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, watermelon, honeydew):** For control of Pickleworm, Melonworm, Cucumber beetles, Squash bugs, Flea beetle, Leafhoppers, Anthracnose, Bacterial wilt, Downy mildew, Scab, Alternate leaf spot, apply  $\frac{3}{4}$  to  $1\frac{1}{4}$  lbs. per 1,000 sq. ft.

**LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH:** For control of Flea beetles, Leafhoppers, Harlequin bug, Downy mildew, Leaf spot, apply  $\frac{3}{4}$  to  $1\frac{1}{4}$  lbs. per 1,000 sq. ft. For Six-spotted Leafhoppers, Corn earworm, Armyworm, Cabbage caterpillar, Spittlebugs, Lygus and Stink bugs, apply  $1\frac{1}{4}$  to 2 lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to 1 lb. per 500 sq. ft.). Do not apply within 14 days of harvest (except head lettuce — 3 days).

**ASPARAGUS:** For control of Asparagus beetle on spears - Apply  $1\frac{1}{4}$  to  $2\frac{1}{2}$  lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to  $1\frac{1}{4}$  lbs. per 500 sq. ft.) Apply during cutting season; do not repeat application within three days. Do not apply within 1 day of harvest.

**CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS:** For control of Flea beetles, Harlequin bugs, Leaf spot, Downy mildew - Apply  $\frac{3}{4}$  to  $1\frac{1}{4}$  lbs. per 1,000 sq. ft. For Corn earworm, Stink bug, Armyworm, Cabbage caterpillar - Apply  $1\frac{1}{4}$  to 2 lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to 1 lb. per 500 sq. ft.) Do not apply within 3 days of harvest.

**ROSES, AZALEA AND OTHER ORNAMENTALS:** For control of Japanese beetle, Bagworm, Flea beetles, Leafhoppers, Scale insects, Tent caterpillars and many others - Apply 2 to  $2\frac{1}{4}$  lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  lb. per 250 sq. ft.).

**GRAPES:** For control of Leafhoppers, Japanese beetles, Grape berry moth, June beetles, Black rot, Downy and Powdery mildew - Apply  $\frac{3}{4}$  to 2 lbs. per 1,000 sq. ft.

**STRAWBERRIES:** For control of Meadow spittle bug, Leaf roller, Strawberry weevil, Leaf spot - Apply  $1\frac{1}{4}$  to 2 lbs. per 1,000 sq. ft. ( $\frac{1}{2}$  to 1 lb. per 500 sq. ft.). Do not apply within 1 day of harvest.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Wear gloves when applying this product. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing after use. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Promptly drink a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Call a physician immediately.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF IN EYES:** Immediately flush eyes with plenty of water for at least 15 minutes, call a physician.

### ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to fish. Keep out of lakes, streams or ponds. This product is extremely toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**STORAGE:** Store only in original container, in a cool dry place inaccessible to children and pets and away from feed and foodstuffs.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Discard bag in trash.

**STAINING PRECAUTION:** Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Manufactured by  
**DRAGON CORPORATION**  
Roanoke, VA 24019

