

DRAGON®

FRUIT TREE SPRAY

ACCEPTED
SEP 26 1988
76-129

CONTAINS:
METHOXYCHLOR, MALATHION, CAPTAN.
A COMPLETE INSECTICIDE,
FUNGICIDE AND MITICIDE
FOR INSECT AND DISEASE CONTROL
ON FRUIT TREES.

BEST AVAILABLE COPY

ACTIVE INGREDIENTS	
Methoxychlor, Technical	10.00%
Captan**	8.62%
Related Derivatives	0.36%
Malathion — 0,0-dimethyl dithiophos- phate of diethyl mercaptosuccinate	5.00%
INERT INGREDIENTS	78.00%
	100.00%

*Equivalent to 8.8% 2,2-bis (p-methoxyphenyl) 1,1-trichloroethane
and 1.2% other isomers and related compound.

**N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER

See back panel for additional precautionary statements.

NET WEIGHT 1 LB.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Captan causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Avoid inhalation of dust or spray mist. Use with adequate ventilation.

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything to an unconscious person. Get medical attention. **If inhaled** — Remove victim to fresh air. Apply respiration if indicated. **If on skin** — Wash contaminated skin with soap and water. **In eyes** — Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

ENVIRONMENTAL HAZARDS — This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

ENDANGERED SPECIES PRECAUTIONS — The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

Stackwater Darter	Alabama (Lauderdale, Limestone, and Madison) Tennessee (Lawrence and Wayne)
Freshwater Mussels	Tennessee (Hancock, Claiborne, Hawkins and Sullivan) Virginia (Smyth, Scott, Lee, Washington and Grayson)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with DIRECTIONS FOR USE and PRECAUTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

STORAGE: Do not store near feed or foodstuffs. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse empty container. Discard container in trash.

Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

SPRAY APPLICATION: One pound of this material will make 10 gallons of fruit tree spray. For smaller quantities, use 5 level tablespoons to each gallon of water. Mix well. Agitate solution while spraying. (DRAGON FRUIT TREE SPRAY may be applied as a dust by use of an efficient duster and directing dust to all portions of the tree being treated.)

AMOUNT OF SPRAY NEEDED PER TREE:

SIZE	DILUTED SPRAY
Young or Dwarf	Approx. 1 to 2 pints
Medium	Approx. 1 to 2 quarts

APPLES, PEARS: To control scab, brooks fruit spot, woolly and green apple aphids, pacific clover and red spider mites, codling moth, curculio, apple maggot, Japanese beetle, and bitter rot, apply as follows:

1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals to 21 days before harvest.

PEACHES: To control brown rot, peach scab, European red mite, red spider, plum curculio, Japanese beetle, European fruit scale, pacific and clover mites, apply as follows:

1. Full bloom.
2. When shucks begin to split.
3. 10 days later.
4. 10 days later.
5. 2 weeks later.
6. Do not apply within 21 days of harvest.

PLUMS, PRUNES: To control pacific, European and red spider mites, plum curculio, mealy plum and green peach aphids and Japanese beetle, apply as follows:


1. When trees are in full bloom.
2. When shucks are falling.
3. 10 days later.
4. 2 weeks later.
5. Continue at 2-week intervals if necessary. Do not apply later than 21 days before harvest.

If control of disease on above fruit crops is necessary within 21 days of harvest, apply a wettable sulfur spray. Remove residues by washing, wiping, brushing or other effective means.

EPA Reg. No. 16-129

EPA Est. 16-VA-1

Manufactured by
DRAGON CHEMICAL CORPORATION
Roanoke, Virginia 24019



BEST AVAILABLE COPY