

JUN 20 1991

Ms. Rebecca M. Horton
Dragon Corporation
P.O. Box 7311
Roanoke, VA 24019

Dear Ms. Horton:

Subject: Label amendments
Dragon Ready To Use Garden Pest Control
EPA Registration Number 16-132
Submission dated September 10, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

1. Delete the logo- Natural Organic Recommended For Organic Gardeners. This pesticide is produced from synthesized material and it is not clear who recommends it.
2. Delete petroleum distillate as an active ingredient and include its percentage with the total figure for INERT INGREDIENTS.
3. Revise the ENVIRONMENTAL HAZARDS statement to read:

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes, and estuaries. Do not contaminate water when disposing of equipment washwaters. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

EPA Registration Number 16-132:

Submit an updated CONFIDENTIAL STATEMENT OF FORMULA for the record.

Sincerely,

 Richard F. Mountfort
Product Manager 10
Insecticide-Rodenticide Branch
Registration Division (H7505C)

3/4
DRAGON

Ready to Use!

GARDEN PEST CONTROL

Fast Acting Organic Spray for Vegetables, Flowers and Ornamentals



ACTIVE INGREDIENTS

INERT INGREDIENTS
TOTAL

Keep Out of Reach of Children

CAUTION

See back panel for additional precautions

NET CONTENTS 32 FL OZ

Kills: Aphids, Whiteflies, Beetles, Leafhoppers, Stink Bugs, Caterpillars and many other listed insects.

Controls Insects on Ornamentals and Delicate House Plants.

May be applied on day of harvest.

Apply with conventional mechanical compressed air equipment or non-aerosol pump.

Do not transport or store under 32° F. If freezing occurs, thaw and agitate vigorously.

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUN 20 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

16-132

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL
Storage: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.
Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.
Container Disposal: Do not reuse empty container. Discard container in trash.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 16 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower leaf surfaces are

contacted. Use caution on tender foliage. Remove shipping cap before use. Sprayer should be rinsed thoroughly and stored empty for storage.

This insecticide may be used in greenhouses during the growing season or as a pre-treatment of other insecticides. Observe any special restrictions and

To control insects on lower leaf surfaces or as needed to control insects applied on day of

USE ON THE FOLLOWING: Beans, Broccoli, Brussel Sprouts, Cucumber, plant, Lettuce, Mustard, Turnips, Peppers, Potatoes, and Tomatoes.

TO CONTROL: Aphids, Leafhoppers, Whiteflies, Beetles, 12-spotted Leafhopper, Cabbage Loopers, Crossed-banded Leafhopper, Harlequin Bugs, Stink Bugs, Greenhouse Fleas, Fireworms, Blister Beetles and Webworms.

USE ON THE FOLLOWING: African Violets, Camellias, Dahlias, Dogwoods, Marigolds, Rose of Sharon, etc.

TO CONTROL: Aphids, Leafhoppers and Whiteflies.

4/4
To kill Gypsy Moth Caterpillars and adults, adult Japanese Beetles and Eastern and Forest Tent Caterpillars on contact.

ONTREES AND SHRUBS: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

ON ROSES AND IN FLOWER GARDENS: Treat plants and flowers as soon as insects appear. Spray plants and flowers, making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moths: apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure, filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at rate of 5 or 6 quarts per acre (4 oz. per 1,000 sq. ft.).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or .18 gallon per 1,000 sq. ft. Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

NOTICE:

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions for use and cautions stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

EPA Reg. No. 16-132

EPA Est. 16-VA-1