

PM-10

Reg # 16-132

194

OCT 29 1996

Ms. Rebecca M. Horton
Dragon Corporation
P.O. Box 7311
Roanoke, VA 24019-7311

Dear Ms. Horton:

Subject: Amendment- Label
Dragon Ready to Use Garden Pest Control
EPA Registration No. 16-132
Your Submission dated August 2, 1996

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. A stamped copy of the label is enclosed for your records.

Under "Environmental Hazards" replace the word "disposal" from the third sentence to read "disposing of equipment washwaters".

Sincerely yours,

RPK

Richard P. Keigwin, Jr.
Product Manager 10
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure:

274

**DRAGON®
READY TO USE GARDEN PEST CONTROL**

[ADDITIONAL BRAND NAMES: DRAGON® Ready-to-Use Vegetable Garden Insect Spray, DRAGON® Gentle Care™ House Plant Spray, Ready-to-Use DRAGON® Rose & Flower Insect Spray, DRAGON® Tomato & Vegetable Insect Spray, DRAGON® Rose & Flower Insect Spray]

Nothing Beats The Dragon!

Insect Protection For Delicate Plants

Fast Acting Organic Spray for Vegetables, Flowers and Ornamentals.

Kills: Aphids, Whiteflies, Beetles, Leafhoppers, Stink bugs, Caterpillars and many other listed insects.

Controls Insects on Ornamental and Delicate House Plants.

May be applied on day of harvest.

Apply with conventional mechanical compressed air equipment or non-aerosol pump.

Do not transport or store under 32°F. If freezing occurs, thaw and agitate vigorously.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., sites listed on label - roses, carnations, listed vegetables, etc.]

ACTIVE INGREDIENTS:

Pyrethrins	0.02%
*Piperonyl Butoxide, Technical	0.20%

INERT INGREDIENTS**:	99.78%
TOTAL	100.00%

*Equivalent to 0.16% (butylcarbityl)(6-propylpiperonyl) ether and 0.04% related compounds.

**Contains petroleum distillates.

**ACCEPTED
with COMMENTS
in EPA Letter Dated**

OCT 29 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
16-132

**Keep Out of Reach of Children
CAUTION**

See side/back/other panel(s) for additional precautions.

NET CONTENTS 8 FL OZ / 1 PT (16 FL OZ) / 1 QT (32 FL OZ)

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

CAUTION: Harmful if swallowed. **IF SWALLOWED:** Call a physician or Poison Control Center immediately.

ENVIRONMENTAL HAZARDS: This product is highly toxic to fish. Do not apply directly to water. Do not contaminate water when disposal of equipment washwaters. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage.

RECD EPA/OPP/DPD1

96 AUG -9 AM 1:38

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOMEOWNER USE ONLY. This insecticide may be used indoors, outdoors or in home greenhouses throughout the growing season, or as a preharvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestation. May be applied on day of harvest.

USE ON THE FOLLOWING CROPS:

Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes.

To Control (Kill): Asparagus beetles, Aphids, Leafhoppers, Whiteflies, Mexican bean beetles, 12-spotted cucumber beetles, Cabbage loopers, Cross-striped cabbageworms, Diamondback moth larvae, Flea beetles, harlequin bugs, Imported cabbageworms, Stink bugs, Green peach aphids, Leaf-tiers, Fireworms, Blister beetles, Colorado potato beetles, and Webworms.

USE ON THE FOLLOWING FLOWERS, ORNAMENTALS & HOUSE PLANTS:

African violets, Aralia, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Dracena, Ferns, Ficus, Geraniums, Gladioli, Marigolds, Palms, Roses, Rubber plants, Schefflera, Spathiphyllum, Wandering Jew, etc.

To Control (Kill): Aphids, Flea beetles, Leafhoppers and Whiteflies.

To Kill Gypsy Moth Caterpillars and adults, adult Japanese beetles and Eastern and Forest Tent Caterpillars on contact.

ON TREES AND SHRUBS: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

ON ROSES AND IN FLOWER GARDENS: Treat plants and flowers as soon as insects appear. Spray plants and flowers, making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moths: Apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure, filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

484

OUTDOOR, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at rate of 4 oz. per 1,000 sq. ft.

To control mosquito larvae, apply at the rate of 1 pint per 703 sq. ft. (approx. 20'x35' area). Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions for use and cautions stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019