

(Front panel)

FRAM  
FERBAM WETTABLE PULVER

Controls certain diseases of tomatoes, apples, pears, peaches, grapes, ornamentals and other plants.

ACTIVE INGREDIENT:  
Ferbam (Ferric dimethyl dithiocarbamate) ..... 76%  
INERT INGREDIENTS: ..... 24%

KEEP OUT OF REACH OF CHILDREN

CAUTION  
See back panel for additional precautionary statements

Net Weight 6 oz.

ACCEPTED  
*with label*  
SEP 9 1982  
Under the Federal Insecticide  
Fungicide, and Rodenticide Act  
as amended and authorized  
registered under  
EPA Reg. No. 16.134

(continued)

#### PREVENTION OF DAMAGE TO PLANTS

Apply to leaves, stems, and fruit.

Caution: May cause irritation of eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing.

#### STATEMENT OF HAZARDOUS REACTIVITY

In case of contact, flush with plenty of water; for eyes, seek medical attention.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specific on this label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

#### PHYSICAL OR CHEMICAL HAZARDS

Keep packages or dust away from sparks, flames or hot surfaces.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container; wrap and place in trash collection.

To mix, place the proper amount of this product into the spray tank. Add enough water to make the product into a thick paste by stirring. Bring to the proper dilution by continuing to stir as the additional quantity of water is added. THE DILUTION RATES GIVEN ARE LEVEL TEASPOONFULS TO BE ADDED TO ONE GALLON OF WATER. Wet plants to the dripping point. Observe days interval between last application and harvest indicated by number in ( ) following crop.

Apples (7) For early season control of Scab, Cedar Rust, Blotch use 1 - 1½ teaspoonfuls in prebloom & calyx stage spray. Use 1 - 1½ teaspoonfuls 10 days after petal fall and then space next two sprays 14 days apart. For Rust control, make an additional bloom application using ½ teaspoonful. For late season control of Scab, Black Rot, Sooty Blotch, Fly Speck and Brook's Spot use ¾ teaspoonful. Where Bitter Rot and Brook's Spot are severe problems, increase dosage to 1½ - 2 teaspoonfuls. On Golden Delicious, Jonathan, Baldwin and other varieties of apples which are especially apt to Russet. Some injury may result from the use of this product if used in the early season or bloom applications. The best time for use is after petal fall. When used in combination with sulfur, do not apply within 10 days to 2 weeks of an oil spray, nor just prior to or during periods of excessively high temperatures.

Cabbage (Plant Beds Only) Downy Mildew - use 1½ teaspoonfuls, making 2 or 3 applications each week.

(Back panel continued)

Grapes (7) Black Rot - Use 2 teaspoonfuls, applying just before bloom, just after bloom and two weeks later.

Ornamentals Black Spot and Rust on Roses; Rust on Begonias and Carnations; Leaf Spot on Chrysanthemums; Botrytis Blight on Peonies and Tulips - Use 1 - 1 1/2 teaspoonfuls, applying at 7 to 10 day intervals during growing season.

Peaches (21) Leaf Curl - Use 1 1/2 teaspoonfuls. Apply in the fall after leaves drop or in the spring before buds begin to swell.

Pears (7) Scab, Sooty Blotch - Use 1 1/2 teaspoonfuls in pink & calyx stages. Use 1 1/2 teaspoonfuls 10 days after petal fall and 14 days later. Spray at rate of 2 tablespoonfuls every 14 days until 7 days before harvest.

Tomatoes Gray Mold (Botrytis) - Use 3 teaspoonfuls. Spray for complete coverage. Make seed bed application just before seedlings break the soil surface and repeat at 7 day intervals. Make garden application when disease threatens and repeat at 7 to 10 day intervals as necessary to maintain control.

NOTICE: Buyer assumes all risk of use, handling or storage of this material not in strict accordance with label directions and precautions. This package is filled by weight, and not by volume. Product settling occurs after product is packaged.

EPA Reg. No. 16 - 134

EPA Est. 16 - VA - 1

Manufactured For

DRAGON CHEMICAL CORPORATION  
Roanoke, VA 24019