

FRONT PANEL

DRAGON

CAPTAN

WETTABLE
FUNGICIDE

Controls Listed diseases on roses,
flowers, fruits and vegetables.

MAY BE USED UP TO DAY OF HARVEST.

ACTIVE INGREDIENTS

Captan N- trichloromethyl. thio -4 cyclohexane-1,2-dicarboximide . . .	48.9%
Related Derivatives	1.1%
INERT INGREDIENTS	50.0%
TOTAL	100.0%


KEEP OUT OF REACH OF CHILDREN

DANGER

See back panel for additional cautions.

NET WEIGHT 8 OZ

ACCURATE
per self art.
 SEP 16 1987 *RFM*
 U.S. ...
 F.A.C. ...
 BSA ...
 16-135



BEST AVAILABLE COPY

BACK PANEL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS
& DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

In eyes - Flush eyes with water. Call physician immediately. If swallowed - Drink promptly a large quantity of milk, egg whites, gelatin solutions or if these materials are not available, drink large quantities of water. Avoid alcohol. Never give anything by mouth to an unconscious person. Call physician. If inhaled - Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention. If on skin - Wash contaminated skin with soap and water.

ENVIRONMENTAL HAZARDS


This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

COMPATIBILITY

This product is compatible with most commonly used insecticides and fungicides except highly alkaline materials such as hydrated lime. Captan should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixture.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE Store in the tightly closed original container in a cool, dry, preferably locked storage area out of reach of children. PESTICIDE DISPOSAL Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL Do not reuse empty container. Wrap and put in trash collection.



BACK PANEL Contd.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Wear chemical resistant gloves when using indoors or outdoors. Remove contaminated clothing and separately launder before reuse. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing.

Unless otherwise specified, use 3 1/3 tablespoonfuls of product mixed in each gallon of water. Use level measure. Mix designated amount in a little water to make a thin paste, add to 1 gallon water and agitated during application. Use a compression type or hose-end sprayer. Spray both upper and lower portions of leaves thoroughly.

ROSES: Black Spot Mix 3 tablespoonfuls in one gallon water at first sign of disease. TUBEROUS BEGONIAS: Powdery Mildew Mix 2 tablespoonfuls in 1 gallon water. Repeat at 7 to 10 day intervals. Apply at first sign of disease. CARNATIONS: Rust, Alternaria (Leaf Spot), CHRYSANTHEMUMS: Botrytis Flower Blight Apply at first sign of disease. Repeat every 7 to 14 days or as necessary. BULBS (Gladiolus, Tuberos Begonias): Rot, Damp-off- Preplanting Mix 8 tablespoonfuls in 1 gallon water. Dip bulbs in diluted mixture 20 or 30 minutes, drain and plant. Keep mixtures stirred.

TOMATOES: Early and Late Blight, Gray Leaf Spot, Anthracnose, Septoria Leaf Spot Apply at first sign of disease. Repeat 5 to 7 day intervals or as necessary to keep disease under control. Severe Blight Conditions - 5 1/3 tablespoonfuls per gallon water. Begin application 30 days after first bloom.

APPLES & PEARS: Scab, Fruit Spot, Bitter Rot Apply in preblossom, petal fall and cover applications. Do not use on D-Anjou pears.

PEACHES: Brown Rot, Peach Scab. CHERRIES: Cherry Leaf Spot, Brown Rot Apply at pink bud, full bloom, and petal fall stages. Repeat at 10 to 14 day intervals after petal fall.

STRAWBERRIES: Botrytis Rot. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe Infections Conditions Continue applications through harvest period, treating immediately after each picking.

NOTICE Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS FOR USE and PRECAUTIONARY STATEMENTS stated on this label.