

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95

United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number 229772
---	---	--

Application for Pesticide - Section I

1. Company/Product Number 16-151	2. EPA Product Manager J. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DRAGON Captan Wettable	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dragon Corporation P. O. Box 7311 Roanoke, VA 24019-7311 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION: Add "marketing copy" to label of record.

NOTIFICATION
201 / 1995

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product				<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Rebecca M. Horton	Title Regulatory & Sales Support Mgr.	Telephone No. (Include Area Code) (540) 362-3657
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Rebecca M. Horton</i>	3. Title Regulatory & Sales Support Manager	
4. Typed Name Rebecca M. Horton	5. Date 9/17/96	

AFTER 2 9 5

[Front panel]

**DRAGON®
CAPTAN WETTABLE**

Nothing Beats The Dragon!

Controls listed diseases on roses, flowers and fruits.

MAY BE USED UP TO DAY OF HARVEST.

Broad-Spectrum Fruit Disease Control
Use on Apples, Peaches, Grapes, Strawberries, and others listed

[Note to EPA: Label front panel may include graphics illustrating product use; e.g., Brown Rot, Powdery Mildew, Scab, Black Spot, etc.]

ACTIVE INGREDIENTS:

*Captan	48.9%
Related Derivatives	1.1%
INERT INGREDIENTS:	<u>50.00%</u>
TOTAL:	100.00%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

**Keep Out of Reach of Children
DANGER**

See back/side/other panels for additional precautionary statements.

NET WEIGHT 8 OZ

NOTIFICATION
JUL 4 1996

[BACK/SIDE/OTHER PANELS AND/OR BOOKLET]


DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

COMPATIBILITY: This product is compatible with most commonly used insecticides and fungicides, except highly alkaline materials such as hydrated lime. Captan should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixture.

DIRECTIONS: FOR HOMEOWNER USE ONLY. Wear goggles or face shield when handling. Wear chemical resistant gloves when using indoors or outdoors. Remove contaminated clothing and separately launder before reuse. When using outdoors, wear long pants and long-sleeved shirt, and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing.

Where foliage applications are directed, mix designated amount of material in a little water to make a thin paste, add to specified amount of water, and keep mixture agitated during application. Use a compression-type or hose-end sprayer.



FRUITS

Amount of Spray Needed Per Tree	
Size	Diluted Spray
Young or Dwarf	1 to 2 pints
Medium	1 to 2 quarts
Large	½ to 3 gallons

APPLES: Scab, Fruit Spot, Bitter Rot, Black Rot, Botrytis Blossom-end Rot, Sooty Blotch, Fly Speck, Black Pox, Botryosphaeria Rot - Use 2 1/5 level Tbs. per 1 gal. water. Apply in pre-blossom, petal fall and cover applications.

APRICOTS: Brown Rot (Twig Blight), Jacket Rot - Mix 1 1/3 level Tbs. in 1 gal. water. Apply in red bud, bloom, 75% petal fall and cover sprays.

BLUEBERRIES: Botrytis Gray Mold or Berry Rot, Mummy Berry - Mix 1 1/3 level Tbs. in 1 gal. water. Start spray program when buds swell or when buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom.

CHERRIES: Brown Rot, Leaf Spot, Botrytis Rot - Mix 1 1/10 level Tbs. in 1 gal. water and apply per previous tree chart. Apply at pink bud, full bloom and petal fall stages. Repeat at 10 to 14 day intervals after petal fall.

GRAPES: Black Rot, Downy Mildew - Mix 1 1/10 level Tbs. in 1 1/2 gal. water to cover an area of 400 sq. ft. (20'x20'). Make first application just before bloom, repeat just after bloom and at 10 to 14 day intervals for 1 to 3 additional applications, depending upon the susceptibility of the grape variety. Dead Arm (current season infection) - Mix 1 1/4 level Tbs. in 1 1/2 gal. water to cover an area of 400 sq. ft. (20'x20'). Apply first application when shoots are 1/2 to 1 inch long. Repeat application when shoots are 4 to 6 inches long.

NECTARINES: Brown Rot, Scab - Mix 2 1/2 level Tbs. in 1 1/2 gal. water. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications throughout harvest if conditions favor brown rot. Corvneum Blight (Peach Blight, Shothole) - Mix 2 1/2 level Tbs. in 2 1/2 gal. water. Apply in pink, bud, full bloom, petal fall and cover sprays as necessary, and as a post-harvest spray (but before leaves drop).

PEACHES: Brown Rot, Scab - Use 2 1/5 level Tbs. per 1 gal. water. Apply in full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays.

PLUMS, FRESH PRUNES: Brown Rot - Mix 1 2/3 level Tbs. in 2 Gal. water. Apply in full pink, bloom, and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage.

STRAWBERRIES: Botrytis Rot (Gray Mold), Leaf Spot - Mix 1 2/3 level Tbs. in 2 gal. water and apply to 400 sq. ft. area (20'x20'). Make first application when new growth starts in Spring before fruit starts to form. Repeat applications weekly. Severe infection conditions - Continue applications through harvest period, treating immediately after each picking.

FLOWERS & ORNAMENTALS

- AZALEAS:** Damping-off of Cuttings - Mix 1½ level Tbs. in 1 gal. water. Dip cuttings before bedding. Petal Blight - Mix ¾ level Tbs. in 1 gal. water. Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.
- BEGONIAS (Tuberous):** Powdery Mildew - Mix 1½ level Tbs. in 1 gal. water. Apply at first sign of disease. Repeat at 7 to 10 day intervals. Damping-off, Tuber Rot - Mix 1½ level Tbs. in 1 gal. water. Dip tubers in diluted mixture 30 minutes, drain and plant.
- CAMELLIAS:** Petal Blight - Mix ½ level Tbs. in 1 gal. water. Apply to drench soil around plants, beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.
- CARNATIONS:** Rust, Alternaria Leaf Spot - Mix ¾ level Tbs. in 1 gal. water. Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews. Damping-off of Cuttings - Mix 1½ level Tbs. in 1 gal. water. Dip cuttings before bedding.
- CHRYSANTHEMUMS:** Botrytis Flower Blight, Septoria Leaf Spot - Mix ¾ level Tbs. in 1 gal. water. Apply at first sign of disease. Repeat at 7 to 14 day intervals, or as necessary. Damping-off of Cuttings - Mix 1½ level Tbs. in 1 gal. water. Dip cuttings before bedding.
- GLADIOLUS (Corms):** Corm Rot and Decay, Damping-off - Mix 3 level Tbs. in 1 gal. water. Dip corms 25 minutes, drain and plant.
- GRASSES (Ornamental in non-pastured areas only):** Leaf Spot, Damping-off, Brown Patch, Melting Out, Seeding Blights, and Brown Spot on St. Augustine grass - Mix ¾ level Tbs. in 1 gal. water. Apply 1 gallon spray per 100 sq. ft. (10'x10'). Begin when growth starts in Spring. Repeat at 7 to 14 day intervals throughout season. Do not feed clippings to domestic animals nor allow them to feed on treated areas.
- ROSES:** Black Spot, Botrytis Blossom Blight - Mix ¾ level Tbs. in 1 gal. water. Apply at first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

STORAGE AND DISPOSAL

- Do not contaminate water, food or feed by storage or disposal.
- STORAGE:** Store in the tightly closed original container in a cool, dry, preferably locked storage area out of reach of children.
- PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.
- CONTAINER DISPOSAL:** Do not reuse empty container. Wrap and put in trash collection.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Flush with water. Call physician immediately.
- IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solutions or, if these materials are not available, drink large quantities of water. Avoid alcohol. Never give anything by mouth to an unconscious person. Call physician.
- IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

575

IF ON SKIN: Wash contaminated skin with soap and water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS FOR USE** and **PRECAUTIONARY STATEMENTS** stated on this label (or, in attached booklet).

THIS PACKAGE IS FILLED BY WEIGHT, NOT BY VOLUME.

Distributed by
DRAGON CORPORATION
Roanoke, VA 24019