

[FRONT PANEL]

**DRAGON®  
FRUIT TREE SPRAY WETTABLE**

Contains: Methoxychlor, Malathion, Captan.

A complete insecticide, fungicide and miticide for listed insect and disease control on apple trees.

[Note to EPA: Front panel of label may include graphics representing product use, e.g., insects/diseases controlled - Japanese beetle, Codling moth, Scab, Aphid, etc.]

**ACTIVE INGREDIENTS:**

- \*Methoxychlor [1,1,1-trichloro-2,2-bis(p-methoxyphenyl)ethane] ..... 10.00%
- Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide) ..... 6.85%
- Related derivatives ..... 0.15%
- Malathion (0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate) ..... 5.00%

**INERT INGREDIENTS:** ..... 78.00%

**TOTAL:** ..... 100.00%


\*Equivalent to 8.8% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.2% other isomers and related compounds.

**Keep Out of Reach of Children**

**DANGER**

See back/side/other panel(s) for additional precautionary statements.

NET WT. 1 LB / 4 LB



[BACK/SIDE/OTHER PANEL(S)]

**BEFORE USE READ CAUTIONS AND NOTICE STATEMENTS CAREFULLY**

**DIRECTIONS FOR USE**


It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Wear chemical resistant gloves when using indoors or outdoors. When using outdoors, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing. Remove contaminated clothing and separately launder clothing before reuse.

**SPRAY APPLICATION:** One pound of this material will make 10 gallons of fruit tree spray. For smaller quantities, use 5 level tsp. to each gallon of water. Mix well. Agitate solution while spraying.

AMOUNT OF SPRAY NEEDED PER TREE:	
SIZE	DILUTED SPRAY
Young or Dwarf	1 to 2 pints
Medium	1 to 2 quarts
Large	½ to 3 gallons

**APPLES:** To control scab, brooks fruit spot, woolly and green apple aphids, pacific clover and red spider mites, codling moth, curculio, apple maggot, Japanese beetle, and bitter rot apply as follows:



1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals to 21 days before harvest.

If control of disease on above fruit crops is necessary within 21 days of harvest, apply a wettable sulfur spray. Remove residues by washing, wiping, brushing or other effective means.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**DANGER:** Captan causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Use with adequate ventilation. Wash thoroughly with soap and water after handling.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Flush eyes with plenty of water. Call a physician.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solutions or, if these are not available, drink large quantities of water. Avoid alcohol. Never give anything by mouth to an unconscious person. Call a physician.

**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**IF ON SKIN:** Wash contaminated skin with soap and water.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

**BEE CAUTION:** This product is highly toxic to bees exposed to direct treatment or residues remaining on treated areas. Do not apply when bees are actively visiting the crop. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**STORAGE & DISPOSAL**

**STORAGE:** Do not store near feed or foodstuffs. Store in the tightly closed original container in a cool, dry preferably locked storage area, out of reach of children.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty container. Wrap and put in trash collection.

**NOTICE:** Buyer assumes all risks and liability of use, storage and/or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Manufactured by  
**DRAGON CORPORATION**  
 Roanoke, VA 24019

**BEST AVAILABLE COPY**

