



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:
16-159

Date of Issuance:
JUL 20 1993

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
Dragon All Purpose
Fungicide

Name and Address of Registrant (Include ZIP Code):

Dragon Corporation
Box 7311
Roanoke VA 24019

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

- a. Add the phrase, "EPA Reg. No. 16-159".
- b. In the Precautionary Statement Section reinstate "Skin sensitizer."
- c. The Storage and Disposal Section must appear in the Directions for Use not in the Precautionary Statements.

Signature of Approving Official:

151

Date:

JUL 20 1993

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

[FRONT PANEL]

JUL 20 1993

DRAGON®
All-Purpose Fungicide

CONTAINS DACONIL 2787®

NET CONTENTS 1 QT (32 OZ)/1 PT (16 OZ)/8 OZ

Under the Federal Insecticide
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

16-159

For control of diseases on Lawns, Shrubs, Trees, Fruits, Vegetables, Roses and flowers.

[Note to EPA: Label front panel may feature graphics representing diseases controlled, e.g., black spot, powdery mildew, rust.]

ACTIVE INGREDIENT:	BY WT.
Chlorothalonil (tetrachloroisophthalonitrile)*	13.9%
INERT INGREDIENTS:	87.1%
TOTAL	100.0%

*Contains 1.14 lbs. Chlorothalonil per gallon.

Keep Out of Reach of Children
CAUTION
See back/side/other panel(s) for additional precautions.

[BACK/SIDE/OTHER PANELS]

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FOR OUTDOOR DOMESTIC (HOMEOWNER) USE ONLY. SHAKE WELL BEFORE USING! Read entire label and use as directed. Apply only to plants specified on this label. Do not apply to plants not listed on this label. Do not combine this product with pesticides, surfactants or fertilizers unless prior use has shown the combination physically compatible and non-injurious under conditions of use.

NOTE TO USER: Do not graze treated areas. Do not feed clippings to livestock. This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Wear long-sleeve shirt, long pants and gloves when using this product.

DRAGON® All-Purpose Fungicide is highly effective in preventing a wide variety of commonly occurring diseases of flowering and foliage plants, shrubs, trees, bulbs, vegetables, lawns and ornamental turf. When disease is present **DRAGON® All-Purpose Fungicide** provides effective control when applied according to label directions.

GRASS & ORNAMENTAL LAWNS

To prevent or control the following listed diseases of grass and ornamental lawns, mix the recommended amount of **DRAGON® All-Purpose Fungicide** in one gallon of water. Apply as a thorough foliage spray to 200 sq. ft. (20'x10') of lawn area (see following chart). Make application when conditions favor disease development, such as wet or humid weather. Use the lowest application rate and longest interval for prevention or when disease conditions are light to moderate. When disease is severe, use the highest rate and shortest interval. Do not mow or water after treatment until lawn is thoroughly dry. Do not graze treated areas. Do not feed clippings to livestock. May cause slight burning or temporary discoloration to grass lawns if used when the temperature is above 90° F.

Disease	Spray Interval	Tbs./Gal. of Water to cover 200 sq. ft.	
		To Prevent Disease	To Control Existing Disease
Copper Spot (Ramulispora)	7-10 days	4-6 Tbs. per gal.	6-7 Tbs. per gal.
Curvularia (including blight, fading out, going out, leaf spot)	7-10 days	2-4 Tbs. per gal.	4-7 Tbs. per gal.
Dollar Spot (Sclerotinia)	7-14 days	2-4 Tbs. per gal.	4-7 Tbs. per gal.
Gray leaf spot (Piricularia)	7-10 days	2-4 Tbs. per gal.	4-7 Tbs. per gal.
Helminthosporium (including leaf spot, melting out)	7-10 days	2-4 Tbs. per gal.	4-7 Tbs. per gal.
Large Brown Patch (Rhizoctonia)	7-10 days	2-4 Tbs. per gal.	4-7 Tbs. per gal.
Red Thread (Corticium)	7-10 days	2-6 Tbs. per gal.	6-7 Tbs. per gal.
Stem Rust of Bluegrass (Puccinia)	7-14 days	4-6 Tbs. per gal.	6-7 Tbs. per gal.
DICHONDRA Alternaria Leaf Spot (California only)	7-14 days	4-6 Tbs. per gal.	6-7 Tbs. per gal.
Anthracnose	7-14 days	4-6 Tbs. per gal.	6-7 Tbs. per gal.

To control Gray Snow Mold (Typhula) in ornamental lawns: Mix 5-10 fl. oz. in 1 gallon of water to cover 200 sq. ft. Apply as a thorough foliage spray in autumn before winter snows. In areas of little snow cover, repeat at monthly intervals as needed for control.

ROSES, FLOWERS, BULBS

ORNAMENTAL PLANTS, WOODY SHRUBS, TREES & VINES

To prevent or control the specified diseases of plants listed below: Mix 4 Tbs. in 1 gallon of water (for Roses use 3 Tbs. in 1 gallon of water). Apply as a thorough foliage spray, taking care to completely wet upper and lower leaf surfaces. Coverage may vary depending on amount of foliage sprayed. Make first application as recommended for each plant and disease. Repeat at 7-14 day intervals as needed to maintain control. Use the longest interval for prevention or when disease conditions are light to moderate. When disease is severe, use the highest rate. **USE PRECAUTION:** Do not apply to tender foliage during hot, sunny weather. Do not apply to wilted plants.

ASH: Leaf spot (Cercospora), Leaf spot (Cercosporidium, Cylindrosporium) — Apply at spring bud break.

AZALEA*, RHODODENDRON*: Ovulinia petal blight — Apply at early bloom. Phytophthora dieback — Apply when new leaves emerge.

BUCKEYE, HORSECHESTNUT: Anthracnose (Glomerella), Leaf blotch (Guidnardia) — Apply at spring bud break.

CARNATION: Alternaria leaf spot and stem rot — Apply when transplanting cuttings. Botrytis blossom blight — Apply when weather is cool and moist or when transplanting cuttings.

CHERRY-LAUREL: Cercospora leaf spot — Apply at petal fall.

CHRYSANTHEMUM, DAISIES: Gray mold blight (Botrytis) — Apply before bloom. Leaf spot (Septoria), Ray blight (Mycosphaerella) — Apply when transplanting cuttings or before bloom.

DRACAENA: Leaf spot (Fusarium) — Apply prior to transplanting.

EUNONYMUS: Anthracnose (Colletotrichum) — Apply at spring bud break.

FIRETHORN: Scab (Fusicladium) — Apply at spring bud break.

) FLOWERING ALMOND, FLOWERING QUINCE, PURPLELEAF SAND CHERRY: Brown rot blossom and twig blight (Monilinia) — Apply at early bloom.

FLOWERING CRABAPPLE: Cedar-apple rust (Gymnosporangium), Leaf spot (Sphaeropsis), Scab (Venturia) — Apply at spring bud break.

FLOWERING DOGWOOD: Leaf spot (Septoria) — Apply at early bloom.

GERANIUM: Gray mold blight (Botrytis), Rust - - Apply when weather is cool and moist.

GLADIOLUS: Botrytis leaf and flower spot, Curvularia leaf and flower spot -- Apply when leaves first emerge.

HAWTHORN: Leaf blight (Fabraea), Rust (Gymnosporangium) — Apply before bloom.

HOLLY: Rhizoctonia web blight — Apply during warm, moist conditions.

) HOLLYHOCK: Rust — Apply to young seedlings.

HYDRANGEA*: Leaf spot (Cercospora, Septoria), Rust (Pucciniastrum) — Apply to foliage only of young plants.

IRIS: Botrytis blossom blight, Leaf spot (Didymellina) — Apply during cool, moist conditions or before bloom.

LEATHERLEAF FERN: Ascochyta blight, Leaf spot (Cercospora), Leaf spot (Cylindrocladium), Rhizoctonia blight — Apply at spring bud break.

LIGUSTRUM (PRIVET): Leaf spot (Cercospora) — Apply during prolonged wet conditions.

LILIES: Gray mold blight (Botrytis) — Apply before bloom.

MOUNTAIN-LAUREL: Leaf spot (Cercospora) — Apply at spring bud break.

6049

OAK (RED GROUP ONLY): Taphrina blister, Leaf spot (Actinopelte), Anthracnose — Apply at dormant budswell.

OREGON GRAPE (MAHONIA): Rust — Apply at spring bud break.

OYSTER PLANT (RHOEO) also known as MOSES-IN-THE-CRADLE: Tan leaf spot (Curvularia) — Apply to young plants.

PARLOR PALM (CHAMAEDOREA): Bipolaris leaf spot (Helminthosporium) — Apply during cool, moist conditions.

PETUNIA*: Crown rot (Phytophthora), Gray mold and blight (Botrytis) — Apply before bloom.

PHILODENDRON: Leaf spot (Dactylaria), Phytophthora blight — Apply during moist conditions.

PIERIS (ANDROMEDA): Phytophthora die-back — Apply when new leaves emerge.

PHOTINIA: Fabraea (Entomosporium) leaf spot — Apply at spring bud break.

POPLAR: Marssonina leaf spot — Apply at spring bud break.

PRAYER PLANT (MARANTA): Helminthosporium leaf spot — Apply to young plants.

ROSES (Use 3 Tbs. in 1 Gal. of water): Black spot (Diplocarpon), Botrytis blossom blight, Suppression of Powdery mildew (Sphaerotheca) — Apply at spring bud break.

STATICE: Anthracnose, Leaf spots (Cercospora, Alternaria), Leaf blight (Botrytis) — Apply at spring bud break.

SYCAMORE, PLANETREE: Anthracnose — Apply at spring bud break.

SYNGONIUM: Cephalosporium leaf spot — Apply during warm, moist conditions.

VIBURNUM: Powdery mildew (Microsphaera) — Apply mid-summer.

ZINNIA: Powdery mildew (Erysiphe) — Apply at first sign of disease.

*Discoloration has been noticed on certain varieties when application is made during bloom.

ORNAMENTAL CONIFERS: To control Rhabdocline needle cast and Swiss needle cast of Douglas Fir, Sirococcus tip blight, and Scleroderris canker of Pine: Mix 3 Tbs. in 1 Gal. of water. To control Rhizosphaera needle cast of spruce and Scirrhia brown spot of Pine: Mix 6 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray in spring when new shoot growth is 1/4 to 2 inches long. Repeat at 3 to 4 week intervals until conditions no longer favor disease development (wet or humid weather).

FRUITS & VEGETABLES

To prevent or control listed diseases in the home garden, apply DRAGON® All-Purpose Fungicide when conditions favor disease development, such as wet or humid weather. Use the lowest rate and longest interval of application when disease conditions are light to moderate. For severe disease conditions, apply the highest rate at the shortest interval. Numbers in parentheses () next to each crop indicate days required between last application and harvest. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area. USE PRECAUTION: Do not apply to any crop not listed on this label. Do not graze treated areas or feed treated plants to livestock.

7097

APRICOT, NECTARINE, PEACH: Brown rot blossom and twig blight (*Monilinia*), Coryneum blight (shothole), Leaf curl—nectarine and peach—(*Taphrina*), Scab (*Cladosporium*) — Mix 1 Tbs. in 1 Gal. of water. For leaf curl, make first application in late autumn or early winter before first freeze. Where shothole occurs, apply once at petal fall or at shuck split. For brown rot blossom and twig blight, spray at popcorn stage and again at full bloom. If conditions favor disease development, make an additional application at petal fall. For scab, apply at popcorn stage, full bloom and shuck split.

BEANS, DRY (blackeye, kidney, lima, navy, pinto) (42): Anthracnose (*Colletotrichum*), *Cercospora* leaf spot—blackeye only, Downy mildew (*Peronospora*), and Rust (*Uromyces*) — Mix 2-3 Tbs. in 1 Gal. of water and apply as a thorough foliage spray at early bloom. Repeat at 7-10 day intervals as needed for adequate control.

BEANS, SNAP (7): Rust (*Uromyces*) — Mix 2-3 Tbs. in 1 Gal. of water. Gray mold blight (*Botrytis*) — Mix 4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray during early bloom or when disease first threatens. Repeat at weekly intervals or as needed to maintain control.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER (all 0): Downy mildew (*Peronospora*), Gray leaf spot (*Alternaria*), Ringspot of Brussels sprouts—California only (*Mycosphaerella*) — Apply as a thorough foliage spray after transplant or when seedlings emerge and when conditions favor disease development. Mix 2-3 Tbs. in 1 Gal. of water. Repeat at 7-10 day intervals or as needed to maintain control.

CARROTS (0): Early blight (*Cercospora*), Late blight (*Alternaria*). Mix 2-3 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray when disease threatens. Repeat at 7-10 day intervals or as needed to maintain control.

CELERY (7): Basal stalk rot (*Rhizoctonia*), Early blight (*Cercospora*), Late blight (*Septoria*), Pine rot suppression (*Sclerotinia*). Mix 3-4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray at transplant. Repeat at 7 day intervals or as needed to maintain control.

CHERRY (Sweet and Sour): Brown rot blossom and twig blight (*Monilinia*), Cherry leaf spot (*Coccomyces*) — Mix 1 Tbs. in 1 Gal. of water. Make first application at delayed dormant - the period just before blossom buds break open - and a second at full bloom. If conditions favor disease development, make an additional application at petal fall. Do not apply after petal fall. ✓

CORN, SWEET (14): *Helminthosporium* leaf blight, Rust — Mix 1½ to 3 Tbs. in 1 Gal. of water. Apply at first sign of disease. Repeat at 4-7 day intervals or as needed to maintain control. Do not apply to sweet corn to be processed.

CUCUMBER (0): Anthracnose (*Colletotrichum*), Downy mildew (*Pseudoperonospora*), Gummy stem blight (*Mycosphaerella*), Leaf blight (*Alternaria*), Powdery mildew—except Southwestern states—(*Erysiphe*), Scab (*Cladosporium*), Target leaf spot (*Corynespora*) — Mix 2½ to 4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray when first leaves develop or when disease threatens. Repeat at 7 day intervals. Shorter intervals may be used when disease is severe.

GARLIC (7); **LEAK** (14); **SHALLOT** (14): *Botrytis* leaf blast, Downy mildew suppression (*peronospora*), Purple blotch (*alternaria*) — Mix 2 to 4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray at first sign of disease or during rainy periods. Repeat at 7-10 day intervals as needed to maintain control. Do not apply more than 3 times per season.

PUMPKIN, SQUASH (Summer and Winter), **WATERMELONS, MELONS** (Cantaloupe, Honeydew, Muskmelon) (all 0): Anthracnose (*Colletotrichum*), *Cercospora* leaf spot, Downy mildew (*Pseudoperonospora*), Gummy stem blight (*Mycosphaerella*), Leaf blight (*Alternaria*), Powdery mildew—except Southwestern states—(*Erysiphe*), Scab (*Cladosporium*) — Mix 3 to 4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray when conditions favor disease development, such as wet or humid weather. Repeat at 7 day intervals. Shorter intervals may be used during severe disease conditions.

7979

ONION, DRY (7): Botrytis leaf blast, Purple blotch (Alternaria) — Mix 2 to 3 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray at first sign of disease or during rainy periods. Repeat at 7-10 day intervals for as long as conditions favor disease development. Use the shorter interval during persistent rainy periods.

ONION, GREEN (14): Botrytis leaf blast, Downy mildew suppression (Peronospora), Purple blotch (Alternaria) — Mix 2 to 4 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray at first sign of disease or during rainy periods. Repeat at 7-10 day intervals for as long as conditions favor disease development. Use the shorter interval during persistent rainy periods.

PAPAYA (0): Anthracnose (Colletotrichum), Fruit spot (Alternaria), Stem-end rot (Phytophthora) — Mix 3 to 5 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray when conditions favor disease development, such as wet or humid weather. Repeat at 14 day intervals to maintain control.

PARSNIP (10): Alternaria leaf spot, Anthracnose (Cylindrosporium), Botrytis blight (gray mold), Bottom rot (Rhizoctonia), Downy mildew (Plasmopara) — Mix 2 to 3 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray at first sign of disease or when conditions favor disease development, such as wet or humid weather. Repeat at 7-10 day intervals. Do not apply more than 4 times per season.

PASSION FRUIT—Hawaii only (0): Alternaria fruit spot. Mix 3 Tbs. in 1 Gal. of water. Apply as a thorough foliage spray when fruit spots appear (April to July). Full coverage of fruit and leaves is important to maintain control. Repeat at 14 day intervals until weather conditions no longer favor disease development.

PLUM, PRUNE: Brown rot blossom blight (Monilinia) — Mix 1 Tbs. in 1 Gal. of water. Make first application at delayed dormant—the period just before blossom buds break open—and a second at full bloom. If conditions favor disease development, make an additional application at petal fall. Do not apply after petal fall.

POTATO (0): Botrytis vine rot, Early blight (Alternaria), Late blight (phytophthora) — Mix 1½ to 2 Tbs. in 1 Gal. of water. Apply when plants are 6-8 inches high or when disease threatens. Repeat at 7-10 day intervals or as needed to maintain control.

TOMATOES (0): Anthracnose (colletotrichum), Black mold rot (Alternaria), Early blight (Alternaria), Fruit rot (Rhizoctonia), Gray leaf spot (Stemphylium), Gray mold blight (Botrytis), Late blight (Phytophthora), Leaf mold (Cladosporium), Septoria leaf spot — Mix 2-4 Tbs. in 1 Gal of water. Apply as a thorough foliage spray when disease threatens. Repeat at 7-10 day intervals. During conditions of severe disease pressure, use the shortest interval.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

skin sensitizes

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or inhaled. Avoid contact with eye, skin or clothing. Avoid inhaling vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water, call physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

220

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

STORAGE & DISPOSAL

STORAGE: Store only in original container, in a cool dry place, inaccessible to children and pets, and away from foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly and discard in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with label directions and cautions.

(Daconil 2787 covered under U.S. Pat. No. 3,290,353 and No. 3,331,735.)

Distributed by
DRAGON CORPORATION
Roanoke, VA 24019

✓
.....
.....