



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number: 16-172
Date of Issuance: JAN 22 1997

Term of Issuance: Until Reregistration

Name of Pesticide Product: Dragon Dursban 1% Granular Insecticide

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Dragon Corporation
P.O. Box 7311
Roanoke, Virginia 24019

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 16-172."
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Date:

JAN 22 1997

2712

[FRONT PANEL]

DRAGON®
DURSBAN® 1% GRANULAR INSECTICIDE

Nothing Beats the Dragon!

Controls [Kills]: Ants, Fleas, Brown Dog Ticks, Crickets, Chinch Bugs, Certain White Grubs, Sod Webworms (Lawn Moths), Cutworms, Earwigs, Sowbugs (Pillbugs), and Hyperodes (Turfgrass) Weevils.

Controls [Kills, Stops] Lawn, Garden & Home Invading Pests.

Ready To Use Granules.

Controls [Kills] Imported Fire Ants. No Watering Necessary. For Spot Treatment of Imported Fire Ant Mounds.

For Outdoor Use Only.

Defend Yourself Against Home Invading Pests!

[10 lb. package] Treats Up to 4,000 Sq. Ft.

Stops Insects from Invading the Home.

Controls [Kills] vegetable soil insects such as: Root Maggots, Onion Maggots and Cutworms.

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 22 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

16-172

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)phosphorothioate]	1.0%
INERT INGREDIENTS:	<u>99.0%</u>
TOTAL:	100.0%

[NOTE TO EPA: Front panel of label may include graphics representing product use; e.g., insects controlled]

Keep Out of Reach of Children
CAUTION

See back/side/other panel(s) for additional precautionary statements.

NET WEIGHT ____ LB

96 OCT 16 AM 0:44

RECD EPA/C/P/DPD1

[BACK/SIDE/OTHER PANEL(S)]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Do not take internally. Avoid inhaling dust. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and domestic water supplies.

3/12

FIRST AID

- IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED:** Remove victim to fresh air if symptoms of cholinesterase inhibition appear, and get medical attention immediately.
- IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.
- IF ON SKIN:** Wash thoroughly with soap and water.
- NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, by injection only, is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

VERSION A

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To kill listed ants, armyworms, billbugs, boxelder bugs, chiggers, chinch bugs, clover mites, corn borers, crickets, cutworms, earwigs, European crane fly larvae, fire ants, fleas, flea beetle larvae, grasshoppers, Hyperodes (turfgrass) weevils, lesser cornstalk borers, root maggots, rootworm larvae, seed corn beetles, sod webworms (lawn moths), sowbugs (pillbugs), spiders, sweet potato beetles, symphylans, ticks (including deer ticks), white grubs, wireworms.

WHERE TO USE: Use on home lawns and other ornamental turf areas, in flower beds, under shrubbery, around buildings and ~~in home vegetable gardens.~~ Do not use this product indoors.

HOW TO USE:

LAWNS

To control ants, boxelder bugs, chiggers, chinch bugs, clover mites, crickets, earwigs, European crane fly larvae*, fire ants*, fleas, grasshoppers, Hyperodes (turfgrass) weevils*, sod webworms (lawn moths), sowbugs (pillbugs), spiders, ticks (including deer ticks)*, white grubs*.
(*See the following specialized application directions.)

Treat when insects first appear. Immediately after application, give the lawn a moderate watering in order to move the insecticide down into the soil.

Apply **DRAGON® Dursban® 1% Granular Insecticide** uniformly to home lawns and other ornamental and recreational turf areas using any hand or rotary-type spreader designed for application of granular materials. Treat when insects first appear using granules at the rate of 2½ lbs. per 1,000 sq. ft. Consult the Application Rate Table to determine the amount of product needed for larger-sized areas. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle.

BEST AVAILABLE COPY

4912

Application Rates	
Area to be Treated	Lbs. of Granules
1,000 sq. ft.	2½ lbs.
2,000 sq. ft.	5 lbs.
4,000 sq. ft.	10 lbs.
5,000 sq. ft.	12½ lbs.
10,000 sq. ft.	25 lbs.

SUGGESTED SPREADER SETTINGS Spreader Type	Delivery Rate in Pounds per 1,000 Sq. Ft.	
	2.5	5
Ortho Broadcast Spreader	4	7½
Ortho Drop Spreader (Model 5000)	6	7½
Scotts Drop Spreader	6	7½
Central Drop Spreader	6¾	—
Scotts Easy Green	27 ½	30
Sears Craftsman	6 ½	8 ½
Republic Broadcast	4	7½
Republic Drop Spreader	6	7½
Hand Held Spreaders		
ORTHO Whirlybird Spdr	4	5
Republic	4	5

Note: Above spreader settings are approximate. Accurately weigh several pounds into your spreader and apply to the recommended area. Adjust spreader setting if necessary to assure the correct rate of application.

2 cups = approximately 1 lb. of granules.

1. **European Crane Fly:** Apply at the rate of 3½ lbs. per 1,000 sq. ft. For increased effectiveness, follow the granule application with a thorough watering to move the insecticide into the soil. Apply the equivalent to ½-1 inch of water by using a sprinkler or other means of applying water to turf. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.
2. **Fire Ants (for individual ant mounds):** Apply at a rate of 2 oz. (4 Tbs.) per active mound. Sprinkle granules gently and evenly over surface of mound, and out approximately 2 feet surrounding the mound. Avoid disturbing ants. For best results, apply when ants are least active; in cool weather, 65°F to 80°F, or in early morning or late evening hours. Treat new mounds as they appear.
3. **Turfgrass weevils (larvae of Hyperodes weevil):** Apply at the rate of 4½ lbs. per 1,000 sq. ft. Apply the granules uniformly over the area to be treated. For best results, treat when grubs are young and actively feeding near the soil surface. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.
4. **Ticks:** Treat perimeter of house and yard by applying under shrubbery, in and around ground cover and around and under patios. Treat edges or perimeter of yard area where grass and other ground cover plants exist and other areas that appear likely to serve as harborage sites for ticks that have removed themselves from their host. Prior to and following treatment, weeds and high grasses should be kept trimmed to reduce tick harborage sites. Rodents can serve as hosts for ticks. Rock piles, wood piles and other rodent habitats should be removed or reduced. Key months for tick control are April through November in most parts of the country. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.
5. **White Grubs (larvae of Japanese beetle, black turfgrass atenius, European chafer and southern masked chafer):** Apply at the rate of 4½ lbs. per 1,000 sq. ft. When high populations of grubs are present in the area to be treated, a repeat application of 4½ lbs. per 1,000 sq. ft. may be made 7-14 days following the initial application. Apply the granules uniformly over the treated area. For increased effectiveness, follow the granule application with a thorough watering to move the insecticide into the soil. Apply the equivalent of ½-1 inch of water by using a sprinkler or other means of applying water to turf. For best results, treat when grubs are young and actively feeding near the soil surface. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.

PERIMETER TREATMENT TO HELP PREVENT INFESTATION OF BUILDINGS

To control ants, crickets, earwigs, fleas, sowbugs (pillbugs), clover mites, ticks, boxelder bugs, spiders, treat a band of soil 6-10 feet wide around and adjacent to buildings where pests are active and may find entrance. Use at the rate of 2½ lbs. per 1,000 sq. ft.

IN FLOWER GARDENS, UNDER AND AROUND SHRUBBERY

To control ants, crickets, cutworms, sowbugs (pillbugs), grasshoppers, earwigs, armyworms, ticks. Apply at a rate of 2½ lbs. per 1,000 sq. ft. evenly over soil within flower beds and under shrubbery.

VEGETABLE GARDENS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CHINESE CABBAGE, COLLARDS, KALE, KOHLRABI, RUTABAGAS, TURNIPS: To control root maggot and other pests, apply 7½-14 oz. (1-1¼ cup) per 100 feet of row by placing granules in a 4 inch band over the row after the seeds have been placed in the furrow and covered. Lightly cultivate the granules into the top ¼-½ inch of soil. Make only 1 application per year at planting time. Do not use rutabaga tops for food or feed purposes.

ONIONS: To control onion maggot apply 5½ oz. (11 Tbs.) per 100 ft. of row by placing granules in the furrow at planting time. Cover with soil. Make only 1 application per year at planting time.

PEANUTS: To control cutworms, wireworms, lesser cornstalk borer, southern corn rootworm larvae. *At-Plant Application:* Apply 11-22½ oz. (1½-2¾ cups) per 100 ft. of row by placing granules in

a 6-12 inch band over the row after the seeds have been placed in the planting furrow and covered. Lightly cultivate the granules into the surface of the soil. *Post-Plant Application:* Apply 11-22½ oz. (1½-2¾ cups) per 100 feet of row at early flowering stage in a 6-12 inch band over the row; or during early pegging in a 10-12 inch band over the fruiting zone. Use the 11 oz. rate only if the 11 oz. rate was applied at planting. Irrigation or rain following application will enhance effectiveness. Do not apply more than a total of 22½ oz. (2¾ cups) per 100 ft. of row per season. Do not harvest within 21 days of last treatment. Do not feed above ground portions of peanut plants to domestic animals.

RADISHES: To control root maggot apply 5 oz. (½ cup) per 100 ft. of row by placing granules in the furrow at planting time. Cover granules and seeds with suitable soil. Make only 1 application per year at planting time.

SWEET CORN: To control wireworms, seed corn maggots, root maggots, seed corn beetle, symphylans, cutworms, fire ants, billbugs, northern and western corn rootworm larvae, southern corn rootworm larvae, lesser cornstalk borer, flea beetle larvae, European and southwestern corn borer, chinch bug, grubs apply 12½-19 oz. (1½-2½ cups) per 100 feet of row. With seed still uncovered in furrow, apply granules in the furrow and along the sides of the furrow for a total of a 10 inch wide band. Cover seed with treated soil. After seed is covered, any remaining surface granules should be lightly cultivated into the top ½-1 inch of soil. In soils containing over 10% organic matter, use the highest rate given (19 oz. (2½ cups)/100 ft. of row). Make only 1 application per year at planting time.

SWEET POTATOES: To control wireworms, flea beetles, sweet potato beetle. *Pre-plant Broadcast Application:* Evenly broadcast 4½ lbs. (9 cups) per 1,000 sq. ft. of planting area over the soil surface and then cultivate the soil down to a depth of 4-6 inches. Plant the sweet potatoes in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time the granules will protect against feeding damage by the listed pests). This product will not control false wireworm of whitefringed beetle and certain other grubs that attack sweet potatoes. Make only one application per year prior to planting. Do not harvest within 125 days of treatment.

SUNFLOWERS: To control cutworms, apply 12½ oz. (1½ cup) per 100 ft. of row by placing granules in a 7 inch band over the row after the seeds have been placed in the furrow and covered. Lightly cultivate the granules into the top 1 inch of soil.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

CONTAINER IS FILLED BY WEIGHT, NOT BY VOLUME. Some settling of contents may occur.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.

DURSBAN® is a registered trademark of DowElanco.

Distributed by
DRAGON CORPORATION
Roanoke, VA 24019

BEST AVAILABLE COPY

VERSION B

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To kill listed ants, armyworms, billbugs, boxelder bugs, chiggers, chinch bugs, clover mites, corn borers, crickets, cutworms, earwigs, European crane fly larvae, fire ants, fleas, flea beetle larvae, grasshoppers, Hyperodes (turfgrass) weevils, lesser cornstalk borers, root maggots, rootworm larvae, seed corn beetles, sod webworms (lawn moths), sowbugs (pillbugs), spiders, sweet potato beetles, symphylans, ticks (including deer ticks), white grubs, wireworms.

WHERE TO USE: Use on home lawns and other ornamental turf areas, in flower beds, under shrubbery, around buildings and ~~in home vegetable gardens.~~ Do not use this product indoors.
in home vegetable gardens.

HOW TO USE:

LAWNS	
Apply DRAGON® Dursban® 1% Granular Insecticide uniformly to home lawns and other ornamental and recreational turf areas using any hand or rotary-type spreader designed for application of granular materials. Treat when insects first appear. Immediately after application, give the lawn a moderate watering in order to move the insecticide down into the soil. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle.	
To Control:	Amt. to Use:
Ants, Boxelder bugs, Chiggers, Chinch bugs, Clover mites, Crickets, Earwigs, European crane fly larvae*, Fire ants*, Fleas, Grasshoppers, Hyperodes (turfgrass) weevils*, Sod webworms (lawn moths), Sowbugs (pillbugs), Spiders, Ticks (including deer ticks)*, White grubs*	2½ lbs. per 1,000 sq. ft. - or - 5 lbs. per 2,000 sq. ft. - or - 10 lbs. per 4,000 sq. ft. - or - 12½ lbs. per 5,000 sq. ft. - or - 25 lbs. per 10,000 sq. ft.
(*See the following specialized application directions.)	

BEST AVAILABLE COPY

SUGGESTED SPREADER SETTINGS Spreader Type	Delivery Rate in Pounds per 1,000 Sq. Ft.	
	2.5	5
Ortho Broadcast Spreader	4	7½
Ortho Drop Spreader (Model 5000)	6	7½
Scotts Drop Spreader	6	7½
Central Drop Spreader	6¾	—
Scotts Easy Green	27 ½	30
Sears Craftsman	6 ½	8 ½
Republic Broadcast	4	7½
Republic Drop Spreader	6	7½
Hand Held Spreaders		
ORTHO Whirlybird Spdr	4	5
Republic	4	5
<p>Note: Above spreader settings are approximate. Accurately weigh several pounds into your spreader and apply to the recommended area. Adjust spreader setting if necessary to assure the correct rate of application. 2 cups = approximately 1 lb. of granules.</p>		

- 1. European Crane Fly:** Apply at the rate of 3½ lbs. per 1,000 sq. ft. For increased effectiveness, follow the granule application with a thorough watering to move the insecticide into the soil. Apply the equivalent to ½-1 inch of water by using a sprinkler or other means of applying water to turf. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.
- 2. Fire Ants (for individual ant mounds):** Apply at a rate of 2 oz. (4 Tbs.) per active mound. Sprinkle granules gently and evenly over surface of mound, and out approximately 2 feet surrounding the mound. Avoid disturbing ants. For best results, apply when ants are least active; in cool weather, 65°F to 80°F, or in early morning or late evening hours. Treat new mounds as they appear.
- 3. Turfgrass weevils (larvae of Hyperodes weevil):** Apply at the rate of 4½ lbs. per 1,000 sq. ft. Apply the granules uniformly over the area to be treated. For best results, treat when grubs are young and actively feeding near the soil surface. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.
- 4. Ticks:** Treat perimeter of house and yard by applying under shrubbery, in and around ground cover and around and under patios. Treat edges or perimeter of yard area where grass and other ground cover plants exist and other areas that appear likely to serve as harborage sites for ticks that have removed themselves from their host. Prior to and following treatment, weeds and high grasses should be kept trimmed to reduce tick harborage sites. Rodents can serve as hosts for ticks. Rock piles, wood piles and other rodent

BEST AVAILABLE COPY

habitats should be removed or reduced. Key months for tick control are April through November in most parts of the country. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.

- 5. **White Grubs** (larvae of Japanese beetle, black turfgrass ataenius, European chafer and southern masked chafer): Apply at the rate of 4½ lbs. per 1,000 sq. ft. When high populations of grubs are present in the area to be treated, a repeat application of 4½ lbs. per 1,000 sq. ft. may be made 7-14 days following the initial application. Apply the granules uniformly over the treated area. For increased effectiveness, follow the granule application with a thorough watering to move the insecticide into the soil. Apply the equivalent of ½-1 inch of water by using a sprinkler or other means of applying water to turf. For best results, treat when grubs are young and actively feeding near the soil surface. Consult your local Agricultural Extension Service specialist for proper timing to treat in your area.

PERIMETER TREATMENT TO HELP PREVENT INFESTATION OF BUILDINGS	
Treat a band of soil 6-10 feet wide around and adjacent to buildings where pests are active and may find entrance.	
To Control:	Amt. to Use:
Ants, Boxelder bugs, Clover mites, Crickets, Earwigs, Fleas, Sowbugs (pillbugs), Spiders, Ticks	2½ lbs. per 1,000 sq. ft.

IN FLOWER GARDENS, UNDER AND AROUND SHRUBBERY	
Distribute granules evenly over soil within flower beds and under shrubbery.	
To Control:	Amt. to Use:
Ants, Armyworms, Crickets, Cutworms, Earwigs, Grasshoppers, Sowbugs (pillbugs), Ticks	2½ lbs. per 1,000 sq. ft.

VEGETABLE GARDENS			
Use on:	To Control:	Amt. to Use:	Directions:
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CHINESE CABBAGE, COLLARDS, KALE, KOHLRABI, RUTABAGAS, TURNIPS	root maggot and other pests	7½-14 oz. (1-1¼ cup) per 100 feet of row	Place granules in a 4 inch band over the row after the seeds have been placed in the furrow and covered. Lightly cultivate the granules into the top ¼-½ inch of soil. Make only 1 application per year at planting time. Do not use rutabaga tops for food or feed purposes.

BEST AVAILABLE COPY

VEGETABLE GARDENS			
Use on:	To Control:	Amt. to Use:	Directions:
RADISHES		5 oz. (½ cup) per 100 ft. of row	Place granules in the furrow at planting time. Make only 1 application per year at planting time.
ONIONS	onion maggot	5½ oz. (11 Tbs.) per 100 ft. of row	Place granules in the furrow at planting time. Make only 1 application per yer at planting time.
PEANUTS	cutworms, wireworms, lesser cornstalk borer, southern corn rootworm larvae	11-22½ oz. (1½-2¾ cups) per 100 ft. of row	<i>At-Plant Application:</i> Place granules in a 6-12 inch band over the row after the seeds have been placed in the planting furrow and covered. Lightly cultivate the granules into the surface of the soil. <i>Post-Plant Application:</i> Apply at early flowering stage in a 6-12 inch band over the row; or during early pegging in a 10-12 inch band over the fruiting zone. Use the 11 oz. rate only if the 11 oz. rate was applied at planting. Irrigation or rain following application will enhance effectiveness. Do not apply more than a total of 22½ oz. (2¾ cups) per 100 ft. of row per season. Do not harvest within 21 days of last treatment. Do not feed above ground portions of peanut plants to domestic animals.

BEST AVAILABLE COPY

VEGETABLE GARDENS			
Use on:	To Control:	Amt. to Use:	Directions:
SWEET CORN	wireworms, seed corn maggots, root maggots, seed corn beetle, symphylans, cutworms, fire ants, billbugs, northern and western corn rootworm larvae, southern corn rootworm larvae, lesser cornstalk borer, flea beetle larvae, European and southwestern corn borer, chinch bug, grubs	12½-19 oz. (1½-2½ cups) per 100 feet of row	With seed still uncovered in furrow, apply granules in the furrow and along the sides of the furrow for a total of a 10 inch wide band. Cover seed with treated soil. After seed is covered, any remaining surface granules should be lightly cultivated into the top ½-1 inch of soil. In soils containing over 10% organic matter, use the highest rate given (19 oz.(2½cups)/100 ft. of row). Make only 1 application per year at planting time.
SWEET POTATOES	wireworms, flea beetles, sweet potato beetle	4½ lbs. (9 cups) per 1,000 sq. ft.	<i>Broadcast Application:</i> Evenly broadcast over the soil surface and then cultivate the soil down to a depth of 4-6 inches. Plant the sweet potatoes in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time the granules will protect against feeding damage by the listed pests). This product will not control false wireworm of whitefringed beetle and certain other grubs that attack sweet potatoes. Make only one application per year prior to planting. Do not harvest within 125 days of treatment.

SUNFLOWERS	
Place granules in a 7 inch band over the row after the seeds have been placed in the furrow and covered. Lightly cultivate the granules into the top 1 inch of soil.	
To Control:	Amt. to Use:
Cutworms	12½ oz. (1½ cup) per 100 ft. of row

12/12

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

CONTAINER IS FILLED BY WEIGHT, NOT BY VOLUME. Some settling of contents may occur.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label **DIRECTIONS** and **PRECAUTIONS**.

DURSBAN® is a registered trademark of DowElanco.

Distributed by
DRAGON CORPORATION
Roanoke, VA 24019-7311

BEST AVAILABLE COPY