6-178

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W Washington, D.C. 20460

	EPA	Reg	Number
--	-----	-----	--------

Term of Issuance:

Date of Issuance.

Conditional

16-178

December 5, 2002



NOTICE OF PESTICIDE:

x Registration

under FIFRA, as amended:

Reregistration

Name of Pesticide Product:

Dragon Carbaryl 4L Concentrate

Insecticide

Name and Address of Registrant (include ZIP Code):

Dragon Chemical Corporation 71 Carolyn Boulevard Farmingdale, NY 11735

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division great to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named post cide is hereby registered reregistered under the Federal Insecticide. Fungicide and Rodentil, 2e

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7) (A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3).

- Make the following changes to the labeling: 1.
 - Revise the EPA Registration Number to read "EPA Reg. No. 16-178." a.
 - Under Environmental Hazards delete the section "NOTE: Do not discharge effluent...or b. Regional Office of the EPA." This language is reserved for manufacturing use products. Also delete the phrase, "For terrestrial uses". Your product only has terrestrial uses.
 - As stated in our July 29, 2002 letter efficacy data must be submitted or cited to support C. control of the following public health pests: all ticks, mosquitoes, fire ants, bees, cockroaches, fleas, scorpions, spiders, and wasps. Either provide efficacy data or delete these pests wherever they appear on the label.

Signature of Approving Official: Date:	,
Mile	12/05/2002

EPA Form 8570-6

- 2. Storage stability and corrosion characteristics studies must be submitted within 12 months from the date of this registration notice.
- 3. Please submit two (2) copies of your final printed label before you release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling.

A stamped copy of the label is enclosed for your records. If you have questions about this letter, please contact of at 703-605-0716 (by phone), 703-305-6596 (by fax), or at mccann.geri@epa.gov (by E-Mail).

Sincerely,

Geraldine R. McCann

Environmental Protection Specialist Insecticide/Rodenticide Branch Registration Division (7505C)

Enclosures: Stamped Label A-79 Form

DRAGON CARBARYL 4L CONCENTRATE INSECTICIDE

For Agricultural or Commercial Use Only. **ACTIVE INGREDIENT:** Carbaryl (1-naphthyl N-methylcarbamate)....... 43.4% **INERT INGREDIENTS:** 56.6% Total This product contains 4 pounds of Carbaryl per gallon. EPA Reg. No.16-EPA Est. No. 16-VA-1 with COMMENTS In EFA Letter Dated KEEP OUT OF REACH OF CHILDREN Under the Federal Insecticide. Fungicide, and Rodenticide Act. **CAUTION** as amended, for the pesticide registered under EPA Reg. No. SHAKE WELL BEFORE USING See First Aid Below

FIRST AID

Net contents

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by a poison control center or doctor.

Do not give anything by mouth to an unconscious or convulsing person.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15 to 20 minutes.

IF INHALED:

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center, the National Pesticide Information Center (800-858-7378) or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

NOTE TO PHYSICIAN: Carbaryl 4L Concentrate Insecticide contains an N-methyl carbamate that inhibits cholinesterase. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. Gastric lavage may be used if this product has been swallowed. Carbaryl poisoning may occur rapidly after ingestion and prompt removal of stomach contents is indicated.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Improve tissue oxygenation as much as possible before administering atropine to minimize the risk of ventricular fibrillation.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended. To aid in confirmation of diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, absorbed through the skin, inhaled, or if in eyes. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Keep out of reach of children and domestic animals. OVEREXPOSURE MAY CAUSE: Salivation, watery eyes, pinpoint eye pupils, blurred vision, muscle tremors, difficult breathing, excessive sweating, abdominal cramps, nausea, vomiting, diarrhea, weakness, headache. IN SEVERE CASES CONVULSION, UNCONSCIOUSNESS AND RESPIRATORY FAILURE MAY OCCUR. SIGNS AND SYMPTOMS MAY OCCUR RAPIDLY FOLLOWING OVEREXPOSURE TO THIS PRODUCT.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride (PVC) or viton, shoes plus socks and chemical resistant headgear for overhead exposure. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Workers Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

NOTE: Do not discharge effluent containing this product into takes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service for further information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Strictly observe label directions and precautions. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State, County or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE), restricted entry interval (REI), and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride (PVC) or viton, shoes plus socks and chemical-resistant headgear for overhead exposure.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

The area being treated must be vacated by unprotected persons. Keep unprotected persons out of treated areas until sprays have dried.

GENERAL PRECAUTIONS AND RESTRICTIONS

DRAGON CARBARYL 4L CONCENTRATE INSECTICIDE is a suspension of microfine carbaryl insecticide in an aqueous medium. It readily disperses in water to form a spray which may be applied by ground equipment. Do not apply this product by aircraft.

Plant Response Precautions: Application to wet foliage or during periods of high humidity may cause injury to tender foliage. Do not use on Boston ivy, Virginia creeper and maidenhair fern as injury may result. Carbaryl may also injure Virginia and sand pines. The use of adjuvants may increase the potential for crop injury to sensitive crops.

PRE-HARVEST AND GRAZING RESTRICTIONS AND LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable carbaryl residues when this product is used in accordance with label directions. If used as directed, treated forage may be grazed or used as feed for dairy and meat animals without causing illegal residues in meat or milk. Do not apply at greater rates or at more frequent intervals than stated on the label. To do so may result in illegal residues in crops, meat and milk. Do not use reclaimed irrigation water from crops treated with carbaryl on crops for which carbaryl tolerances are not established. Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

APPLICATION STATEMENTS

Calibrate and adjust application equipment to ensure proper rate and accurate placement. To clean spray system after use, drain and flush with water and detergent mixture. Rinse thoroughly with clean water. Refer to the "STORAGE AND DISPOSAL" section for disposal instructions.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Spray deposits on painted or stained surfaces or finishes (i.e., cars, houses, trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid applications to surfaces where visible spray residues are objectionable.

RESISTANT SPECIES NOTICE: All references to armyworms on the crops listed on this label refer to the species, *Pseudaletia unipuncta*, often called the "true armyworm". Except where indicated otherwise, this product is not registered for the control of other armyworm species. Regional differences have been noted in the susceptibility of certain strains of Colorado potato beetles, diamondback moths, fall armyworms and Southern green stink bugs to carbaryl. If local experience indicates inadequate control, use an alternative pesticide.

MIXING, LOADING AND HANDLING INSTRUCTIONS

TO ASSURE A UNIFORM SUSPENSION, AGITATE, STIR OR RECIRCULATE ALL CONTAINERS OF THIS PRODUCT PRIOR TO USE. Remove oil, rust, scale, pesticide residues and other foreign matter from mix tanks and entire spray system. Flush with clean water. Fill spray or mix tank with ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of this product and then the remaining volume of water. Include rinse water from container. Prepare only as much spray mixture as can be applied on the day of mixing. MAINTAIN CONTINUOUS AGITATION DURING MIXING AND APPLICATION TO ASSURE A UNIFORM SUSPENSION. DO NOT STORE SPRAY MIXTURE FOR PROLONGED PERIODS OR DEGRADATION OF CARBARYL MAY OCCUR. Local water conditions may also accelerate the degradation of spray mixtures containing carbaryl. See the following "COMPATIBILITY STATEMENT".

COMPATIBILITY STATEMENT

CARBARYL 4L CONCENTRATE INSECTICIDE, when diluted with at least an equal volume of water, is compatible with a wide range of pesticides. It is not compatible with diesel fuel, kerosene, fuel oil or aromatic solvents. If compatibility of CARBARYL 4L CONCENTRATE with another product and the resulting crop response is unknown, the mixture should be first tested on a small scale. Curdling, precipitation, greasing, layer formation or increased viscosity are symptoms of incompatibility. Incompatibility will reduce insect control and may cause application and handling difficulties or plant injury. Observe all precautions and limitations on labeling of all products used in mixtures. WHEN PREPARING COMBINATION SPRAYS, FIRST ADD CARBARYL 4L CONCENTRATE TO AT LEAST AN EQUAL VOLUME OF WATER, MIX THOROUGHLY AND THEN ADD COMBINATION PRODUCTS TO THE MIXTURE. DO NOT APPLY TANK MIX COMBINATIONS UNLESS YOUR PREVIOUS EXPERIENCE INDICATES THE MIXTURE IS EFFECTIVE AND WILL NOT RESULT IN APPLICATION PROBLEMS, EXCESSIVE RESIDUES OR PLANT INJURY.

Carbaryl is unstable under highly alkaline conditions and mixtures with strong bases, such as Bordeaux, lime-sulfur and casein-lime spreaders and will result in chemical degradation of the insecticide. Do not use this product in water with pH values above 8.0 unless a buffer is added. If necessary, water should be buffered to neutral (pH = 7.0) before adding this product to the spray tank. Overhead irrigation with alkaline or muddy water after application will also accelerate chemical degradation and may result in reduced insect control.

APPLICATION PROCEDURES AND PRECAUTIONS

On all crops, use sufficient gallonage to obtain thorough and uniform coverage. Observe crop label instructions for **SPECIFIC DIRECTIONS** regarding spray volume where they occur. Calibrate spray equipment to deliver the required amount. The flow rate of CARBARYL 4L CONCENTRATE diluted 1:1 with water is similar to the flow rate of water. Use of 50 mesh slotted strainers in spray system and 25 mesh slotted strainers behind nozzles is recommended.

Ground Application: Apply in sufficient volume for adequate coverage on all crops and sites. To prepare small volumes of spray mixture, use 1/3 fluid ounce (approximately 2 teaspoons) of this product in an adequate amount of water and apply to 500 square feet, where rates of 1 quart per acre are indicated.

Sprinkler Irrigation Systems: Apply this product only through sprinkler irrigation systems including center pivot, and solid set. Do not apply this product through any other type of irrigation system.

Spray Preparation: First prepare a suspension of this product in a mix tank. Fill tank with $\frac{1}{2}$ to $\frac{3}{4}$ the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of this product and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label, per 1 to 4 gallons of water, are recommended.) Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of this product into the irrigation water line so as to deliver the desired rate per acre. The suspension of this product should be injected with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. If you should have any other questions about calibration, you should contact your State Extension Service Specialists, equipment manufacturers or other experts.

Note: When treatment with this product has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

General Precautions for Applications through Sprinkler Irrigation Systems: Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

SPECIFIC USE DIRECTIONS

Crops/Sites: Asparagus, Broccoli, Brussels sprouts, Cabbage, Cauliflower, Kale, Kohlrabi, Mustard greens, Field corn, Popcorn, Sweet corn, Cucumbers, Melons, Squash, Flax, Pasture grass, Rangeland (non grazing), Eggplant, Peppers, Tomatoes, Celery, Dandelion, Endive, Lettuce (Head, Leaf), Spinach, Fresh and Dried beans, Fresh and Dried Peas, Non-cropland (Hedgerows, Rights-of way, Roadsides, Wasteland), Okra, Peanuts, Carrots, Garden beets, Parsnips, Potatoes, Radishes, Grapes, Raspberries, Strawberries, Tobacco, Citrus fruits, Apples, Apricots, Cherries, Nectarines, Peaches, Plums, Forested areas and Rangeland trees, Trees and Ornamentals, Turfgrass, control of specific pests across multiple sites (Grasshoppers, Imported fire ants, adult Mosquito control, Ticks which vector Lyme disease), and pest control around buildings.

INSECT CONTROL

Begin application when insect populations reach recognized economic threshold levels. Consult the Cooperative Extension Service, consultants or other qualified authorities to determine appropriate threshold levels for treatment and specific use information in your area. Where a dosage range is indicated, use the lower rate on light to incorrate infestations, young plants and early instars and use the higher rate on heavy infestations, mature plants, advanced instars and adults. Thorough and uniform spray overage is essential for effective control.

ASPARAGUS

Crop	Pest	Qts. of This Product Per Acre
Asparagus	Apache cicadas, Asparagus	1 to 2
	beetles, Cutworms	
	SPECIFIC DIRECTIONS: Repeat	application as necessary up to a total of
	3 times prior to harvest or a total	of 5 times per crop, but not more often
	than once every 3 days. For cutwo	orm control, this product is most effective
	against species which feed on the	upper portions of the plant.
	Apache cicadas, Asparagus	2 to 4
	beetles, Cutworms	
	SPECIFIC DIRECTIONS: Applica	ation to ferns or brush growth following
	harvest of spears: Repeat applica	tions as necessary, but not more often
	than once every 7 days. Do not n	nake more than a total of 5 applications
	per year to spears and ferns combi	ned.
RESTRICTIONS AND P	RECAUTIONS: Do not apply within 1 day of harv	vest. Do not apply more than a total of 6
ots. Per acre before harv	rest of spears. Do not apply more than a total of 1	0 gts, per acre per vear.

BRASSICA LEAFY VEGETABLE CROPS (Broccoli, Brussels sprouts, Cabbage, Cauliflower, Kale, Kohlrabi, Mustard greens only)

Repeat applications as needed, up to a total of 4 times, but not more often than once every 7 days.

Crop	Pest	Qts. of This Product Per Acre
Broccoli, Brussels sprouts, Cabbage, Cauliflower, Kale,	Flea beetles, Harlequin bugs, Leafhoppers	½ to 1
Kohlrabi, Mustard greens	Armyworms, Aster leafhoppers, Corn earworms, Diamondback moths, Fall armyworms, Imported cabbageworms, Lygus bugs, Spittle bugs, Stink bugs, Tarnished plant bugs	1 to 2

RESTRICTIONS AND PRECAUTIONS: For Broccoli, Brussels sprouts, Cabbage, Cauliflower and Kohlrabi, do not apply within 3 days of harvest. For Kale and Mustard greens, do not apply within 14 days of harvest. Do not apply more than a total of 6 qts. per acre per crop.

CEREAL GRAIN CROPS (Field corn and Popcorn only)

Repeat applications as needed, up to a total of 4 times, but not more often than once every 14 days. Optimum timing and good coverage are essential for effective control

Crop	Pest	Qts. of This Product Per Acre
Field corn, Popcorn	Armyworms, Chinch bugs, Corn	
	earworms, Corn rootworm adults,	
	Fall armyworms, Flea beetles,	1 to 2
	Japanese beetles, Sap beetles,	
	Southwestern corn borers,	
	Leafhoppers	
	SPECIFIC DIRECTIONS: For optim	
	equipment to apply at least 20 gais.	
	toward stalk to provide thorough cover	erage.
	European corn borers	1½ to 2
	SPECIFIC DIRECTIONS: For optime	•
	not apply in less than 15 gals, of wat	er per acre.
	Cutworms, Western bean cutworms	2
	SPECIFIC DIRECTIONS: For optim	um cutworm control, apply in a 12-inch
	band over the row using sufficient vo	lume of water to obtain thorough
	coverage. For broadcast application	
		effective against species which feed
		or western bean cutworms, treat when
		o 100% tassel emergence. Treatment
	after 100% silk emergence will reduc	e effectiveness.

Field corn, Popcorn (continued)

RESTRICTIONS AND PRECAUTIONS: Observe Bee Caution. Do not apply more than a total of 8 qts per acre per crop. Do not apply within 48 days of harvest of grain and fodder or within 14 days of harvest or grazing of forage or silage.

SWEET CORN

Observe Bee Caution. Repeat applications as necessary up to a total of 8 times, but not more than once every 3 days. Optimum timing and good coverage are essential for effective control.

For insects attacking silks and ears, insecticide sprays should be applied starting when first silks appear and continuing until silks begin to dry. During silking, the minimum re-treatment interval (3 days) may not provide adequate levels of protection under conditions of rapid growth or severe pest pressure. The use of an alternative product should be considered in conjunction with this product.

Crop	Pest	Qts. of This Product Per Acre
Sweet corn	Armyworms, Chinch bugs, Corn	
	earworms, Corn rootworm adults,	
	Fall armyworms, Flea beetles,	1 to 2
	Japanese beetles, Leafhoppers,	{
	Sap beetles, Southwestern corn	
	borers	
		mum chinch bug control, use ground
		s. of water per acre and direct spray
	toward stalk to provide thorough or	
	European com borers	1½ to 2
	}	mum European corn borer control, do
	not apply in less than 15 gals. of w	
	Cutworms, Western bean cutworm	
		mum cutworm control, apply in a 12-inch
		volume of water to obtain thorough
		on, use at least 20 gals, per acre. For
	cutworm control, this product is mo	st effective against species which feed
	on the upper portions of the plant.	For western bean cutworms, treat when
	infestation averages 15% and at 9	to 100% tassel emergence. Treatment
	after 100% silk emergence will red	
RESTRICTIONS AND PRI	ECAUTIONS: Do not apply within 2 days of ha	
	in 48 days of harvest of fodder. Do not apply r	
crop.	с ==, с =: //=: : с = ; о = = ; о = = ; о = ; o =	and the state of the draw bot make be-

CUCURBIT VEGETABLES (Cucumbers, Melons, Squash only)

Repeat applications as necessary, up to a total of 6 times, but not more often than once every 7 days.

Crop	Pest	Qts. of This Product Per Acre
Cucumbers, Melons, Squash	Melonworms, Pickleworms	½ to 1
	Cucumber beetles, Flea beetles, Leafhoppers, Squash bugs	1
	morning or late afternoon.	coverage and time sprays for early
RESTRICTIONS AND PRECAUT	IONS: Do not apply within 3 days of har	vest. Do not apply more than a total of
6 qts. per acre per crop. Observe	plant response precautions.	

FLAX

Up to 2 applications per crop may be made, but not more often than once every 14 days.

Crop	Pest	Qts. of This Product Per Acre
Flax	Armyworms	1 to 1½
(DO NOT USE IN CALIFORNIA)		
RESTRICTIONS AND PRECAUTIO	NS: Do not apply within 42	days of harvest for seed or straw. Do not apply
more than a total of 3 qts. per acre p	per crop.	

PASTURE GRASS

Up to 2 applications per year may be made, but not more often than once every 14 days.

Crop	Pest	Qts. of This Product Per Acre
Pasture grass	Armyworms, Chinch bugs, Essex	
	skippers, Fall armyworms, Striped	1 to 1½
	grass loopers, Thrips, Range	
	caterpillars, Range crane flies, Ticks	
	SPECIFIC DIRECTIONS: To control	thrips in grasses grown for seed, use
	high spray pressure to improve penet	ration into the boot. Carefully mark
	swaths to avoid over-application.	
RESTRICTIONS AND PR	ECAUTIONS: Do not apply within 14 days of harv	est or grazing. Do not exceed a total
of 3 qts. per acre per year		

RANGELAND

Do not make more than 1 application per year. Carefully mark swaths to avoid over-application.

Site	Pest	Qts. of This Product Per Acre
Rangeland (non-grazing)	Black grass bugs, Grasshoppers, Mormon crickets, Range caterpillars, Range crane flies	½ to 1
	Ticks	1
RESTRICTIONS AND PRECAL	JTIONS: May be harvested or grazed the	same day of treatment. Do not apply
more than 1 qt. per acre per yea	ar.	

FRUITING VEGETABLES (Eggplant, Peppers, Tomatoes only)

Repeat applications as necessary up to a total of 7 times, but not more often than once very 7 days. When disease transmission is suspected, monitor fields following application and re-treat if reinfestation occurs, but not more often than once every 7 days.

Crop	Pest	Qts. of This Product Per Acre
Eggplant, Peppers, Tomatoes	Colorado potato beetles, European corn borers, Fall armyworms, Lace bugs, Stink bugs (suppression), Tarnished plant bugs, Thrips (suppression), Tomato fruitworms, Tomato hornworms, Tomato pinworms	1 to 2
	SPECIFIC DIRECTIONS: Thorough suppress Stink bugs.	coverage is essential to effectively
	Flea beetles, Leafhoppers	½ to 1
	Cutworms	2
	SPECIFIC DIRECTIONS: For cutwo effective against species which feed	

LEAFY VEGETABLES (Celery, Dandelion, Endive, Lettuce, Spinach only)

Repeat applications as necessary, up to a total of 5 times, but not more often than once every 7 days.

tles, Harlequin bugs, pers ½ to 1 ms, Aster leafhoppers,
ms Aster leafhonners
worms, Fall armyworms, I cabbageworms, Lygus I to 2 wittlebugs, Stink bugs, I do plant bugs

RESTRICTIONS AND PRECAUTIONS: Do not apply within 14 days of harvest. Do not apply more than a total of 6 qts. per acre per crop.

LEGUME VEGETABLES (Fresh and Dried beans, Fresh and Dried peas only)

Repeat applications as necessary, up to a total of 4 times, but not more often than once every 7 days. Use lower rates for light to moderate populations and smaller instars and to provide maximum survival of beneficial insects

and spiders. Use the higher rates for heavy populations and larger instars

	heavy populations and larger instars.	
Crop	Pest	Qts. of This Product Per Acre
Fresh and Dried beans (Phaseolus species including Kidney, Navy and	Bean leaf beetles, Blister beetles, Cucumber beetles, Grape colapsis,	
Snap), Fresh and Dried peas	Green cloverworms, Japanese	½ to 1
(Pisum species)	beetles, Mexican bean beetles,	
_	Velvetbean caterpillars	· ·
	Corn earworms	½ to 1½
	Alfalfa caterpillars, Colorado potato	
	beetles, Flea beetles, Leafhoppers,	1
	Three cornered alfalfa hoppers,	
	Thrips, Western bean cutworms	
	Armyworms, Cutworms, European	4.1412
	corn borers, Fall armyworms, Stink	1 to 1½
	bugs, Tarnished plant bugs, Webworms	
	SPECIFIC DIRECTIONS: For cutwor	m control this product is most
	effective against species which feed of	•
	Alfalfa loopers (suppression),	Straine appel postions of the plant.
	Cowpea curculios (suppression),	
	Painted ladies (Thistle caterpillars),	1½
	Pea leaf weevils, Pea weevils,	
	Saltmarsh caterpillars, Woollybean	
	caterpillars, Yellowstriped	
	armyworms	
	CALIFORNIA ONLY:	
	Corn earworms (suppression),	
	Limabean podborers (suppression),	1½
	Lygus bugs (suppression), Stink	
DESTRICTIONS AND RECAUTION	bugs (suppression) IS: Do not apply within 14 days of graz	ing a harman factor of a second second second
RESTRICTIONS AND PRECAUTION	ia: Do not abbiy within 14 days of dtaz	ing of narvest for torage of Within 3

RESTRICTIONS AND PRECAUTIONS: Do not apply within 14 days of grazing or harvest for forage or within 3 days of harvest of fresh beans or peas or within 21 days of harvest of dried beans or peas, seed or hay. Do not apply more than a total of 6 qts. per acre per crop. Observe plant response precautions.

NON-CROPLAND

Up to 2 applications per year may be made, but not more often than once every 14 days. Carefully mark swaths

to avoid ov	erapplication.
-------------	----------------

Site	Pest	Qts. of This Product Per Acre
Wasteland, Rights-of-Way, Hedgerows, Roadsides	Black grass bugs	1/4 to 1/2
	Morman crickets, Range caterpillars, Range crane fly	½ to 1
	Ticks	1 to 1½
RESTRICTIONS AND PRECAU	TIONS: Do not apply within 14 days of	

RESTRICTIONS AND PRECAUTIONS: Do not apply within 14 days of grazing or harvest for forage or hay. Do not apply more than a total of 3 qts. per acre per year.

OKRA

Apply in sufficient volume to ensure good coverage.

Crop	Pest	Qts. of This Product Per Acre
Okra	Corn earworms, Stink bugs	1 to 1½
	grasshopper control, refer to the ACROSS MULTIPLE SITES" sec	
RESTRICTIONS AND	PRECAUTIONS: Do not apply within 3 days of h	narvest. Do not apply more than a total
6 qts. per acre per se	ason.	

PEANUTS

Repeat applications as necessary, up to a total of 5 times, but not more often than once every 7 days.

Crop	Pest	Qts. of This Product Per Acre		
Peanuts	Blister beetles, Mexican bean	½ to 1		
	beetles			
	Alfalfa caterpillars, Bean leaf	1		
	beetles, Cucumber beetles, Green			
	cloverworms, Japanese beetles.			
	Leafhoppers, Rednecked			
	peanutworms, Three cornered			
	alfalfa hoppers, Thrips, Velvetbean			
	caterpillars.			
	SPECIFIC DIRECTIONS: For optimu	SPECIFIC DIRECTIONS: For optimum control of thrips, use directed or		
	banded sprays with hollow cone spray	y nozzles. Ensure adequate		
	coverage for the underside of leaves.			
	Armyworms, Corn earworms, Fall	1 to 1½		
	armyworms, Stink bugs, Webworms			
	Whitefringed beetle adults.	2		
	Cutworms			
	SPECIFIC DIRECTIONS: For cutwor	SPECIFIC DIRECTIONS: For cutworm control, this product is most		
	effective against species which feed of	effective against species which feed on the upper portions of the plant.		
RESTRICTIONS AND F	PRECAUTIONS: Observe plant response precaution	is. Do not apply within 14 days of		
iarvest. Do not apply n	nore than a total of 8 gts. per acre per crop.			

ROOT AND TUBER CROPS (Carrots, Garden beets, Parsnips, Potatoes, Radishes only)

Repeat applications as necessary,	up to a total of 6 time	nes, but not more	often than once every 7 days.

Pest	Qts. of This Product Per Acre
Flea beetles, Leafhoppers	½ to 1
Armyworms, Aster leafhoppers,	
Colorado potato beetles, Corn	
earworms, Cutworms, European	1 to 2
corn borers, Fall armyworms, Lace	
bugs, Lygus bugs, Spittlebugs,	
Stink bugs, Tarnished plant bugs	
SPECIFIC DIRECTIONS: For cutworm control, this product is most	
effective against species which feed on the upper portions of the plant.	
	Flea beetles, Leafhoppers Armyworms, Aster leafhoppers, Colorado potato beetles, Corn earworms, Cutworms, European corn borers, Fall armyworms, Lace bugs, Lygus bugs, Spittlebugs, Stink bugs, Tarnished plant bugs SPECIFIC DIRECTIONS: For cutwo

SMALL FRUITS AND BERRIES (Grapes, Raspberries, Strawberries only)

Repeat applications as necessary, up to a total of 5 times, but not more often than once every 7 days.

Crop	Pest	Qts. of This Product Per Acre
Grapes, Raspberries, Strawberries	European fruit lecaniums, European raspberry aphids, Flea beetles, Grape leaffolders, Grape leafrollers, Japanese	
	beetles, Leafhoppers, Leafrollers, Meadow spittlebugs, Omnivorous leafftiers, Rose chafers, Snowy tree crickets, Strawberry bud weevils, Strawberry clippers, Strawberry fruitworms, Strawberry leafrollers, Strawberry weevils, Western grapeleaf skeletonizers, Western yellowstriped	1 to 2
	armyworms	
	SPECIFIC DIRECTIONS: In grapes, for grabefore the first brood larvae emerge from ro	
	Blueberry maggots, Cherry fruitworms, Cranberry fireworms, Cranberry fruitworms, Cranberry twig girdlers, Elm spanworms, Gypsy moths, Spaganothus worms, Tarnished plant bugs	1½ to 2
	Eight-spotted foresters, Cytworms, Grape berry moths, June beetles, Omnivorous leafrollers, Orange tortrix, Raspberry fruitworms, Raspberry sawflies, Redbanded leafrollers, Saltmarsh caterpillars	2
	SPECIFIC DIRECTIONS: For cutworm con effective against species which feed on the	
	Change Bas Couties Do not anniversity	

RESTRICTIONS AND PRECAUTIONS: Observe Bee Caution. Do not apply within 7 days of harvest. Do not apply more than a total of 10 qts. per acre per crop.

Caution: Carbaryl may injure Early dawn and Sunrise varieties of strawberries. In grapes, do not concentrate spray on the bunch or visible residues may result.

TOBACCO

Plant bed and Field Treatment – Repeat treatments as necessary, up to a total of 4 times per crop, but not more often than once every 7 days. Use lower rate on young plants (up to knee height). Use at least 10 gallons of prepared spray per acre. Begin treatments when worms are small.

Crop	Pest	Qts. of This Product Per Acre
Tobacco	Budworms, Fall armyworms, Hornworms, Japanese beetles, June beetles, Suckflies, Tobacco flea beetles	1 to 2
	CAUTIONS: Tobacco may be harvested on the day of to	reatment. Do not apply more

TREE FRUIT CROPS (Citrus fruits only)

On all tree fruit crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, density and other factors. Repeat applications as necessary, up to a total of 8 times, but not more often than once every 14 days.

Crop	Pest	Qts. of This Product Per Acre
Citrus fruits	Avocado leafrollers, California orangedogs, Citrus cutworms, Fruittree leafrollers, Orange tortrix, Western tussock moths	2 to 3
	Citrus rust mites, Eriophyid mites, Plant bugs, Scale insects [Black scales, Brown soft scales, California red scales (except in California), Citrus snow scales, Yellow scales (except in California)]	3 to 5
	SPECIFIC DIRECTIONS: For scale control, a present. For best control of Eriophyid mites, in apply when pest populations are low.	· · •
	Apopka weevils (adult), Citrus root weevils (adults), Fuller rose beetles, Little leaf notchers (adult)	5 to 7½
	California Only: California red scales, Yellow scales	5 to 16
SPECIFIC DIRECTIONS: Do not make more than 1 season for California red scales. Apply when crawler		

RESTRICTIONS AND PRECAUTIONS: Observe Bee Caution. Do not apply within 5 days of harvest. Do not apply more than a total of 20 qts. per acre per crop.

POME FRUITS (Apples only)

Repeat applications as necessary, up to a total of 8 times per crop (including thinning sprays on apples), but not

more often than once every 14 days.

Crop	Pest	Qts. of This Product Per Acre	
Apples	White apple leafhoppers	½ to 1½	
	Apple aphids, Codling moths	1 to 3	
	Apple aphids, Apple maggots, Apple mealybugs, Apple rust mites, Bagworms, California pearslugs (Pear sawflies), European apple sawflies, Eyespotted bud moths, Fruittree leafrollers, Gypsy moths, Japanese beetles, Lesser appleworms, Lygus bugs, Orange tortrix, Pearleaf blister mites, Pear psyllas, Pear rust mites, Periodical cicadas, Plum curculios, Redbanded leafrollers, Rosy apple aphids, Scale insects (Forbes scales, Lecanium scales, San Jose scales), Tarnished plant bugs, Tentiform leafminers, Wooly apple aphids,	1½ to 3	
	Yellowheaded fireworms		
	bloom until 30 days after full bloom unless f pest control during this period also may res	SPECIFIC DIRECTIONS: On apples, avoid use during the period from full bloom until 30 days after full bloom unless fruit thinning is desired. Use for pest control during this period also may result in fruit removal. For psylla control apply when eggs hatch or young nymphs are present. For scale control, apply when crawlers are present.	
Apples Only, for fruit thinning		1 to 3	

SPECIFIC DIRECTIONS: Observe Bee Caution. Apply between 80% petal fall and 16 mm fruit size. Use the higher rates on hard to thin varieties at the early timing which is the 80% petal fall to 6 mm fruit size. Use the lower rates on easy to thin varieties and at the later thinning period which is the 10 to 16 mm fruit size. The rate to use per acre will depend on varieties, tree size, row spacing, weather conditions at the time of and following applications. Consult with the local fruit thinning experts in your area for the proper rate to use under your conditions.

The optimum spray gallonage will depend on the tree size, planting density, row spacing and amount of foliage. Use sufficient spray volume to ensure adequate coverage (100 to 400 gals, per acre). Avoid spraying to the point of runoff. Reduce spray coverage to the lower portion of the tree since overthinning may occur here. Factors such as climatic temperature, high humidity, frost, tree age, variety, nutrition, previous crop, pruning and bloom may influence fruit thinning results with the product. Exercise caution to avoid possible overthinning. For the most effective results, apply under good drying conditions and when daytime temperatures (°F) will be 70 to low 80's for the following 1 to 3 days. Application with daytime temperatures in excess of 80°F may result in overthinning. This product may be mixed with other fruit thinners; however, use caution to avoid overthinning and other adverse effects. Consult with local fruit thinning experts in your area for recommendations. Refer to the other product labels for specific use directions.

Consult with local fruit thinning experts in your area for advice on the proper use of this product on your varieties under your growing conditions.

PRECAUTIONS: The use of this product may result in fruit deformity under certain environmental conditions. Before using on any variety of apples, the user must weigh the risk versus benefits when using this product, particularly when using between 30% petal fall and 6 mm fruit size. Red Delicious are more sensitive to this phenomenon and in particular, the varieties Bisbee, Red Chief and Vallee Spur are very susceptible to conditions causing fruit deformity. Precipitation and temperatures below 65°F increase the possibility of fruit deformity. The use with summer spray oils and wetting agents may increase the risk of fruit deformity and injury.

POME FRUITS Continued

RESTRICTIONS AND PRECAUTIONS: Do not apply within 3 days of harvest. Do not apply more than a total of 15 qts. per acre per crop. Do not make more than a total of 8 applications per crop.

FOR PROTECTION OF HONEYBEES: Remove all bee hives from orchard to be treated prior to application. Do not apply this product if bees are actively foraging in orchard. If weed bloom is present, mow the cover crop on the orchard floor prior to applying this product.

STONE FRUITS (Apricots, Cherries, Nectarines, Peaches, Plums only)

STONE PROTTS (Aprilcots, Cherries,		Qts. of This
Сгор	Pest	Product Per Acre
Apricots, Cherries, Nectarines,	Apple pandemis, Black cherry aphids,	
Peaches, Plums	Cherry fruitworms, Cherry maggots	
	(Cherry fruit flies), Codling moths,	0.40
	Cucumber beetles, Eastern tent	2 to 3
	caterpillars, Eyespotted bud moths,	
	European earwigs, Fruittree leafrollers,	
	Green fruitworms, Gypsy moths,	
	Japanese beetles, June beetles, Lesser	
	peachtree borers, Mealy plum aphids,	
	Orange tortrix, Oriental fruit moths, Peach	
	twig borers, Periodical cicadas, Plum	
	curculios, Prune leafhoppers, Redbanded	
	leafrollers, Rose chafers, Scale Insects	
	(Brown soft scales, Forbes scales,	
	Lecanium scales, Olive scales,	
	Oystershell scales, San Jose scales), Tarnished plant bugs, Tussock moths,	
	Variegated leafrollers	
	SPECIFIC DIRECTIONS: Observe Bee Ca	ution Popost applications as
	necessary, up to a total of 3 times per crop,	
	every 7 days. An additional application at the	
	timing may be made. For optimum scale co	
	present. For lesser peachtree borers, best	
	thoroughly spraying limbs and tree trunks at	
	flight.	would make adming would
	CALIFORNIA ONLY: Black cherry	
	aphids, Cherry fruitworms, Cherry	
	maggots (Cherry fruit flies), Codling	
	moths, Cucumber beetles, Eyespotted	3 to 4
	bud moths, European earwigs, Fruittree	
	leafrollers, Green fruitworms, Mealy plum	
	aphids, Orange tortrix, Oriental fruit	
	moths, Peach twig borers, Scale insects	
	(Brown soft scales, Forbes scales,	
	Lecanium scales, Olive scales,	
	Oystershell scales, San Jose scales),	
	Tarnished plant bugs, Tussock moths	
	SPECIFIC DIRECTIONS: Observe Bee Ca	ution. Repeat applications as
	necessary, up to a total of 3 times per crop,	
	every 14 days. An additional application at	
	dormant timing may be made. For optimum	scale control, apply when
	crawlers are present.	
	Peach twig borers, Scale insects (Brown	
	soft scales, Forbes scales, Lecanium	4 to 5
	scales, Olive scales, Oystershell scales,	
	San Jose scales)	
	SPECIFIC DIRECTIONS: For dormant or d	
	combination with a recommended dormant	
	product label for additional use directions ar	
RESTRICTIONS AND PRECAUTION	S: Do not apply within 3 days of harvest, exc	cept in California. In California,

RESTRICTIONS AND PRECAUTIONS: Do not apply within 3 days of harvest, except in California. In California, do not apply within 1 day of harvest. Do not apply more than a total of 14 qts. per acre per crop. Do not apply more than a total of 5 qts. per acre at the dormant or delayed dormant timing. Do not apply more than a total of 9 qts. per acre during the production season.

FORESTED AREAS AND RANGELAND TREES

Apply in sufficient volume for adequate coverage. This will vary depending on the tree size, density and stage of

growth. Crop/Site	Pest	Qts. of This Product Per Acre
Non-urban forests, Tree plantations,	Ants, Apple aphids, Armyworms, Ash	
Planted Christmas trees, Rangeland	whiteflies Azalea leafminers, Bagworms,	
trees	Balsam twig aphids, Birch leafminers,	1
-	Blister beetles, Boxelder bugs, Boxwood	
•	leafminers, Browntail moths,	
•	Cankerworms, Catalpa sphinx, Chiggers,	
	Cooley spruce gall adelgids, Cutworms,	
	Cypress tip moths, Douglas fir tussock	
,	moths, Eastern spruce gall adelgids, Elm	
	leaf aphids, Elm leaf beetles, Elm	
	spanworms, Eriophyld mites, European	
	pine shoot moths, Fall armyworms, Flea	
	beetles, Fuchia gall mites, Fuller rose	
	beetles, Gall midges, Gall wasps,	
	Greenstriped mapleworms,	
	Grasshoppers, Hackberry nipplegall	
	makers, Holly bud moths, Holly	
	leafminers, Jackpine budworms,	
	Japanese beetles, Jeffrey pine	
	needleminers, June beetles, Lace bugs,	
	Leafhoppers, Leafrollers, Locust borers,	
	Maple Leafcutters, Mealybugs, Mimosa	
	webworms, Nantucket pine tip moths, Oak	
	leafminers, Oak moths, Oak	
	skeletonizers, Oakworm (complex),	
	Oleander caterpillars. Olive ash borers,	
	Orangestriped oakworms, Periodical	
	cicadas, Pine loopers, Pine sawflies, Pine	
	spittlebugs, Pitch pine tip moths, Plant	
	bugs, Poinsettia hornworms, Psyllids,	
	Puss caterpillars, Redhumped oakworms,	
	Rose aphids,/Rose chafers, Rose slugs,	
	Saddled prominents, Sawflies (exposed),	
	Scale insects (crawlers), Sowbugs,	
	Spiney elm caterpillars, Springtails,	
	Spruce budworms, Spruce needleminers,	
	Subtropical pine tip moths, Tent	
	caterpillars, Thornbugs, Thrips (exposed),	
	Ticks, Walnut caterpillars, Webworms,	
	Western hemlock loopers, Western	
	spruce budworms, Willow leaf beetles,	
	Woolly gall aphids, Yellow-poplar weevils	

FORESTED AREAS AND RANGELAND TREES Continued

SPECIFIC DIRECTIONS: Observe plant response precautions. Obtain thorough coverage of upper and lower leaf surfaces. The addition of a sticker may improve residual control. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. For optimum worm control, treat when pests are small. Do not use on syrup-producing sugar maples where sap is harvested. Applications for control of maple leafcutters on sugar maple should be made when larvae are in 2nd instar after mining and as cases are being formed. Repeat treatments as necessary, up to a total of 2 times per year, but not more often than once every 7 days.

Gypsy moths 3/4 to

SPECIFIC DIRECTIONS: For gypsy moth control, use the higher rate for heavy infestations.

Elm bark beetles, lps engraver beetles, Mountain pine beetles, Roundheaded pine beetles, Spruce beetles, Western pine beetles 2% solution (5 fl. ozs. per gal.)

SPECIFIC DIRECTIONS: Direct Trunk Treatment – Effective as a preventative treatment only. Repeat annually as required to prevent beetle attacks. Apply 1 gal. of spray per 50 sq. ft. of bark prior to beetle flight or host-tree attack. Treat tree trunk from ground level up, until trunk diameter is less than 5 inches. For elm bark beetles, apply approximately 20 to 30 gals. of spray mixture for each 50 ft. of elm tree for thorough coverage of all bark surfaces on trunks, limbs and twigs. Do not make more than 2 applications per year or repeat applications more often than once every 6 months.

RESTRICTIONS AND PRECAUTIONS: Do not make more than 2 applications per year. To prepare small volumes of spray mixture, use 1/3 fl. oz. (approx. 2 tsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1 qt. per acre are indicated. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.

NOTE: All references to armyworms on the preceding list of plants refer to the species *Pseudaletia unipuncta*, often called the "true armyworm". Except where indicated otherwise, this product is not registered for control of other Armyworm species. Regional differences have been noted in the susceptibility of certain strains of fall armyworms to carbaryl. If local experience indicates inadequate control, use an alternative pesticide.

TREES AND ORNAMENTALS

Apply in sufficient volume for adequate coverage. This will vary depending on the plant or tree size, density and stage of growth. Do not use on Boston ivy, Virginia creeper and maidenhair fern. During early season, it may also injure Virginia and sand pines.

Cranisita	Pont	Qts. of This
Crop/Site	Pest	Product Per Acre
Trees, Ornamentals: (Roses,	Ants, Apple aphids, Armyworms, Ash	4
Flowers, Other Herbaceous plants),	whiteflies, Azalea leafminers, Bagworms,	1 qt.
Woody plants, Shrubs in	Birch leafminers, Blister beetles, Boxelder	-or-
commercial and residential areas	bugs, Boxwood leafminers, Browntail	1 fl. oz per
and in nursery production areas	moths, Cankerworms, Catalpa sphinx,	3 gals. of spray
	Chiggers, Cooley spruce gall adelgids,	-or-
	Cutworms, Cypress tip moths, Eastern	2 tsps. per
	spruce gall adelgids, Elm leaf aphids, Elm	gal. of spray
	leaf beetles, Elm spanworms, Eriophyid	
	mites, European pine shoot moths, Fall	
	armyworms, Flea beetles, Fuchia gall	
	mites, Fuller rose beetles, Gall midges,	
	Gall wasps, Greenstriped mapleworms,	
	Grasshoppers, Hackberry nipplegall	
	makers, Holly bud moths, Holly	
	leafminers, Jackpine budworms,	
	Japanese beetles, Jeffrey pine	
	needleminers, June beetles, Lace bugs,	
	· · · · · · · · · · · · · · · · · · ·	
	Leafhoppers, Leafrollers, Locust borers,	
	Maple leafcutters, Mealybugs, Mimosa	
	webworms, Nantucket pine tip moths, Oak	
	leafminers, Oak moths, Oak	
	skeletonizers, Oakworm (complex),	
	Oleander caterpillars, Olive ash borers,	
	Orangestriped oakworms, Periodical	
	cicadas, Pine toopers, Pine sawflies, Pine	
	spittlebugs, Pitch pine tip moths, Plant	
	bugs, Poinsettia hornworms, Psyllids,	
	Puss caterpillars, Redhumped oakworms,	
	Rose aphids, Rose chafers, Rose slugs,	
	Saddled prominents, Sawflies (exposed),	
	Scale insects (crawlers), Sowbugs, Spiney	
	elm caterpillars, Springtails, Spruce	
	needleminers, Subtropical pine tip moths,	
	Tent caterpillars, Thornbugs, Thrips	
	(exposed), Ticks, Walnut caterpillars,	
	Webworms, Western hemlock loopers,	
	Willow leaf beetles, Wooly gall aphids,	
	Yellow-poplar weevils	
	SPECIFIC DIRECTIONS: Observe plant res	•
	thorough coverage of upper and lower leaf s	
	sticker may improve residual control. To co	
	stems and twigs in addition to plant foliage.	For optimum worm control,
	treat when pests are small. Do not use on s	syrup-producing sugar maples
	where sap is harvested. Applications for co	
	sugar maple should be made when larvae a	
	as cases are being formed. Repeat treatme	
	of 2 times per year, but not more often than	
	5. 2 thirds por your, but not more often than	once every i days.

TREES	AND	ORN	AMENT	CALS	Continued

TREES AND URNAMENTALS CONT	ilined	
	Balsam twig aphids, Cooley spruce gall	1 to 2 qts.
	adelgids, Douglas-fir tussock moths,	-or-
	Spruce budworms, Western spruce	2 to 4 tsps. per gal.
	budworms	of spray
	SPECIFIC DIRECTIONS: Use higher rate v	vhen large larvae or heavy
	populations are present.	
	Gypsy moths	3⁄4 to 1
	SPECIFIC DIRECTIONS: For Gypsy moth	control, use the higher rate for
	heavy infestations.	
,.	Elm bark beetles, lps engraver beetles,	2% solution
	Mountain pine beetles, Roundheaded	(5 fl. ozs.
i	pine beetles, Spruce beetles, Western	per gal.)
	pine beetles	
	SPECIFIC DIRECTIONS: Direct Trunk Trea	atment – Effective as a
	preventative treatment only. Repeat annual	ly as required to prevent beetle
	attacks. Apply 1 gal. of spray per 50 sq. ft. of	of bark prior to beetle flight or
	host tree attack. Treat tree trunk from grour	
	is less than 5 inches. For elm bark beetles,	
	gals, of spray mixture for each 50 ft, of elm t	
	all bark surfaces on trunks, limbs and twigs.	
	applications per year or repeat applications	i
	, , , , , , , , , , , , , , , , , , , ,	inote offer than office every o
	months.	

RESTRICTIONS AND PRECAUTIONS: Do not make more than 2 applications per year. To prepare small volumes of spray mixture, use 1/3 fl. oz. (approx. 2 tsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1 qt. per acre are indicated. DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.

TURFGRASS

Apply the recommended rates using pressure-type equipment in sufficient spray volume for thorough coverage and thatch penetration.

Site	Pest	Qts. of This Product Per Acre	
Golf turf, Sports	Ants, Armyworms, Centipedes, Chiggers,		
fields, Domestic	Cutworms, Earwigs, Essex skippers, European		
and	chafers, Fall armyworms, Fiery skippers,	2 to 4	
Commercial	Grasshoppers, Green June beetle larvae,	(1½ to 3 fl. ozs. per 1,000 sq. ft.)	
lawns,	Leafhoppers, Lucerne moths, Millipedes,		
Cemeteries	Mosquitoes (adults), Scarab beetle adults (May		
}	beetles, June beetles, Japanese beetles, Green		
	June beetles), Sowbugs, Spittlebugs, Springtails,	[
	Ticks, Yellowstriped armyworms		
	SPECIFIC DIRECTIONS: For armyworm, cutworm as		
ĺ	irrigate treated areas within 24 hours following insection		
	beetle grub (larvae) control: Make applications when		
	Water or irrigate turfgrass soon after treatment. Repe	at applications as necessary, up to a	
}	total of 4 times a year, but not more often than once e	very 7 days.	
	Chinch bugs, Sod webworms (Lawn moths)	6 to 8	
		(4½ to 6 fl. ozs. per 1,000 sq. ft.)	
ļ	SPECIFIC DIRECTIONS: For chinch bug control: Treat entire grass area rather than just		
	damaged areas. Irrigation prior to treatment will aid in penetration of insecticide into		
	turfgrass. Do not irrigate treated areas within 24 hour		
	Repeat applications as necessary, up to 2 times per y 7 days.	ear, but not more often that once every	
	Bluegrass billbug larvae, European crane flies,		
	Fleas, White grubs (such as Japanese beetles,	8	
	Chafer beétles and <i>Phyllophaga</i> spp. larvae), Black	(6 fl. ozs. per 1,000 sq. ft.)	
	turfgrass ataenius beetle larvae, Hyperodes weevil		
	(Bluegrass weevil) larvae		
	SPECIFIC DIRECTIONS: For European crane fly cor	ntrol: Treatments should be applied in	
	early Spring (April 1 to April 15) or as recommended by	y local Agricultural Extension Service	
	agents. Water or irrigate turfgrass soon after treatment. For white grub control: Applications		
	should be made when grubs are feeding near the soil surface, usually during late March		
	through May, or July through early September, or as recommended by local Agricultural		
	Extension Service agents. Water or irrigate turfgrass		
	applications as necessary, up to 2 times per year, but		
	days.		
	AND PRECAUTIONS: DO NOT ALLOW PUBLIC USE	OF TREATED AREAS DURING	
APPLICATIONS (OR UNTIL SPRAYS HAVE DRIED.		

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES

GRASSHOPPERS

Crop	Pest	Qts. of This Product Per Acre
All crops on this	Grasshoppers	½ to 1½ *
label	SPECIFIC DIRECTIONS: Apply ½ to ¾ qt. per plants or sparse vegetation. Apply 1 to 1½ qts. applications to dense foliage or if extended resivolumes are appropriate to assure adequate contents.	per acre for mature grasshoppers or idual control is desired. Be certain spray
	ndividual site listing elsewhere on this label for u listed for the site or exceed other use restrictions	

CONTROL OF TICKS WHICH VECTOR LYME DISEASE

For control of juvenile and adult ticks which vector Lyme disease, apply the recommended amount in sufficient volume for thorough coverage:

Crop/Site	Pest	Qts. of This Product Per Acre
All crops on this label, Pastures, Forested areas, Wasteland, Rights-of-Way, Hedgerows, Roadsides, Trees	Ixodes spp. (Deer ticks, Bear ticks, Black legged ticks), Amblyomma spp. (Lone star ticks)	1 to 2 * (2 to 4 tsps. per gal.)
and Ornamentals, Turfgrass, Lawns and Recreational Turfgrasses (including Lawns and Perimeters, Golf courses, Sports fields, Cemeteries, and Pastures), Shrubs, Wooded areas	(Lone star ticks) SPECIFIC DIRECTIONS: Use the higher rate for heavy tick infestations: Use higher spray volumes for dense ground cover or heavy leaf litter. Target applications for nymphal control in late spring or early summer. Control of adult ticks can be obtained with late summer and fall application Do not use spot treatments. Treat entire area and perimeter areas where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Re-treat as necessary to maintain adequate contilevels*.	

RESTRICTIONS AND PRECAUTIONS FOR CONTROL OF TICKS WHICH VECTOR LYME DISEASE:

* Note: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions.

DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATION OR UNTIL SPRAYS HAVE DRIED.

IMPORTED FIRE ANTS

Crop/Site	Pest	Qts. of This Product Per Volume of Water		
Pastures, Rangelands,	Imported fire ants % fl. oz. per gal.			
and Ornamentals, Turfgrass, Wasteland, Lawns, Cemeteries, Golf courses	SPECIFIC DIRECTIONS: Apply a total of 2 gals. of the diluted solution over surface of each mound or at least 1 qt. per 6 inches of mound diameter us bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding areas to a 4-ft diameter (12 sq. ft.). Do not disturb mound principle.			
Nursery stock, Vegetable transplants*, Foliage	le Imported fire ants 1½ qts. per 100 gals.			
plants, Bedding plants (Outdoor use only)	SPECIFIC DIRECTIONS: Avoid contact with foliage and treat only the growing media when using on bedding plants. Do not make more than one application either as a root dip or drench treatment (applied to the point of saturation).			
RESTRICTIONS AND PRI	CAUTIONS: DO NOT ALLOW PUBL	IC USE OF TREATED AREAS DURING		

RESTRICTIONS AND PRECAUTIONS: DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL DRENCH HAS DRIED. DO NOT USE IN GREENHOUSES.

To prepare small amounts, use $\frac{3}{4}$ fl. oz. (approx. $\frac{1}{2}$ tbsps.) of this product per each gal. of mix where $\frac{1}{2}$ qts. per 100 gals, are indicated.

* Note: DO NOT USE ON ANY FOOD CROP NOT LISTED ON THIS LABEL.

Refer to specific crop section for additional restrictions and precautions.

ADULT MOSQUITO CONTROL

Apply in sufficient gallonage for thorough coverage.

Crop/Site	Pest	Qts. of This Product Per Acre
Pastures, Rangeland,	Mosquitoes (adults)	1/4 to 1*
Forested lands, Trees and Ornamentals, Turfgrass, Wastelands	SPECIFIC DIRECTIONS: Observe Bee Ca adult mosquitoes congregate. Treat when a mornings or late evenings. Repeat applicat 100 gals. in mistblowers and 1 qt. per acre in For Residual Control in Subtropical Region 2,000 sq. ft. of surface area. Repeat in 3 to	adult mosquitoes are active in early tions as necessary*. Use ¼ to ½ qt. per in low pressure ground sprayers. ions – Apply 4 gals. of prepared spray per
PESTRICTIONS AND P	RECAUTIONS: DO NOT ALLOW PUBLIC U	SE OF TREATED AREAS DURING

APPLICATION OR UNTIL SPRAYS HAVE DRIED. Caution: May kill shrimp and crabs. Do not use in areas where these are important resources. To prepare small volumes of spray mixture, use 1/3 fl. oz. (approx. 2 tsps.) of this product in an adequate amount of water and apply to 500 sq. ft. where rates of 1 qt. per acre are indicated. * Note: Refer to individual site listings elsewhere on this label for use limitations and restrictions. Do not use rates higher than listed for the site or exceed other use restrictions.

NUISANCE PEST CONTROL

For use as a perimeter treatment to prevent infestations of buildings, apply the specified dosage in a band 6 to 10 feet wide around the outside perimeter of the structure.

Pest	Rate
Ants, Bees, Brown dog ticks, Centipedes, Cockroaches, Crickets, Darkling beetle (lesser mealworm or litter beetle), Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Spiders, Wasps	1% solution or 2½ fl. ozs. per gal. of water

SPECIFIC DIRECTIONS: Observe plant response precautions. Apply as a coarse wet spray in a band 6 to 10 feet wide around the outside perimeter of buildings. Confine applications to areas immediately adjacent to the building. Direct application to the structure should be minimal and restricted to cracks, crevices, and areas where insects tend to congregate. For use in and around flowerbeds and ornamental plantings, apply as a coarse wet spray in and around plant beds. May be applied to outdoor sleeping quarters of pets. Do not treat animals. Do not allow animals to use treated area until sprays have dried.

RESTRICTIONS AND PRECAUTIONS: NUISANCE PEST CONTROL

DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED. Do not use in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed.

Note: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits on painted or stained surfaces of finishes (e.g. cars, house trailers, boats, etc.) should be removed immediately by washing to prevent discoloration. Avoid application to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

PESTICIDE STORAGE: store unused product in original container only, in cool, dry areas out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F. If container is damaged, before cleaning up, put on Personal Protective Equipment (PPE).

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of applications, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumers all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is condition of sale by the Seller and is accepted as such by the Buyer.