

SINCE
1859

COSTELLO

SINCE
1859

DIRECTIONS

Use at the rate of one gallon with water to make 100 gallons of spray. Apply with pressure, hand or power sprayer in sufficient quantity to give thorough coverage of the weeds without excessive run-off or wetting of the soil. For turf, approximately 150 to 200 gallons per acre (1 gallon to 200 sq. ft.) will be required. More luxuriant foliage will require more spray. If extremely efficient spray equipment is available, it may be possible to obtain thorough coverage with a smaller volume of spray mixture, in which case the concentration of Weed Killer in the spray should be proportionately increased.

The recommended concentration should be doubled for the difficult to control weeds, such as yarrow, goldenrod, poison ivy growing in sunny locations. Deep-rooted perennial weeds such as bindweed, Canada thistle, and leafy spurge are more difficult to destroy and may require repeated applications.

Apply during warm weather when temperature is at least 60° F. and when weeds are growing actively. Apply to lawns between mowings so that a maxi-

CAUTION: KEEP OUT OF REACH OF CHILDREN
CAUTION: KEEP OUT OF REACH OF CHILDREN
DISTRIBUTOR

WEED KILLER

MAR 6, 1967
WATER-SOLUBLE HERBICIDE CONTAINING THE TRIETHANOLAMINE SALT OF 2,4-DICHLOROPHEN OXYACETIC ACID

Kills

Dandelion, Poison Ivy, Poison Oak, Ragweed, Plantain, Thistles, Docks, Wild Morning Glory, Shepherd's Purse, Wild Carrot, Mallow, Mustard, Daisy, and many other broad-leaved weeds without injury to grasses.

Use on golf courses, cemeteries, parks, playgrounds, tennis courts, along highways, railroad rights of way, fire lanes, airfields, ditches, fence rows, factory areas, school grounds, pasture lands, orchards, vineyards and around farm buildings and roadside advertising signs.

ACTIVE INGREDIENT

*Triethanolamine Salt of 2,4-Dichlorophenoxyacetic Acid 10%

INERT INGREDIENTS 90%

Total 100%

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 6%

Licensed under U. S. Pat. 2,322,761 and 2,326,471

CAUTION: May cause skin irritation. Avoid contact with eyes, skin and clothing.

NET CONTENTS: U.S. GALLON
USDA NO.

IMPORTANT

Avoid application to vegetables, flowering plants, shrubbery, trees or other desirable type of plants. This is particularly necessary when treating weeds in orchards, vineyards and nurseries.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. Use a coarse spray, as this type of sprayer is less likely to cause drift. Likewise, do not store near fertilizers, seeds, insecticides or fungicides.

Avoid contamination of water used for irrigation and domestic purposes.

Do not use on bent grass lawns, except for spot spraying.

Always clean spraying equipment thoroughly before using for any other purpose. Rinse several times with hot water containing soap or washing powder (1 lb. to 25 gallons of water) allowing the tank to stand filled with water overnight is recommended. Be sure to clean out hose and spray nozzle.

WARNING: Do not dispose of poison ivy by burning after it has been killed. Smoke from the burning vine is very poisonous on contact with the skin.

Seller makes no warranty of any kind, express or implied, concerning the use of this product.

Buyer assumes all risks of use or handling, whether in accordance with directions or not.

COSTELLO MFG. CO

ESTABLISHED
1859

ST. LOUIS, MISSOURI