

DIRECTIONS FOR USE

Recommend Dilution: 1 oz. per gallon of water.

FOR MOPPING FLOORS:

Increase dilution proportion for unusual conditions. Mop or scrub, wet all surfaces to be disinfected.

FOR USE AS A SPRAY:

On toilet seats, urinals, lavatories, fixtures and all other surfaces to be disinfected. Use recommended dilution (hot water will increase effectiveness). Spray with hand or electric sprayer until surface to be disinfected is entirely wet. Avoid breathing spray mist.

USDA REG. NO. 39-23

SANATROL DISINFECTANT

CONCENTRATED LIQUID DISINFECTANT
GERMICIDE DEODORANT

Recommended for general hospital, industrial and institutional sanitation.

Phenol Coefficient by A.C.A.C. Method at 20°C

- Salmonella Typhosa.....16
- Staphylococcus Aureus.....18

ACTIVE INGREDIENTS

- Sodium-ortho-benzyl-para-chlorophenate.....14.2%
- Isopropanol..... 7.5%
- Dodecyl-benzene-Sulfonate..... 5.0%

INERT INGREDIENTS

- Water..... 73.3%

CAUTION: Keep away from foodstuffs and avoid contact with the skin and eyes.

ANTIDOTE: If accidentally swallowed, give milk, mucilage, corstarch paste, raw egg white or gruel and follow with a tablespoon of mustard in a tumbler of water.

CALL A PHYSICIAN

If undiluted disinfectant cleaner gets on the skin, wash it off with plenty of water.

DIRECTIONS FOR USE

Recommended Dilution: 1 oz per gallon of water.

FOR USE IN SICK ROOM:

Use recommended dilution in washing solution. Wash the bed, furniture, clothes, linens, walls, floor, utensils and the hands of those in attendance.

INDUSTRIAL & INSTITUTIONAL DISINFECTING:

Remove all filth and dirt from the premises and use the recommended dilution for disinfecting floors, walls, fixtures and rest room fixtures.

NET CONTENTS: _____

CAUTION: KEEP OUT OF REACH OF CHILDREN

